Effects of the Cellcultured *Acanthopanax senticosus* Extract on Antioxidative Defense System and Membrane Fluidity in the Liver of Type 2 Diabetes Mouse

Jung-Hee Hong¹, Youn-Soo Cha², and Soon-Jae Rhee^{3,*}

¹The Center for Traditional Microorganism Resources, Keimyung University,
1000 Shndang-Dong, Dalseo-Gu, Daegu 704-701, Korea

²Department of Food Science and Human Nutrition, and Institute for Molecular Biology and Genetics,
Chonbuk National University, Chonju 561-756, Korea

³Department of Food Science and Nutrition, Catholic University of Daegu, Gyung-san-si Gyeongbuk 712-702, Korea

Received 16 December, 2008; Accepted 26 January, 2009

Summary This study examined the effect of cellcultured *Acanthopanax senticosus* (*A. senticosus*) extract on the antioxidative defense system, oxidative stress and cell membrane fluidity in the liver of type 2 diabetes in the C57BL/6J mouse as an animal which is genetically prone to develop insulin resistance and obesity/diabetes. C57BL/6J mice were randomly divided, control diet (N-C), high fat diet (DM-C), control diet plus *A. senticosus* extract (N-CASM), and high fat diet plus *A. senticosus* extract (DM-CASM). The mice were orally administered an *A. senticosus* extract (0.5 g/kg body weight) in the N-CASM and DM-CASM groups once a day for 12 weeks, and distilled water in the N-C and DM-C groups. Cellcultured *A. senticosus* extract was found to be excellent for strengthening the antioxidative defense system, reducing the generation of reactive oxygen species (ROS) and damaging oxidative substances, and maintaining membrane fluidity (MF) in the liver of type 2 diabetes mouse.

Key Words: Cellcultured *Acanthopanax senticosus* extract, C57BL/6J mice, antioxidative defense system, oxidative stress, membrane fluidity

Introduction

As a consequence of the increased prevalence of obesity, type 2 diabetes mellitus (T2D) has become a global epidemic, with more than 300 million individuals worldwide projected to be afflicted with T2D by the year 2025 [1]. Previous studies demonstrated that chronic consumption of a high-fat diet induces obesity, insulin resistance, dyslipidemia, and T2D in genetically C57BL/6J mice [2–4]. Reactive oxygen species (ROS) and the oxidative damage that they cause are thought to play a key role in the pathogenesis of diabetes and its complications [5]. It has been proposed that hyper-

glycemia contributes to the generation in various tissues of free radicals that, in turn, can lead to multiple complications and, at least in part, to the induction of insulin resistance [6, 7]. It has been demonstrated that chronic exposure of cells to high glucose increases intracellular ROS [8–10], and diminished superoxide dismutase (SOD), glutathione peroxidase (GSHpx), catalase (CAT) represent important components of the antioxidant defense enzyme [11] has been found in high-fat diet induced obesity mice.

Acanthopanax senticosus (*A. senticosus*) is a common Asian herb known as “Siberian Ginseng” or “Eleutherococcus senticosus” that is used as an adaptogenic medicine [12]. The major active components of *A. senticosus* are eleutherosides, chiisanosides, isofraxidin, acanthosides, daucosterine, β -sitosterol, sesamine, and savinine [13]. Crude *A. senticosus* extracts have been used as popular health supplements to

*To whom correspondence should be addressed.
Tel: +82-53-850-3523 Fax: +82-53-850-3504
E-mail: jung-hee0322@hanmail.net

treat stress-induced physiological changes [14, 15] as well as antioxidant [16, 17] various allergic conditions [18], inflammation [19], cancer [20] chronic bronchitis, hypertension, ischemic heart disease, and gastric ulcers.

But, crude *A. senticosus* fertility is weaker than other plants, for these reasons, now under development through cell cultivation method.

Therefore, in this study, it is intended to prove that seedlings of *A. senticosus* Max. using cell cultivation method has the anti-diabetes effect in the type II diabetes. The present study observed the effect of cellcultured *A. senticosus* Max. on the antioxidative defense system, oxidative stress and cell membrane fluidity (MF) in the liver of the C57BL/6J mouse which is genetically prone to develop insulin resistance and obesity/diabetes.

Materials and Methods

Materials

All chemicals were obtained from Sigma Chemical Co. (St Louis, MO) unless otherwise indicated. The enzyme assay kits used to determine blood glucose was purchased from Asan Pharm (Asan Pharmaceutical Co., Republic of Korea).

Preparation of *Acanthopanax senticosus* extract

Cultured *A. senticosus* cells with a torpedo shape were supplied by Microplants Co., Ltd (Yusung, Korea). *A. senticosus* cell were allowed to grow in an MS medium (2% sugar) without growth regulators in bio-reactors equipped with an airlift. Culturing for a period of about 10–15 days led the matured somatic embryos to develop plantlets [21]. The *A. senticosus* cells were dried, ground to a fine powder, and extracted with deionized water (thirty times volume) for 9 h at 80°C. The resulting extracts were filtered, concentrated under vacuum at 60°C to 70 Brix.

Animals and diet

Four-week old male C57BL/6J mice were purchased from Jackson Laboratories (Bar Harbor, ME). The animals were maintained on a chow diet (Jeil-jedang, Suwon, Korea) for 1 week, and then randomly divided into 4 groups, control diet (N-C), high fat diet (DM-C), control diet plus *A. senticosus* extract (N-CASM), and high fat diet plus *A. senticosus* extract (DM-CASM). The animals were randomly put into groups ($n = 10$) so that the average weight in each group was comparable. The mice were allowed free access to the diets and water. The mice were orally administered *A. senticosus* extract (0.5 g/kg body weight) in the N-CASM and DM-CASM groups once a day for 12 weeks, and distilled water in the N-C and DM-C groups. Research Diets (New Brunswick, NJ) manufactured the diets. The animals were housed in a temperature controlled environment with a 12 h

light/dark cycle. The food consumption and body weight were measured daily and weekly, respectively.

Sample collection and preparation

All mice were fasted for 12 h prior to sacrificial. Animals were anesthetized using pentothal sodium (40 mg/kg body weight). Their livers were removed immediately, washed in ice-cold saline, and weighted. The removed liver was cut into small pieces and homogenized with a glass Teflon homogenizer in ice-cold 0.1 M potassium phosphate buffer (pH 7.5) equivalent to four times the liver weight. This homogenate was used for the determination of thiobarbituric acid reactive substances (TBARS) contents. The homogenate was centrifuged at 600 g for 10 min, and the resulting supernatant was recentrifuged at 10,000 g for 20 min, in both cases at 4°C. The resulting precipitate, the mitochondrial fraction, was suspended in the above buffer and used as the source of SOD, CAT, superoxide radical, hydrogen peroxide, and carbonyl value and the supernatant was further centrifuged at 105,000 g for 60 min, at 4°C. The supernatant obtained at this point, the cytosolic fraction, was used as the source of the enzymes GSHpx, and hydrogen peroxide and to yield the microsomal fraction used as the source of superoxide radical and carbonyl value. Protein in the liver tissues was determined using the method of Lowry *et al.* [22].

Activities of defense enzymes

The SOD activity was spectrophotometrically measured using a modified version of the method developed by Marklund and Marklund [23]. Briefly, SOD was detected on the basis of its ability to inhibit superoxide mediated reduction. One unit was determined as the amount of enzyme that inhibited the oxidation of pyrogallol by 50%. The activity was expressed as unit/mg protein/min. The GSHpx activity was measured by the method of Lawrence and Burk [24]. The reaction mixture contained 1 mM glutathione reductase in a 0.1 M Tris-HCl (pH 7.2) buffer. The reaction was initiated by adding 2.5 mM H₂O₂ and the absorbance was measured at 340 nm for 1 min. A molar extinction coefficient of 6.22 mM⁻¹ cm⁻¹, was used to determine the activity. The activity was expressed as nmol NADPH/mg protein/min. The CAT activity was measured using Aebi's method [25] a slight modification wherein the hydrogen peroxide decomposition yielding water and oxygen was measured. The mitochondria pellet was dissolved in 1.0 ml of the same homogenization buffer. Ten microliters of the mitochondria solution was added to a cuvette containing 2.89 ml of a 50 mmol/l potassium phosphate buffer (pH 7.4), then the reaction was initiated by adding 0.1 ml of 30 mmol/l H₂O₂ to make a final volume of 3.0 ml at 25°C. The decomposition rate of H₂O₂ was measured at 240 nm for 5 min using a spectrophotometer. A molar extinction coefficient of 0.041 mM⁻¹ cm⁻¹, was used to determine the CAT activity. The activity was

defined as the nmol H₂O₂ decreased/mg protein/min.

Determination of ROS

Superoxide radical content were determined by the Azzi method *et al.* [26]. The reaction mixture contained 50 mM phosphate buffer (pH 7.5), 90 mM succinate, 150 mM KCl, 30 nM KCN, 0.3 mM cytochrome C and mitochondria or microsome added in that sequence. The mixture was shaken vigorously and the absorbance of the mixture was recorded at 550 nm. After incubated at 37°C. for 2 min the absorption was measured again at 550 nm. The liver cytosolic and mitochondrial hydrogen peroxide levels were measured by Wolff's method [27]. Fox 1 (Ferrus Oxidation with Xylenol orange) reagent was prepared as the following mixture with 100 µM xylenol orange, 250 µM ammonium ferrus sulfate, 100 mM sorbitol, and 25 mM H₂SO₄. Fifty microliters of the test sample was added to 950 µL Fox 1 reagent, vortexed, and incubated at room temperature for a minimum of 30 min at which color development is virtually complete. The absorbance was read at 560 nm and the standard was linear in the 0–5 µM H₂O₂ concentration range. The unit was expressed as micromoles of H₂O₂ per mg cytosolic or mitochondrial protein.

Measurement of oxidative damage and age

As a marker of lipid peroxidation, the TBARS concentrations were measured in liver homogenates using method of Park *et al.* [28]. Briefly, 500 µl of a 10%(w/v) tissue homogenate solution was mixed with 2.5 ml of 10% TCA, vortexed, and incubated for 10 min at room temperature. The mixture was centrifuged at 3000 rpm × g for 15 min and the pellet was mixed with 2.5 ml of 0.05 M H₂SO₄, vortexed and recentrifuged at 3000 rpm × g for 15 min. The pellet was heated at 95°C for 30 min after the addition of 2.5 ml of 0.05 M H₂SO₄ and 3 ml of TBA. After cooling the reaction, 3 ml of butanol:pyridine (15:1) solution was added and vortexed. The mixture was centrifuged at 3000 rpm × g for 10 min and the resulting coloured layer was measured at 530 nm using malondialdehyde (MDA) made by the hydrolysis of 1,1,3,3-tetramethoxypropane (TMP) as standard. The content of lipofuscin in the liver tissue was measured in compliance with the Fletcher method *et al.* [29].

Measurement of MF

The quantitative measurement of MF employed the fluorescence polarization technique described by Yu *et al.* [30] with TMA-DPH (1,4-(trimethyl-ammoniumphenyl)-6-phenyl 1-1,3,5-hexatriene) sensing as a fluorescence probe. Membrane preparations (50 µg protein) were suspended in 50 mM Tris-HCl buffer (pH 7.4), mixed with TMA-DPH prepared from a stock solution of 5 mM TMA-DPH solubilized in tetrahydrofurans, and incubated at 37°C for 30 min. Fluorescence polarization was determined using

a Perkin Elmer LS-50 fluorescence spectrophotometer equipped with rotating polarizing filters with samples exposed to a wide range of temperatures (15–40°C). Samples were excited at 360 nm, and emission intensity was read at 435 nm. Calculation of the values for polarization (P) and fluorescence anisotropy (r) of the samples were done by the Yu *et al.* [30].

Statistical analysis

Data from individual experiments are expressed as the mean ± standard deviation. All statistical analysis was performed using SAS software (SAS Institute, Cary, NC). The data were analyzed by two-way analysis of variance (ANOVA) (diet × *A. senticosus* extract). Pairwise statistical significance was established using a *post hoc* Student's *t* test. When significant differences were indicated found, mean values were compared by Tukey's test and repeated-measures ANOVA. Statistical significance is defined as *p* < 0.05.

Results

Body weight, food intake, food efficiency ratio, liver weight and blood glucose level

The high fat diet group gained more weight and faster than the control diet group (Table 1). Although, food intake of the groups were not significantly different, the body weights of the mice fed high fat diets were significantly higher than those of the mice fed control diets. *A. senticosus* extract administration showed no effects on weight in the control diet group; however, in the high fat diet group, the DM-CASM group gained significantly less weight than the DM-C group. The food efficiency ratio (FER) (Table 1) was significantly higher in the high fat diet group than in the control group. *A. senticosus* extract administration significantly decreased the FER, in both control and high fat diet groups. Weight of hepatic, was significantly increased in the DM-C group than that of N group, but DM-CASM group was significantly decreased compared with HF group. The blood glucose level in the DM-C group was 124% higher than in the N-C group. However, the level of in the DM-CASM group was 18% lower than in the DM-C group (Table 1).

Antioxidative defense enzyme activities

Table 2 shows the activity of SOD, an antioxidant enzyme that reduces superoxide radicals to H₂O₂, which in turn is excreted as H₂O based on the activity of GSHpx and CAT, thereby protecting the body from oxygen toxicity. DM-C group had a decreased than the N-C group, whereas, DM-CASM significantly increased than the DM-C group. The GSHpx activity decreased considerably in the DM-C group compared to the N-C group, whereas, the DM-CASM group

Table 1. Effects of the Cellcultured *Acanthopanax Senticosus* extract on body weight gains, food intake, food efficiency ratio and blood glucose in C57BL/6J mice

	Normal		Diabetic		ANOVA ¹		
	N-C	N-CASM	DM-C	DM-CASM	D	A	D × A
Initial body weight (g)	19.52 ± 7.13	19.08 ± 6.91	19.77 ± 9.27	20.23 ± 8.61	NS	NS	NS
Weight gain (g/12 weeks)	5.82 ± 0.78 ^c	5.04 ± 0.36 ^c	16.69 ± 4.27 ^a	12.49 ± 3.45 ^b	0.0001	0.0185	NS
Food intake (g/12 weeks)	198.24 ± 0.06	223.44 ± 0.27	199.08 ± 0.17	194.04 ± 0.07	NS	NS	NS
FER ²	0.029 ± 0.007 ^c	0.023 ± 0.0024 ^d	0.086 ± 0.0065 ^a	0.065 ± 0.0019 ^b	0.0001	0.0001	0.0062
Liver weight (g)	0.85 ± 0.08 ^b	0.88 ± 0.14 ^b	1.54 ± 0.47 ^a	0.97 ± 0.11 ^b	0.0001	0.0163	NS
Blood glucose (mg/dl)	95.98 ± 7.36 ^c	93.49 ± 9.92 ^c	215.56 ± 16.72 ^a	176.92 ± 26.39 ^b	0.0001	0.0007	1

The values represent the mean ± SD of 10 mice per group. Different superscripts in the same row indicate significant differences ($p < 0.05$) among groups by Tukey's test.

¹The degree of significance resulting from the two-way ANOVA is shown with effects of diet (D), administration A. senticosus (A), and the interaction of diet and administration of A. senticosus (D × A) being expressed as the numerical value or as not significant (NS) when $p < 0.05$.

²Food efficiency ratio (FER) was calculated as weight gain weight gain (g/12 weeks)/food intake (g/12 weeks).

Table 2. Effects of the Cellcultured *Acanthopanax Senticosus* extract on SOD, GSHpx, and CAT activity in the liver of type 2 diabetes mouse

		SOD (unit/mg protein/min)	GSHpx (nmol NADPH/mg protein/min)	CAT (nmol/mg protein/min)
Normal	N-C	0.48 ± 0.02 ^a	170.05 ± 4.27 ^a	69.00 ± 5.30 ^a
	N-CASM	0.49 ± 0.03 ^a	179.97 ± 2.20 ^a	75.05 ± 3.89 ^a
Diabetic	DM-C	0.40 ± 0.01 ^c	150.80 ± 5.65 ^b	49.90 ± 5.48 ^b
	DM-CASM	0.45 ± 0.03 ^b	168.74 ± 1.04 ^a	67.90 ± 5.48 ^a

The values represent the mean ± SD of 10 mice per group. Values with different superscript letters are significantly different at $p < 0.05$ as shown by Tukey's test.

was 12% higher than the DM-C group, which remained at the normal level. The CAT activity was 28% lower in the DM-C group, compared to the N-C group, whereas, the DM-CASM group remained at the normal level.

Changes in ROS contents

The liver mitochondria and microsome fraction superoxide radical content (Fig. 1) in the DM-C group were 35% and 49% higher than in the N-C group, respectively. However, the activity in the DM-CASM group were 14% and 25% lower than in the DM-C group, respectively. No significant difference was found in the H₂O₂ content (Fig. 2) in the mitochondria fraction between the experimental groups. But in the case of the cytosol fraction, DM-C group was 19% higher than in the N-C group, whereas, the content in the DM-CASM group was 13% lower than in the DM-C group, which remained at the normal level.

Observation of oxidative damage

The result of TBARS contents as an index oxidative damage was shown in (Table 3). The TBARS concentration

in DM-C group was 49% higher than in the N-C group. However, the concentration in the DM-CASM group was 17% lower than in the DM-C group. The lipofuscin contents (Table 3) in the DM-C group was 38% higher than in the N-C group, whereas, in the DM-CASM group was 14% lower in the DM-C group.

Observation of MF

Since the cell membrane requires good fluidity to maintain homeostasis and metabolism in the body, fluidity is an effective index of adult disease. The cell membrane fluidity in the mitochondria and microsome fractions (Fig. 3) of the liver were 26% and 24% lower in the DM-C group, respectively, compared to the N-C group. However, both fractions of the liver were 17% and 14% higher in the DM-CASM group, respectively, compared to the DM-C group.

Discussion

The current study examined the effects of cellcultured *A. senticosus* extract on the antioxidative defense system,

Fig. 1. Effects of the Cellcultured *Acanthopanax Senticosus* extract on mitochondria (A) and microsomes (B) superoxide radicals in the liver of type 2 diabetes mouse. The values represent the mean $\pm$ SD of 10 mice per group. Values with different superscript letters are significantly different at $p < 0.05$ as shown by Tukey's test.

Fig. 2. Effects of the Cellcultured *Acanthopanax Senticosus* extract on mitochondria (A) and cytosol (B) hydrogen peroxide (H₂O₂) in the liver of type 2 diabetes mouse. The values represent the mean $\pm$ SD of 10 mice per group. Values with different superscript letters are significantly different at $p < 0.05$ as shown by Tukey's test.

Table 3. Effects of the Cellcultured *Acanthopanax Senticosus* extract on TBARS and lipofuscin in the liver of type 2 diabetes mouse

		TBARS (MDA nmol/mg protein)	Lipofuscin (\u00b5g/mg protein)
Normal	N-C	11.80 $\pm$ 0.82 ^c	11.80 $\pm$ 0.82 ^c
	N-CASM	10.97 $\pm$ 1.61 ^c	10.97 $\pm$ 1.61 ^c
Diabetic	DM-C	17.62 $\pm$ 1.54 ^a	17.62 $\pm$ 1.54 ^a
	DM-CASM	14.72 $\pm$ 0.79 ^b	14.72 $\pm$ 0.79 ^b

The values represent the mean $\pm$ SD of 10 mice per group. Values with different superscript letters are significantly different at $p < 0.05$ as shown by Tukey's test.

Fig. 3. Effects of the Cellcultured *Acanthopanax Senticosus* extract on mitochondria (A) and microsome (B) membrane fluidity in the liver of type 2 diabetes mouse. The values represent the mean $\pm$ SD of 10 mice per group. Values with different superscript letters are significantly different at $p < 0.05$ as shown by Tukey's test.

oxidative stress and cell membrane fluidity in the liver of the C57BL/6J mouse as an animal whose auto obesity/diabetes was caused by a high fat diet. These results support previous studies in which a high fat diet increased body weight [31, 32]. There were no significant difference found among the experimental groups, the high fat diet led to significantly higher weight gain compared with the control diet, which resulted in higher FER. Thus mice fed a high fat diet showed a more rapid growth than the mice fed control diet. However, the higher FER exhibited by the *A. senticosus* extract administration, respectively, in control diet and high fat diet group whereas there was no significant effect on body weight in the control diet group. These results suggest that cellcultured *A. senticosus* extract can suppress the increase of weight gain induced by a high fat diet. Also, it was observed that *A. senticosus* extract decreased blood glucose in DM-CASM group. As reported previously by Park *et al.* [33], *A. senticosus* stem bark ethanol extract-treated mice showed marked decreases in plasma glucose level relative to the obese control mice. Oxidative stress means the production stress means the production of highly reactive oxygen radicals that are toxic to the cells, particularly to the cell membrane in which these radicals interact with the lipid bilayer and produce lipid peroxides [34]. Endogenous antioxidant enzymes (e.g., SOD, GSHpx and CAT) are responsible for the detoxication of deleterious oxygen radicals [35, 36]. From the current results of observing the activities of the antioxidative defense system in the liver, the SOD activity in the DM-C group was significantly decreased than that of N-C group, but the DM-CASM group was significantly increased than that of DM-C group. The reason for the decrease in the DM-C group, was suggested that the induction of oxidative stress by high fat diet. Therefore, it is possible that the hydrogen peroxide generated from superoxide radical directly reduces the levels of SOD [37]. GSHpx and CAT activities decreased considerably in the DM-C group compared to the N-C group, respectively.

However, the DM-CASM groups remained the same as that in the N-C group. An increase in GSHpx and CAT activities indicates that *A. senticosus* extract helps in the restoration of vital molecules such as NAD, cytochrome, and glutathione. The inactivity of the antioxidant enzymes GSHpx and CAT in the diabetes groups was attributed to peroxidative damage to the tissues caused by being fed a high fat diet, while the administration of *A. senticosus* extract contributed to maintaining the optimum condition of the cell membrane organelles, essential for enzyme activity, by protecting them from peroxidation. This result confirms that *A. senticosus* acts as an excellent antioxidant even for the liver, which plays an important role in glucose metabolism, and it is a major site of insulin clearance [38]. The phenolic compounds in *A. senticosus*, such as isofraxidin and eleutherosides B and E from the stem bark [39]; eleutheroside E2 and isomaltol 3-*O*- α -D-glucopyranoside from the roots [40]; chiisanoside, chiisano-genin, and hyperin from the leaves [41]; as well as protocatechuic acid, syringin, chorogenic acid, caffeic acid, liriiodendrin, and isofraxidin in ethanol extract of whole *A. senticosus* [42], have a protective effect against oxidative damage. In addition, Lee *et al.* reported that there were no changes in GOT and GPT levels after 6 months treatment with *A. senticosus* [12]. From this data suggested that *A. senticosus* supplementation has no side effects.

ROS was generated as free radicals in the body, such as superoxide anion, hydroxyl radical, and hydrogen peroxide, which are known to induce many adult diseases and promote the aging process. For example, Laura's report [43] stated that ROS induce Alzheimer's disease. Recent evidence showed that increased flux of free fatty acid (FFA), glucose, or hexosamine could raise mitochondrial ROS production, leading to increased intracellular oxidative stress [44, 45]. Among the free radicals which are known to induce aging or adult diseases, the superoxide radical content in the mitochondria and microsome fractions of the liver was

found to be considerably higher in the DM-C group compared to the N-C group, which coincides with a previous report of increased ROS in diabetes [46]. However, in the both fractions, the activity in the DM-CASM group was significantly decreased compared to the DM-C group, respectively. In the case of cytosol fraction, showed that the hydrogen peroxide concentration was increased in DM-C group compared to the N-C group, but the DM-CASM group was significantly decreased compared to the DM-C group.

Because type 2 diabetes is a chronic disease, prevention of oxidative stress needs to be achieved not only periodically but as long as the hyperglycemia persists. Lipid peroxidation is recognized as the most important substance which damaged by various toxicity compounds and drugs or pathophysiological phenomenon by disease. This is caused by the augmentation of oxidative stress and the decrease of an antioxidative defense system *in vivo*. The contents of TBARS in DM-C group was significantly higher than in the N-C group. However, the concentration of the DM-CASM group was significantly lower than that of the DM-C group. The observed effect of *A. senticosus* extract on lipofuscin that is consumptiveness aging pigment produced by free radicals that estimate the degree of aging and leads to diseases in adult people by toxicity function in the cell and promotes aging. The lipofuscin contents in the DM-C group was higher than in the N-C group whereas, in the DM-CASM group was significantly lower than in the DM-C group. In the current study, when the fluidity in the cell membrane of the liver was observed, mitochondria and microsome fractions both showed a decrease in the DM-C group compared to the N-C groups. It would appear that the cell membranes connected by double bonds of polyunsaturated fatty acid in the diabetes-induced groups lost fluidity due to the generation of ROS, such as superoxide radical and hydrogen peroxide. whereas the activity of antioxidant enzyme increased with the administration of *A. senticosus* extract, resulting in significantly increased cell membrane fluidity in the DM-CASM group compared to the DM-C group.

In conclusion, *A. senticosus* extract strengthened the antioxidative defense system with an increased activity of antioxidant enzymes, such as SOD, GSHpx and CAT, in the liver of the type 2 obesity/diabetes experimental mice. Accordingly, *A. senticosus* extract was found to reduce the accumulation of ROS, such as superoxide radical and H₂O₂, which decrease the generation of oxidative damage substances, such as TBARS and lipofuscin, increase the membrane fluidity lowered by oxidative damage.

Acknowledgments

This work was supported by the Ministry of Health & Welfare Grant (03-PJ1-PG10-22000-0015).

Abbreviations

CAT, catalase; FER, food efficiency ratio; GSHpx, glutathione-peroxidase; MDA, malondialdehyde; MF, membrane fluidity; NIDDM, non-insulin dependent diabetes mellitus; ROS, reactive oxygen species; SOD, superoxide dismutase; TBARS, thiobarbituric acid reactive substances.

References

- [1] Wei-jian, J.: Sirtuins: novel targets for metabolic disease in drug development. *Biochem. Biophys. Res. Commun.*, **373**, 341–344, 2008.
- [2] Wouters, K., van Gorp, P.J., Bieghs, V., Gijbels, M.J., Duimel, H., Lutjohann, D., Kerksiek, A., van Kruchten, R., Maeda, N., Staels, B., van Bilsen, M., Shiri-Sverdlov, R., and Hofker, M.H.: Dietary cholesterol, rather than liver steatosis, leads to hepatic inflammation in hyperlipidemic mouse models of nonalcoholic steatohepatitis. *Hepatology*, **48**, 474–486, 2008.
- [3] Stewart, L.K., Soileau, J.L., Ribnicky, D., Wang, Z.Q., Raskin, I., Poulev, A., Majewski, M., Cefalu, W.T., and Gettys, T.W.: Quercetin transiently increases energy expenditure but persistently decreases circulating markers of inflammation in C57BL/6J mice fed a high-fat diet. *Metabolism*, **57**, S39–S46, 2008.
- [4] Shin, C.C., Lin, C.H., and Lin, W.L.: Effects of *Momordica charantia* on insulin resistance and visceral obesity in mice on high-fat diet. *Diabetes Res. Clin. Pract.*, **81**, 134–143, 2008.
- [5] Fiala, E., Westendorf, J., and West, I.C.: Radicals and oxidative stress in diabetes. *Toxicology*, **146**, 83–92, 2000.
- [6] Edelstein, D., Dimmeler, S., Ju, Q., Sui, C., and Brownlee, M.: Biochemistry and molecular cell biology of diabetic complications. *J. Clin. Invest.*, **108**, 1341–1348, 2001.
- [7] Marfella, R., Quagliaro, L., Nappo, F., Ceriello, A., and Giugliano, D.: Acute hyperglycemia induces an oxidative stress in healthy subjects. *J. Clin. Invest.*, **108**, 635–636, 2001.
- [8] Ha, H. and Lee, H.B.: Reactive oxygen species as glucose signaling molecules in mesangial cells cultured and under high glucose. *Kidney Int.*, **77**, S19–S25, 2000.
- [9] Park, S.H., Choi, H.J., Lee, J.H., Woo, C.H., Kim, J.H., and Han, H.J.: High glucose inhibits renal proximal tubule cell proliferation and involves PCK, oxidative stress, and TGF-beta 1. *Kidney Int.*, **59**, 1695–1705, 2001.
- [10] Peiro, C., Lafuente, N., Matesanz, N., Llergo, J.L., Vallejo, S., Rodriguez-Manas, L., and Sanchez-Ferrer, C.F.: High glucose induces cell death of cultured human aortic smooth muscle cells through the formation of hydrogen peroxide. *Br. J. Pharmacol.*, **133**, 967–974, 2001.
- [11] Lee, Y.M., Choi, J.S., Kim, M.H., Jung, M.H., Lee, Y.S., and Song, J.: Effects of dietary genistein on hepatic lipid metabolism and mitochondrial function in mice fed high-fat diets. *Nutrition*, **22**, 956–964, 2006.
- [12] Lee, Y.J., Chung, H.Y., Kwak, H.K., and Yoon, S.: The

- effects of *A. senticosus* supplementation on serum lipid profiles, biomarkers of oxidative stress, and lymphocyte DNA damage in postmenopausal women. *Biochem. Biophys. Res. Commun.*, **375**, 44–48, 2008.
- [13] Davydov, M. and Krikorian, A.D.: Eleutherococcus senticosus (Rupr. & Maxim.) Maxim. (Araliaceae) as an adaptogen: a cross look. *J. Ethnopharmacol.*, **72**, 345–393, 2000.
- [14] Fujikawa, T., Yamauchi, A., Morita, L., Takeda, H., and Nishibe, S.: Protective effects of *Acanthopanax senticosus* Harms from Hokkaido and its components on gastric ulcer in restrained cold water stressed rats. *Biol. Pharm. Bull.*, **19**, 1227–1230, 1996.
- [15] Gaffney, B.T., Hugel, H.M., and Rich, P.A.: Panax ginseng and Eleutherococcus senticosus may exaggerate an already existing biphasic response to stress via inhibition of enzymes which limit the binding of stress hormones to their receptors. *Med. Hypotheses*, **56**, 567–572, 2001.
- [16] Lee, S.H., Son, D.W., Ryu, J.Y., Lee, Y.S., Jung, S.H., Kang, J.G., Lee, S.Y., Ki, H.S., and Shin, K.H.: Anti-oxidant activities of *Acanthopanax senticosus* stems and their ligand components. *Arch. Pharm. Res.*, **27**, 106–110, 2004.
- [17] Lee, Y.S., Jung, S.H., Lim, S.S., Ji, J., Lee, S., and Shin, K.H.: Effects of the water extract from the stem bark of on hyperlipidemia in rats. *Korean J. Pharm.*, **32**, 103–107, 2001.
- [18] Yi, J.M., Hong, S.H., Kim, H.K., Song, H.J., and Kim, H.M.: Effect of *Acanthopanax senticosus* stem on mast cell-dependent anaphylaxis. *J. Ethnopharmacol.*, **79**, 347–352, 2002.
- [19] Lin, Q.Y., Jin, L.J., Cao, Z.H., and Xu, Y.P.: Inhibition of inducible nitric oxide synthase by *Acanthopanax senticosus* extract in RAW264.7 macrophage. *J. Ethnopharmacol.*, **23**, 231–236, 2008.
- [20] Yamazaki, T., Shimosaka, S., Sasaki, H., Matsumura, T., Tukiya, T., and Tokiwa, T.: (+)-Syringaresinol-di-O-beta-D-glucoside, a phenolic compound from *Acanthopanax senticosus* Harms, suppresses proinflammatory mediators in SE982 human synovial sarcoma cells by inhibiting activating protein-1 and/or nuclear factor-kappaB activities. *Toxicol. In Vitro*, **21**, 1530–1537, 2007.
- [21] Choi, Y.E., Kim, J.W., and Yoon, E.S.: High frequency of plant production via somatic embryogenesis from callus or cell suspension cultures in *Eleutherococcus senticosus*. *Ann. Bot (Lond.)*, **83**, 309–314, 1999.
- [22] Lowry, O.H., Roseborough, N.J., Farr, A.L., and Randall, R.J.: Protein measurement with the folin phenol reagent. *J. Biol. Chem.*, **193**, 265–275, 1951.
- [23] Marklund, S. and Marklund, G.: Involvement of the superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *Eur. J. Biochem.*, **47**, 469–474, 1974.
- [24] Lawrence, R.A. and Burk, R.F.: Glutathione peroxidase activity in selenium deficient rat liver. *Biochem. Biophys. Res. Commun.*, **71**, 952–958, 1976.
- [25] Aebi, H., Wyss, S.R., Scherz, B., and Skvaril, F.: Heterogeneity of erythrocyte catalase. Isolation and characterization of normal and variant erythrocyte catalase and their subunits. *Eur. J. Biochem.*, **48**, 137–145, 1974.
- [26] Azzi, A., Montecucco, C., and Richter, C.: The use of acetylated ferric cytochrome c for the detection of superoxide radicals produced in biological membranes. *Biochem. Biophys. Res. Commun.*, **65**, 597–603, 1975.
- [27] Wolff, S.P.: Ferrous ion oxidation in presence of ferric ion indicator xylenol orange for measurement of hydroperoxides. *Methods Enzymol.*, **223**, 182–189, 1994.
- [28] Park, S.Y., Bok, S.H., Jeon, S.M., Park, Y.B., Lee, S.J., Jeong, T.S., and Choi, M.S.: Effect of rutin and tannic acid supplements on cholesterol metabolism in rats. *Nutr. Res.*, **22**, 283–295, 2002.
- [29] Fletcher, B.L., Dillard, C.J., and Tappel, A.L.: Measurement of fluorescent lipid peroxidation products in biological systems and tissues. *Anal. Biochem.*, **52**, 1–9, 1973.
- [30] Choi, J.H. and Yu, B.P.: Unsuitability of TBA test as a lipid peroxidation maker due to prostaglandin synthesis in the aging kidney. *Age*, **13**, 61–64, 1990.
- [31] Cha, Y.S., Soh, J.R., and Kim, J.W.: *Acanthopanax senticosus* extract prepared from cultured cells improves lipid parameters in rats fed with a high fat diet. *J. Korean Soc. Nutraceut. Food*, **8**, 40–45, 2003.
- [32] Lin, S., Thomas, T.C., Storlien, L.H., and Huang, X.F.: Development of high fat diet-induced obesity and leptin resistance in C57BL/6J mice. *Int. J. Obes. Relat. Metab. Disord.*, **24**, 639–646, 2000.
- [33] Park, S.H., Lee, S.G., Kang, S.K., and Chung, S.H.: *Acanthopanax senticosus* reverses fatty liver disease and hyperglycemia in ob/ob mice. *Arch. Pharm. Res.*, **29**, 768–776, 2006.
- [34] Sakaguchi, S. and Furusawa, S.: Oxidative stress and septic shock: metabolic aspects of oxygen-derived free radicals generated in the liver during endotoxemia. *FEMS. Immunol. Med. Microbiol.*, **47**, 166–177, 2006.
- [35] Del Maestro, R.F.: An approach to free radicals in medicine and biology. *Acta Physiol. Scand. Suppl.*, **492**, 153–168, 1980.
- [36] Stern, L.Z., Ringel, S.P., Ziter, F.A., Menander-Huber, K.B., Ionasescu, V., Pellegrino, R.J., and Snyder, R.D.: Drug trial of superoxide dismutase in Duchenne's muscular dystrophy. *Arch. Neurol.*, **39**, 34234–34236, 1982.
- [37] Hodgson, E.K. and Fridovich, I.: The interaction of bovine erythrocyte superoxide dismutase with hydrogen peroxide: chemiluminescence and peroxidation. *Biochemistry*, **14**, 5299–5303, 1975.
- [38] Duckworth, W.C., Hamel, F.G., and Peavy, D.E.: Hepatic metabolism of insulin. *Am. J. Med.*, **85**, 71–76, 1988.
- [39] Nishibe, S., Kinoshita, H., Takeda, H., and Okano, G.: Phenolic compounds from stem bark of *Acanthopanax senticosus* and their pharmacological effect in chronic swimming stress rats. *Chem. Pharm. Bull. (Tokyo)*, **38**, 1763–1765, 1990.
- [40] Li, Q., Jia, Y., Xu, L., Wang, X., Shen, Z., Liu, Y., and Bi, K.: A new lignan glycoside from *Eleutherococcus senticosus*. *Planta Med.*, **67**, 776–778, 2001.
- [41] Lee, S., Shin, D.S., Oh, K.B., and Shin, K.H.: Antibacterial compounds from the leaves of *Acanthopanax senticosus*. *Arch. Pharm. Res.*, **26**, 106–110, 2003.

- [42] Li, Q., Jia, Y., Xu, L., Wang, X., Shen, Z., Liu, Y., and Bi, K.: Simultaneous determination of protocatechuic acid, syringin, chlorogenic acid, caffeic acid, liriiodendrin and isofraxidin in *Acanthopanax senticosus* Harms by HPLC-DAD. *Biol. Pharm. Bull.*, **29**, 532–534, 2006.
- [43] Laura, J.H., Michael, A.T., and Juan, C.T.: Increased susceptibility of Alzheimer's disease temporal cortex to oxygen free radical-mediated process. *Free Radic. Biol. Med.*, **23**, 183–190, 1997.
- [44] Brownlee, M.: Biochemistry and molecular cell biology of diabetic complications. *Nature*, **414**, 813–820, 2001.
- [45] Nishikawa, T., Edelstein, D., Du, K.L., Yamagishi, S., Matsumura, T., Kaneda, Y., Yorek, M.A., Beebe, D., Oates, P.J., Hammes, H.P., Giardino, I., and Brownlee, M.: Normalizing mitochondrial superoxide production blocks three pathways of hyperglycaemic damage. *Nature*, **404**, 787–790, 2000.
- [46] Kwag, O.G., Kim, S.O., Choi, J.H., Rhee, I.K., Choi, M.S., and Rhee, S.J.: Vitamin E improves microsomal phospholipase A₂ activity and the arachidonic acid cascade in kidney of diabetic rats. *J. Nutr.*, **131**, 1297–1301, 2001.