Oxidative Stress and Digestive Diseases

Guest Editor: Hidekazu Suzuki

Introduction to serial reviews: Oxidative stress and digestive diseases

Hidekazu Suzuki*

Fellowship Training Center and Medical Education Center, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan

(Received 21 November, 2017; Accepted 22 November, 2017; Published online 11 April, 2018)

The digestive tract is always in contact with the outside world and is an organ that is constantly exposed to foreign antigens because of feeding behavior⁽¹⁾ and continuous exposure to microorganisms.^(2,3) Furthermore, because the liver is a large-scale metabolic factory of ingested nutrients and drugs, it is also easily exposed to oxidative stress because extensive oxygen is always used in the process of metabolism.^(4,5) Oxidative stress in the pathogenesis of digestive diseases is an important theme that has been discussed for a long time.^(6,7) Inflammation and carcinogenesis, which are dual characteristics of the pathology of the digestive organs, also always involve this concept of oxidative

stress. The aim of this serial review was to promote researchers' and clinicians' understanding of the oxidative aspects of the pathogenesis of digestive diseases and to foster research activities in the fields of gastroenterology and hepatology by cultivating the novel approaches or ideas to the oxidative stress-related disease process. In this serial review, I invite several outstanding researchers in this field to summarize their work and to review their peers' scientific work, and to organize this knowledge according to their opinions. I thank all the anonymous and volunteer reviewers of these articles for their excellent insights.

References

- 1 Han JY, Miura S, Akiba Y, *et al.* Chronic ethanol consumption exacerbates microcirculatory damage in rat mesentery after reperfusion. *Am J Physiol Gastrointest Liver Physiol* 2001; **280**: G939–G948.
- 2 Suzuki H, Mori M, Seto K, *et al.* *Helicobacter pylori*-associated gastric pro- and antioxidant formation in Mongolian gerbils. *Free Radic Biol Med* 1999; **26**: 679–684.
- 3 Nishizawa T, Suzuki H. Gastric carcinogenesis and underlying molecular mechanisms: *Helicobacter pylori* and novel targeted therapy. *Biomed Res Int* 2015; **2015**: 794378.
- 4 Suematsu M, Suzuki H, Ishii H, *et al.* Early midzonal oxidative stress preceding cell death in hypoperfused rat liver. *Gastroenterology* 1992; **103**: 994–1001.
- 5 Suematsu M, Kato S, Ishii H, *et al.* Intralobular heterogeneity of carbon tetrachloride-induced oxidative stress in perfused rat liver visualized by digital imaging fluorescence microscopy. *Lab Invest* 1991; **64**: 167–173.
- 6 Suzuki H, Nishizawa T, Tsugawa H, Mogami S, Hibi T. Roles of oxidative stress in stomach disorders. *J Clin Biochem Nutr* 2012; **50**: 35–39.
- 7 Suzuki H, Matsuzaki J, Hibi T. Ghrelin and oxidative stress in gastrointestinal tract. *J Clin Biochem Nutr* 2011; **48**: 122–125.

*To whom correspondence should be addressed.
E-mail: hsuzuki.a6@keio.jp