	stenosis of the lesion related to MI at the initial study				
	≤25%	50%	75%	90%	99%
Group I(n=16)	7	4	4	1	0
Group II(n=6)	3	2	0	1	0
Group III(n=6)	2	2	1	1	0
Group IV(n=1)	0	1	0	0	0

These results suggest that 1) myocardial infarction is associated with sudden progression from the lesion with less than 75% stenosis in the extent and severity of coronary obstructive lesions and 2) there is no significant relationship between patterns of onset in MI and degree of stenosis in the lesions evaluated at the initial CAG.

-33-

CLINICAL PROFILES OF ACUTE PHASE IN MYOCARDIAL INFARCTION WITH NO SIGNIFICANT STENOSIS DURING CHRONIC PHASE

Keigo Dote, Takashi Sueda, Toshiaki Uchida
Hironobu Tateishi, Hikaru Satoh and
Kohshiroh Sasaki

Hiroshima City Hospital CCU

To describe how coronary artery occlusion can resolve with time and to assess the effect of spontaneous recanalization of infarct vessel on LV function, we analyzed 41 cases of myocardial infarction with no significant stenosis (≥50%) during chronic angiogram (NSMI). NSMI were divided into 2 groups according to acute angiographic evidence; patients who had stenosis exceeding 90% (25 cases Group 1) and who had 75% or less (16 cases Group 2). The factor related to regression of stenosis were thrombolysis with PTCR in G1 and relief of spasm with intracoronary NTG in G2. Acute LVEF were similarly low in both group. However, chronic EF were significantly lower and max CPK were significantly higher in G1 than G2 (acute EF G1:56±14, G2: 61±15 NS, chronic EF G1:55±13, G2:78±8 P<0.01, max CPK G1:3696±2158, G2:495±422 P<0.01). Moreover, G2 were divided into 2 groups according to acute EF; patients with or without asynergy. Both subgroups revealed normal chronic EF and slightly elevation of CPK. These data indicate that the difference of main factor related to coronary artery occlusion-thrombus or spasm-is an important determinant of myocardial injury by ischemia-irreversible necrosis or stunning.

-34-

AN IMMUNORADIOMETRIC ASSAY OF SERUM CARDIAC MYOSIN LIGHT CHAIN I -A NEW METHOD OF MYOCARDIAL INFARCT SIZING-

Yoshio Yazaki, Mitsuaki Isobe, Ryoza Nagai
Kazuhide Yamaaki, Fumimaro Takaku

The Third Department of Internal Medicine,
University of Tokyo

We have reported that serum cardiac myosin light chain I (LCI) analysis by radioimmunoassay (RIA) could provide an accurate method for sizing myocardial infarction. Furthermore, the determination of serum LCI could evaluate quantitatively the effect of intracoronary thrombolysis on the extent of myocardial damage in the patients with acute myocardial infarction, since the liberation of LCI from damaged myocardium into the serum did not appear to be affected by coronary reperfusion. However, in spite of its clinical usefulness, RIA of LCI in serum could not be applied as a generally used method for infarct sizing, because of the limitation of the antibody preparation for the assay. Recently, immunoradiometric assay (IRMA) has been applied for the determination of protein concentrations with special reference to its specificity, short reaction time and large scaled

productivity with relative ease. Thus, we have developed a sensitive IRMA of serum LCI using two types of monoclonal antibodies (MLM508, MLM544) specific for LCI. In this assay, 100 µl of serum was added with 200 µl of 125I-labeled MLM508 (5x 10⁴cpm) to the tubes coated with MLM544 and incubated for 2 hours. Amount of bound LCI in serum was quantitated by counting the tubes after washing. One ng/ml of serum LCI could be measured by this method. We determined serially serum LCI by this IRMA in 75 patients with acute myocardial infarction within 12 hours after the onset of chest pain. A good correlation was observed between serum LCI concentrations determined by IRMA and RIA (r=0.92). Again, we observed that peak LCI levels correlated with left ventricular ejection fraction (r=0.79). Thus, an immunoradiometric assay of serum LCI can provide a clinically useful and accurate method for myocardial infarct sizing even after intracoronary thrombolysis.

-35-

CORRELATION BETWEEN ANATOMIC MYOCARDIAL INFARCT SIZE, TOTAL CPK RELEASE AND A QRS SCORING SYSTEM IN 12-LEADS ECG

Hironori Ezaki, Satoru Matsushita,
Genji Toda, Shuji Oda, Akinori Hattori, Makoto Sakai, Shinichirou Ohkawa, Keiji Ueda and
Kizuku Kuramoto.

Department of Cardiology, Tokyo Metropolitan Geriatric Hospital

Myocardial infarct size is definitely related to cardiac function and prognosis. For critical evaluation of the infarct size estimation, we weighed infarcted myocardium in 44 autopsy cases (24 men and 20 women, mean age of 76.8±8.7 y.o.), and compared with peak serum CPK activity (peak-CPK), peak serum CPKMB isoenzyme activity (peak-CPKMB), total CPK release (ΣCPK), and a QRS scoring system in 12-leads ECG modified by Wagner et al., respectively. Mean infarcted myocardial weight (MI weight) of 44 cases were 38.4±19.1g. Mean value of peak-CPK, peak-CPKMB and ΣCPK were 2487±2344, 221±208, 4597±3750 IU/ml, respectively, and mean QRS score was 7.2±3.4 points. Interval between serial CPK determination, ECG recording and autopsy averaged 130.1 and 52.4 days, respectively. There were significant correlation (p<0.01) between MI weight and peak-CPK (r=0.63, n=17), peak-CPKMB (r=0.79, n=17), ΣCPK (r=0.72, n=11), and a QRS scoring system (r=0.64, n=39), respectively. Especially in cases of non-transmural myocardial infarction, a QRS scoring system showed good correlation with MI weight (r=0.82, n=11, p<0.01).

We concluded that peak-CPK, peak-CPKMB, ΣCPK, and a QRS scoring system were useful for the estimation of myocardial infarct size.

-36-

COMPARATIVE STUDY OF RECONSTRUCTION AND ECG FINDINGS OF INFARCT HEART AUTOPSY CASES

-FIRST REPORT: Q WAVES AND INFARCT SIZES-

Tomohiko Iwasaki, Noboru Kaneko, Koshichiro Hirosawa, Makoto Takeshi*, Kunihiro Teraoka*,