-217-

CLINICAL APPLICATION OF AMBULATORY HOLTER TAPE RECORDING OF ATRIAL ELECTROGRAM TO THE PATIENTS WITH WPW SYNDROME

Takuro Misaki, Keiichi Mukai, Yutaka Kuzawa
Shigeho Iida, Eiichirou Kamata
Makoto Tsubota, Makoto Shinagawa,
Takeshi Watanabe, Takashi Iwa

Department of Surgery Kanazawa University
School of Medicine

Detection of the P wave during tachyarrhythmia is often difficult in the patients with WPW syndrome when we use conventional ambulatory Holter ECG tape recording. To obtain sharp and stable atrial amplitude recording, a J shaped bipolar catheter was introduced percutaneously and positioned in right atrial appendage. The condition of the filter of second channel was changed into new setting: frequencies of 100Hz with time constant of 0.002 sec. A surface ECG was recorded simultaneously on first channel.

In all 15 patients with WPW syndrome, sharp and clear atrial electrograms were obtained. The atrial rate (AA or FF interval) could be counted automatically. Spontaneous tachyarrhythmias were seen in all during the recording period in normal active life. Tachyarrhythmias could be easily diagnosed: 9 as reentrant tachycardia, 2 as atrial flutter with wide QRS tachycardia, 4 as atrial fibrillation with pseudo VT. The mechanism of initiation and termination of tachyarrhythmia could be identified easily. Premature atrial contraction was found to be much more common to initiate reentrant tachycardia even when premature ventricular contraction was strongly suspected by a surface ECG. It could be easily differentiated AV block from VA block at the time of termination. There were no complications at all.

In conclusion: Ambulatory Holter tape recording of atrial electrogram in addition to conventional ECG recording is a useful method to know not only the diagnosis but also the mechanism of spontaneous tachyarrhythmias in the patients with WPW syndrome. With our newly designed Holter recorder, sharp and stable atrial amplitude could be recorded even on exercise.

-218-

CRYOSURGERY FOR CARDIAC ARRHYTHMIAS

Shigeho Iida, Takuro Misaki,
Keiichi Mukai, Eiichiro Kamata,
Michio Kawasuji, Go Watanabe,
Takashi Iwa

Department of Surgery (1), Kanazawa University
School of Medicine

Cryosurgery was introduced for the surgical treatment of tachyarrhythmias in 1977. In our institute, this technique has been used since 1980 with very good results. In this paper, we will demonstrate our clinical study of cryosurgery of the heart.

Cryothermic technique was used in 140 patients. These include 118 pts of WPW syndrome, 19 of ventricular tachycardia, 2 of rapid left atrial rhythm and one of atrial flutter.

WPW syndrome; A total of 254 pts of WPW syndrome have been operated on since 1973. Cryothermic technique was used in 118 pts (46%), aged 7 to 68 years, classified into left cardiac type 65, right 20, septal 11, multiple 11 and concealed 11. The cryoprobe was applied on the ventricular wall after an endocardial atrial separation from the annulus being performed with a scalpel. With this new method, the accessory pathways

were successfully divided in 117 of 118 pts (99.2%), while 118 of 136 pts (86.8%) with our conventional method.

VT; Twenty-three pts with sustained VT underwent surgical treatment. They were 21 with nonischemic and 2 with ischemic heart disease. Cryosurgery was used in 19 pts. The main surgical method for nonischemic right ventricular type was excision of the earliest excitation area and cryosurgery around the excised area. For nonischemic left ventricular type incision and cryosurgery of the LV apex was performed. One pt of ischemic type had endocardial cryosurgery in addition to aneurysmectomy and AC bypass. Seventeen out of 19 pts were cured. Fifteen pts became completely free from medication, while 2 pts need medication. Two pts died from recurrence of VT.

In conclusion, cryosurgery offers considerable advantages in the surgical treatment of cardiac arrhythmias.

Chairmen

219-227 A. Suzuki, Tokyo
T. Ban, Kyoto

-219-

SURGERY FOR THE VENTRICULAR SEPTAL DEFECT FOLLOWING MYOCARDIAL INFARCTION

Hirotaka MURATA, Yoshizumi OKA, Keiichi AO-
KI, Shinsho MAEDA, Katsuhiko YAMASHITA, Koji KITAI
Hiroshi TERAII, Hideo SOH, Makoto TANAKA, Hideki
YAO, Takashi YASUOKA, Takashi MIYAMOTO

Department of Thoracic Surgery, Hyogo College of Medicine

Ventricular septal perforation (VSP) is an uncommon but catastrophic complication of acute myocardial infarction (AMI). Since 1963 coronary artery surgery was operated on in 125 patients (pts) at our institute. Among these, 8 pts with post infarction VSP were operated upon within 17 days after AMI (3 female & 5 male pts with an average of 66 years). Septal perforation was found on the average 4 days after AMI (anterior 4, apical 3 & posterior 1). Operation was undertaken with single patch method through the anterolateral aneurysmal incision in 3 & through the posterior left ventricular infarct in 1. Other 3 pts were repaired with amputation of apical portion. Associated coronary bypass grafting was done in 1 pt. Operative mortality was 50%. We concluded that once diagnosed, urgent operation is recommended before deterioration occurs.