Hidemi Shinomiya, Yoshio Atagi, Kazuto Hamai,
Naoki Nishimura, Hiroyoshi Mori,

The Second Department of Internal Medicine, School
of Medicine, The University of Tokushima.

To evaluate the endocardial mappings of Torsade de pointes (TdP), we attempted the induction of TdP by ventricular pacing in 23 anesthetized dogs. Quinidine (30 mg/kg, i. v.) was given and premature or rapid stimulation of the ventricle was used to induce TdP. When TdP was not induced by the ventricular stimulation, the left anterior descending coronary artery (LAD) was occluded and the same procedure of ventricular pacing was performed. Endocardial mappings in left ventricle were made from 21 electrograms recorded simultaneously by bipolar electrodes. 15 ventricular tachycardias in 7 dogs met criteria for the diagnosis of TdP. At each 6 sites in left and 3 sites in right ventricle, ERP after administration of quinidine was significantly longer than that before quinidine. Δ ERP(max.ERP-min.ERP) increased significantly after administration of quinidine. The characteristics of endocardial mappings during TdP were as follows. 1) The wave front of excitement spread radially from the earliest endocardial activation site in the left ventricle (LV). When the earliest endocardial activation sites in LV were similar in some mappings, these conduction velocities differed. 2) With the change in QRS axis, the earliest endocardial activation site in LV moved from one site to the another. These findings suggested that increased Δ ERP was necessary for the induction of TdP and a reentry was considered to be a possible mechanism to TdP.

-373-

THE QT INTERVAL MAPPING IN THE NORMAL SUBJECTS AND PATIENTS WITH PRIOR MYOCARDIAL INFARCTION

Michiyasu Yamaki, Isao Kubota, Kozue Ikeda, Hide Igarashi, Akira Igarashi, Sukehiko Kawashima, Seiji Yasumura, Tsuyoshi Konta, Kanji Hanashima, Kazuharu Nakamura, Kai Tsuiki and Shoji Yasui

The First Department of Internal
Medicine, Yamagata University, School of
Medicine

The QT interval is a clinically important electrocardiographic measurement. In this study, we attempted to determine whether the QT interval had spatially a narrow range of values on a torso, and whether these spatial pattern of the QT interval were altered by regional abnormalities of the left ventricle. Then, we studied 20 normal subjects and 40 patients with prior myocardial infarction (20 anterior, 20 infero-posterior). Body surface mapping was performed using a HPM 5100 system from 87 torso electrodes. In each patient, the QT interval was determined for each lead by a computer-assisted automated-manual method reported by Mirvis. The QT interval had spatially a wide range of values on a torso. The differences between the longest and shortest interval were 64 ± 16 msec (mean $\pm$ SD) in normal subjects, 96 ± 20 msec in anterior MI group and 76 ± 22 msec in infero-posterior MI group. In the normal subjects, long QT interval were located over the left anterior chest, and short QT interval were positioned over the lower right chest. In patients with prior MI, these spatial distribution were altered. long QT interval were found on the upper anterior chest in the anterior MI group associated with the site of abnormal Q wave. In the infero-posterior MI group, long QT interval were located over the lower left lateral and back. Thus, the body surface distribution of the QT interval had characteristic pattern in the normal and prior MI patients.

-374-

EFFECTS OF CALCIUM ANTAGONISTS ON VENTRICULAR ARRHYTHMIAS AND ELECTRICAL ALTERNANS OF ST SEGMENT DURING ACUTE CORONARY OCCLUSION IN DOGS

Daiju Morishita, Takayuki Ikeda,
Toshisuke Terakawa, Akira Higashino,
Satoshi Hirota, Tetsuya Nishida,
Koji Kubota, Shigeo Takata,
Nobu Hattori

The 1st Department of Internal Medicine
School of Medicine, Kanazawa University

Calcium antagonists have been reported to suppress ventricular arrhythmias (VA) during acute ischemia. We, therefore, studied the effects of calcium antagonists (nifedipine, verapamil and diltiazem) on the ventricular arrhythmias and electrical alternans of ST(STA) during acute myocardial ischemia. We recorded 48 unipolar electrograms simultaneously from the epicardial surface of the ventricle in 56 anesthetized dogs during 30 min after occlusion of the left anterior descending coronary artery.

In 31 dogs calcium antagonists were administered intravenously 20 min prior to coronary occlusion, and followed by continuous administration over 60 min. STA was observed in all control dogs, and VA occurred in 12 of 25 dogs during coronary occlusion. And there was significant difference ($p < 0.01$) in the size of the area recorded STA 15 min after coronary occlusion between VA group (12 dogs) and no VA group (13 dogs). The frequency of VA during coronary occlusion was reduced by the administration of calcium antagonists markedly attenuated the degree of STA and reduced the size of the area recorded STA during coronary occlusion. STA was appeared in the ischemic and the border zone within 2 min after coronary occlusion in control dogs, but was not observed even after the longer period of occlusion by the administration of each of calcium antagonists. There were no significant differences of the effects to the appearance of STA and the frequency of VA by each of calcium antagonists.

In conclusion, calcium antagonists may attenuate the degree of STA and the size of the area recorded STA, and prevent the occurrence of VA.

-375-

VECTORCARDIOGRAPHIC FINDING OF RIGHT VENTRICULAR HYPERTROPHY IN 1 VESSEL CORONARY ARTERY DISEASES

Masakazu Motomura, Yoshihiro Kawaguchi,
Iwao Mashiro, Yoshimichi Kudo,
Satoru Sakoda, Kenichi Mitsunami,
Masahiko Kinoshita, Seiichi Kawakita

The 1st Department of Internal Medicine,
Shiga University of Medical Science

We aimed in this study to investigate the modification of vectorcardiographic finding of right ventricular hypertrophy (RVH) due to ischemia in coronary artery disease. Sixty-two patients with 1 vessel coronary artery diseases (38 of LAD, 16 of RCA and 8 of LCx) and 21 control subjects were studied. All of them were performed coronary angiography, left ventriculography, vectorcardiography (VCG) and echocardiography (UCG). Pattern recognition of Chou