Reply

Dr. Dunlop could be right in interpreting some of our results as demonstrating a VRM “acquired within 100–130°C of room temperature”. The experimental evidence for this is the existence of the double peak which Dr. Dunlop points out in our Fig. 2. However, two further considerations are relevant.

First, if the VRM above 200°C does, as Dunlop suggests, represent a consolidated VRM acquired within 100–130°C of room temperature, then occasionally we should observe a *reversed* consolidated VRM, acquired during a reversed field period. The only hint we have ever had of this came from heating the NRM of specimens from Mull (Wilson, 1969), which had certainly had a complex thermal history, having been in the pneumatolytic zone of the Mull caldera. So in general there is a lack of evidence of reversed consolidated VRM's, which counts against Dunlop's suggestion.

The second consideration concerns the rate of lowering of temperature from 100–130°C to room temperature, perhaps as the specimen becomes uncovered by erosion. If this rate of lowering of temperature is slow enough, covering a few million years (of Tertiary time) then the field will be reversing its polarity many times during the cooling, and one would expect that no net consolidated VRM would exist, because the overall effect would average to zero. After the specimen reached room temperature, the VRM acquired henceforth would take place in the usual way. This may well explain:

- (a) The apparent lack of PNRM in many specimens at intermediate temperatures of a few hundred degrees Centigrade.
- (b) The existence of consolidated VRM's which do exist in rocks that were re-heated during Permo-Carboniferous times when the Kiaman reversed polarity state endured for tens of millions of years, permitting a slow cooling without polarity changes. Under this hypothesis, we need not invoke a unique epoch of remagnetization of rocks just during Kiaman times.

Dr. Dunlop's strongest point is that the existing theory plus a more reasonable assumption (than ours) about the form of $J_s H_c(\theta)$ near room temperature, denies the possibility of a room temperature VRM which could have blocking temperatures as high as 300°C. If we were forced to the point, we would not fail to consider the possibility that the theory could be wrong while we would at the same time agree that his assumptions about $J_s H_c(\theta)$ are better than ours. However, the clash is not as great as it may seem to be. Our highest θ_{demag} values (VRM blocking temperatures) lie *between* 270 and 320°C, and may well tend toward the lower end of this range. Further, the assessment of this temperature is not absolutely objective, since it involves judgement of where a curve becomes tangent to a straight line, with experimental errors as a complication. It is therefore conceivable that we have estimated our θ_{demag} on the high side. These two considerations could reduce the discrepancy between our experimental results and Dunlop's theoretical ones, perhaps to an acceptable level.

Reference

Wilson, R.L., Magnetic properties and normal and reversed natural magnetization in Mull lavas.
Geoph. J. Roy. Astr. Soc., 8, 4, 424–439, 1964.

R.L. WILSON and P.J. SMITH
Department of Geophysics
The Oliver Lodge Laboratory
The University of Liverpool
Liverpool 3, England