Observations of Rapid Auroral Fluctuations

Takasi OGUTI

*Geophysics Research Laboratory, University of Tokyo,
Tokyo 113, Japan*

(Received February 7, 1978)

Rapid auroral fluctuations are categorized into several groups according to their deformation modes, and a composite distribution of these modes is given with respect to the auroral oval. The distribution could impose some restrictions on the nature of plasma in the magnetosphere involved in the auroral precipitations.

1. Introduction

The rapid fluctuation of the aurora, in its luminosity and in its pattern as well, is a fairly common condition of auroral displays and has long been studied, since as early as the end of the 9th century. The early studies have inevitably been based on visual observations so that these have mostly been qualitative descriptions on the most remarkable features with a rather limited range in periodicity. These observations have been summarized by CAMPBELL and REES (1961).

The second phase of the research was opened by the introduction of photomultipliers. Since then, a quantitative discussion of the temporal variations has become possible. Subsequently the research activity on this subject has greatly increased and great progress has been made. That progress has been summarized in several comprehensive papers, e.g., CAMPBELL and REES (1961), SHEPHERD and PEMBERTON (1968) and KVIFTE and PETERSEN (1969) and books such as *Atmospheric Emissions* (OMHOLT *et al.*, 1969), *The Optical Aurora* (OMHOLT, 1971) and *Aurora* (VALLANCE-JONES, 1974).

These studies, in spite of various locations of observation, seem to approach general agreement on several points regarding the rapid auroral fluctuations such as latitude dependence and local time dependence of the occurrence (HEPPNER, 1954; CAMPBELL and REES, 1961; AKASOFU, 1964, 1968; VICTOR, 1965; JOHANSEN and OMHOLT, 1966; OMHOLT and BERGER, 1967; OMHOLT *et al.*, 1969; and KVIFTE and PETERSEN, 1969) spectral feature of temporal variation (e.g., CAMPBELL and REES, 1961; VICTOR, 1965; JOHANSEN and OMHOLT, 1966; PAULSON and SHEPHERD, 1966; PAULSON *et al.*, 1967; SHEPHERD and PEMBERTON, 1968), and phase differences between several characteristic emissions (PAULSON and SHEPHERD, 1965; EVANS and VALLANCE-JONES, 1965). The relation to the variation in temperature in aurora (HILLIARD and SHEPHERD, 1966), relation to geomagnetic pulsations (CAMPBELL and REES, 1961; VICTOR, 1965; PAULSON *et al.*, 1967; CAMPBELL, 1970), coherency be-

tween the fluctuations from separated fields (SHEPHERD and PEMBERTON, 1968; CAMPBELL, 1970) have been studied also during this phase of the research.

However, photometers alone are not suitable for the determination of rapid deformation modes of aurora. These also give important information on the physical processes of acceleration and/or pitch angle scattering of electrons in the magnetosphere.

The third phase of research thus is indebted to the introduction of high sensitive TV cameras (DAVIS, 1966, 1967; SCOURFIELD and PARSONS, 1969a) that enabled us to record the rapid fluctuations with a sampling rate as high as 60 frames per second. CRESSWELL and DAVIS (1966) have confirmed the occurrence region of pulsating aurora to be the equatorward boundary of the auroral activity, BEACH *et al.* (1968) have visualized the flickering aurora, defining it as a group of small patches (1–5 km) that usually move very rapidly (~ 5 km/sec), and HALLINAN and DAVIS (1970) have found the so-called “auroral curl.” Fast auroral waves noted by CRESSWELL and BELON (1966) and by CRESSWELL (1968) on the basis of the data from separated field photometers have been confirmed by TV imaging by SCOURFIELD and PARSONS (1971a). SCOURFIELD and PARSONS (1971b) also found the expansion mode of pulsating patches, and CRESSWELL and DAVIS (1966) and SCOURFIELD *et al.* (1972) have shown the independence of fluctuations between different patches, and in contrast to this, a certain coherency between the different parts of a single patch. OGUTI (1975a), and OGUTI and WATANABE (1976) have shown the existence of a poleward propagation mode and its close relation to geomagnetic pulsations.

In spite of these increased observations and several morphological studies (e.g., CRESSWELL, 1972; THOMAS *et al.*, 1973; OGUTI, 1975a) on the basis of TV observation, the classifications of variation modes seem still not to be in general agreement, and even a simple morphological picture is still missing as to the distribution of each deformation mode. In this paper, the author tentatively proposes a simple classification by re-examining previous results, and subsequently synthesizes distribution of rapid auroral fluctuations with respect to the global auroral pattern.

2. Classifications of Fluctuation Modes

While not everywhere the same at a given level of activity the aurora is generally understood to follow time sequence involving the substorm growth phase, expansion phase and recovery phase. This idea suggests that the terms “expansion” and “recovery,” at least, define the nature of regions rather than of the temporal phases as a whole. Furthermore this means that the aurora left in the wake or in at lower latitudes could be already in a state of recovery or constitute after effects, while the auroras at the front of an expansion are still in the expansion state.

Following this idea, the aurora is first placed into three categories according to three typical regions, namely the region along and near the expansion front, in the wake, both of them in the dusk to midnight sectors, and in the equatorward region

in the dawn sector. Thus the tentative classifications of rapid auroral fluctuations are proposed as follows:

- 1) In and near discrete forms (Expansion Region)
 - Splitting-unfolding or folding over (including outward streaming rotation or leading-rotation)
 - Flickering
- 2) In smoke-like forms (Wake Region)
 - Rotational variations (curl or trailing rotation, fluttering)
 - Folding and meandering (including some of fluttering)
- 3) In pulsating forms (Recovery or After Effect Region)
 - Pulsation
 - Expansion (including core expansion, streaming and flaming)
 - Poleward propagation of arcs (including streaming)
 - Equatorward propagation of arc fragments (auroral waves)
 - Flash

Examples are shown below.

3. Splitting-Unfolding and/or Folding-over

The left half of Fig. 1 shows an example of what the author calls “splitting-unfolding” of a discrete band. The sequence of photographs is taken every 1/2 sec. The right half schematically illustrates how the splitting and unfolding occur. This also indicates how the protrusion to the upper side (high-latitude side) propagates toward the right (eastward) and how it reverses on the lower side (low-latitude side) during a course of a splitting-unfolding. The length of the arc shown in these pictures is about 80 km, so that the characteristic scale of the deformation here is about 50–60 km.

The splitting-unfolding or folding-over can occur in various scales in the fashion shown (OGUTI, 1975c). Among them, the smallest (1–5 km) and the fastest (several to 20 Hz) often occurs along the discrete arc prior to and during the active phase of auroral expansion; the expansion not necessarily of global scale but of local and small scale. An example is shown in Fig. 2, where the left hand side sequence of photographs, reproduced at the rate of 60 frames per second, indicates deformations of a very thin arc. The direction of this arc is roughly east to west; the left is toward the east, and the top is toward low latitude. The length of arc covered in the pictures here is about 100 km. In the right half of the diagram, the solid lines are traces of the shift of identified peeled-off structures that protrude on the upper side (low latitude side) and the dotted lines are those on the high latitude side, both deduced from the sequential pictures at left. The direction of propagation evidently is reversed beyond the main band, and the trends reveal a strong sheer along the arc clockwise viewed along the magnetic field and consistent with clockwise unfolding shown in Fig. 1.

This mode is apt to be regarded as the same group as flickering, but is classified

Fig. 1. Typical "splitting-unfolding" deformation of an arc seen near magnetic zenith; 03:40:13-03:40:22 UT on May 18, 1971, at Syowa Station. Pictures on the left hand side are sequential reproductions from video tape with a time interval of 1/2 sec and with an exposure of 1/60 sec each. The direction of the arc is roughly in east-west; left is to west and the top is to high latitude. The length of the arc covered here is about 80 km. The right half schematically illustrates how the splitting-unfolding proceeds along this arc. Protrusions on each side of the arc propagate in reverse direction as shown by solid and dotted arrows respectively.

Fig. 2. An example of rapid splitting-unfolding along a discrete arc seen near the zenith. The pictures on the left are sequential reproductions from video tape at the rate of 60 fields/sec; 22:32:03:20/60–22:32:03:59/60 UT on Aug. 25, 1971 at Syowa Station. The direction of the arc is roughly in east-west; left is to east and the top is to equatorward. The right half indicates the shifts of protrusions (split ends) in the same fashion as in Fig. 1. Solid lines are the traces of upper (equatorward) protrusions and the dotted lines are of lower (poleward) protrusions.

here otherwise because the mode is different from the flickering defined by BEACH *et al.* (1968). The periodic splitting here always appears to be a feature of the discrete band itself whereas the flickering of patches appears mostly in the remnants just behind the expansion aurora or behind the fluttering of discrete band. This category could include also outward streaming rotation or leading rotation (OGUTI, 1974) because it can be a kind of splitting-unfolding process where the original arc

is much fainter than the splitting-unfolding part. This kind of fluctuation usually is related to auroral hiss emissions (OGUTI, 1975b).

3.1 Flickering

The condition of flickering is well defined by BEACH *et al.* (1968), such that the flickering aurora consists of a number of interconnected, rapidly moving diffuse spots with scale of 1–5 km in diameter. This condition of the display is often found in the remnant diffuse arc behind the expanding discrete band or behind a rapid fluttering of discrete band. The frequency of the flickering aurora is roughly the same or a little less than that of rapid splitting-unfolding described in the previous section (several to 10 Hz).

Sometimes the spot appears to rotate clockwise around a fixed point or to rotate with coherent periodic change in ellipticity. On other occasions, it appears to be illuminated by fainter, wave-like structures periodically. Sometimes bright spots can even be stationary and exist without change in luminosity, and the fainter wave-like structures only represent the periodic variations. In any case flickering reveals itself as a vibrating column along field line when viewed perpendicular to the magnetic field.

The coherency of the fluctuation is usually good so far as one spot is concerned. The spot changes periodically not only in its luminosity but also in its shape, size and position. The brightening and the motion of different spots are not independent of each other and sequential brightenings usually occur from one spot to the adjacent. The situation is seen in Fig. 3 which indicates the temporal and spatial variations of flickering spots in a form of time-position display. The upper part of the figure is obtained through a slit in NE-SW direction and the lower comes from a slit in NW-SE direction, both 40 km in length. The several upward moving traces in the

Fig. 3. Time-position display of a flickering aurora seen near the magnetic zenith, 23:05:30–23:05:35 UT on Aug. 25, 1971 at Syowa Station. The top and bottom figures are obtained through slits in front of monitor TV in NE-SW and in NW-SE directions respectively. The ordinates of both pictures cover about 40 km. The darkness represent the auroral luminosity. Sequential upward shifts of brightness from spot to spot are seen during the period from 23:05:33 to 23:05:34 in the top figure.

upper part from 23:05:33 to 23:05:34 represent 4 different spots; the upward motion of the traces indicates the sequential brightening of these spots, from NE to SW.

3.2 Rotational variations (curls or trailing rotations)

The curl (HALLINAN and DAVIS, 1970) or trailing rotations (OGUTI, 1974) some-

Fig. 4. An example of q - p occurrence of curls (or trailing rotations) with protrusions seen near the magnetic zenith, 23:20:22:0–23:20:36:5 UT on Aug. 25, 1971, at Syowa Station. The pictures are reproductions from video tape at the rate each 1/2 sec with exposure of 1/60 sec. Length covered here is about 80 km, the left is to west and the top is to poleward. q - p generations and leftward (westward) propagations of the rotational structures are evident.

times show a characteristic quasi-periodic condition both in space and in time. The structure usually emerges from a main arc as a protrusion that consists of one or a few rotational cores. An example of such a repetition of the trailing rotation is shown in Fig. 4. The arc here is roughly in east-west direction; left is toward west and the top is to high latitude; the length is about 80 km. Several groups of small trailing rotations are found to repeatedly appear along an arc in a form of protrusions and they propagate to the left (west). The separation between each group of trailing rotations is found to be about 30 km, the period is about 5 sec and the speed here is about 8 km/sec in this particular example. During the period of occurrence of bright and rapid trailing rotations, structureless q - p chorus emission (similar to the post midnight chorus described by TSURUTANI and SMITH, 1974) often can be observed, but the one-to-one correspondence between the rotation and the chorus emission is not yet confirmed.

3.3 *Sheer folding and meandering*

Another typical mode in smoke-like aurora in the wake is folding likely due to sheer motion, or meandering, without discernible rotations. This mode sometimes is periodic in space and quasi-periodic in time. This may be regarded as a less active and a larger "splitting-unfolding" mode with less prominent rotational unfolding motion, or propagating trailing rotations with no prominent rotations.

Although the four modes mentioned above show differences in appearance, two common features are found with regard to the sense of the protrusion and the sense of rotation. That is to say, a protrusion, when it generates from a core or from an arc and grows outwards, always curves toward the right-hand side when viewed parallel (not anti-parallel) to the magnetic field. Hence the auroral arc usually forms a saw-tooth structure with protrusions that protrude toward the west and toward high latitude or that protrude toward the east and toward low latitude. Even when the main arc is not in the east-west direction, the relative configuration is as described. The sense of rotation also remains clockwise when viewed parallel to the magnetic field. These two are the most essential features of aurora both in the poleward expanding front and the wake behind it in the dusk to midnight sectors. The varieties of appearances could be elaborated upon by considering the relative speeds, relative scale of sizes and relative luminosities of protrusions and rotations.

3.4 *Pure pulsation*

The term pure pulsation here defines independently pulsating patches and bands with quasi-periodic change in luminosity without remarkable change in pattern and in position, except the effect of slow drift as a whole. This is the typical condition of auroral activity in the dawn sectors near the low latitude border of the auroral activity, as noted by CRESSWELL and DAVIS (1966). It has a broad spectral peak around 10 seconds, distributed in a range of periods from a few seconds to a few tens of second, and, more importantly, the period of this mode seems not to depend much upon geomagnetic latitude, although the latitude of its occurrence undoubtedly

varies according to the general activity. The persistence of this 10-sec band is evident in a fairly wide range of latitudes, e.g., from Churchill (IMS observation on Feb. 19, 1977) to Saskatoon (e.g., SHEPHERD and PEMBERTON, 1968). Another important point is that it often includes a period other than and much shorter than 10 sec. That is to say, the 'switch-on' level of luminosity of a pulsating patch often fluctuates with period from 0.1 to 1 sec. This has been noted by OMHOLT *et al.* at Tromsö (1969) and by OGUTI at Syowa Station (1976).

The pulsating patches and arc fragments usually form several groups that may be called "core of pulsations" (see Fig. 6). The cores usually occur aligned along the low latitude border of the auroral activity, and are separated some hundreds of kilometers from each other. This is the most active part of the pulsating aurora. This mode usually relates to geomagnetic pulsations observed under or near the patches, but the correlation between them is usually poor because a number of independently pulsating patches can incoherently affect geomagnetic variations.

3.5 *Expansion mode*

Although very few authors have reported on the expanding patch of aurora (e.g., SCOURFIELD and PARSONS, 1971b) this mode really exists in the core region mentioned above. This might be placed into a pulsation category since the pulsating patch usually shows more or less expansion tendency during the period of its increasing luminosity, but the association of chorus groups with this mode, as seen in Fig. 5, could indicate that it is qualitatively different from the pulsation that usually does not relate to chorus emissions. This mode could appear also as a streaming mode when expansion occurs along the elongated direction of patches or along the interconnection structures between patches or cores.

Another expansion is the expansion of the core. While the individual patches pulsate with about 10-sec period, a core can expand by increasing the number of the member patches or by expansion of individual patches with a longer period, about 10–50 sec in a quasi-periodic fashion. The period of the core expansion likely depends on latitude, being longer in higher latitudes, in contrast to the persistence of the period of 10-sec band. Radar aurora observed from Ottawa (McDIARMID and McNAMARA, 1972) also indicates the existence of non-propagating, periodic echoes with the same period range.

3.6 *Poleward propagating mode*

The poleward propagating mode of aurora is fairly common condition at Syowa Station (OGUTI, 1975a; OGUTI and WATANABE, 1975) although few have been reported at other locations. This mode is a common feature of radar aurora, too (KANEDA *et al.*, 1964; McDIARMID and McNAMARA, 1972). The poleward propagating aurora at Syowa (OGUTI and WATANABE, 1975) shows characteristic linear relations between repetition frequency and propagation speed similar to that of radar aurora found by McDIARMID and McNAMARA (1972). The similarity of the characteristics seems to be sufficient to conclude the identity of the two poleward propa-

July 22 1971

Fig. 5

Fig. 5. Time-position display of "core" expansions and poleward propagating modes: 02:28:43–02:29:16 UT on July 22, 1971 at Syowa Station (top panel). Ordinate covers about 40° of elevation from the equatorward horizon. The core expansion is seen in a horizontal trace and poleward propagating mode is seen in up-going traces, both of them being in q - p fashion. The chorus emissions in a time-frequency spectrogram in the bottom panel indicate that they are related to expansion mode but not to the poleward propagation mode.

Fig. 6

Fig. 6. Distributions of rapid auroral fluctuations. Splitting-unfolding and flickering are common conditions in and near an active discrete arc, and pulsation and flash near low latitude borders of active diffuse aurora in the dawn quadrant as well. The propagating modes (both of poleward and equatorward) occur quite frequently but not always.

gating phenomena. The poleward propagating mode appears in the region somewhat higher in latitude than the region of pure pulsation (core), usually in a shape that envelops the poleward side of the cores. The period of the propagating mode is longer than both the period of pulsating patches and the core expansion occurring at the same time. Very close association with geomagnetic pulsations is an important point of this mode (OGUTI and WATANABE, 1976). When the propagation has east-west components rather than poleward, the streaming condition appears along auroral structures usually elongated in east-west direction.

Figure 5 (top) shows the typical poleward propagating aurora along with the q - p expansion of a core in a form of time-position display again; where the top is to high latitude and the slit covers 40° of elevation from the equatorward horizon. At the middle of the figure, the q - p expansion (q - p increase in number of patches in a core) of a core is seen as q - p thickening of the traces without change in its position, while the propagating mode is seen as traces moving upward with time.

The existence of the poleward propagating mode indicates that the quasi-periodic variations of the plasma in the magnetosphere, some of them at least, are generated in the magnetosphere, in contrast to the belief that all come from outside.

3.7 Equatorward propagating mode

This mode is called "fast auroral wave" by CRESSWELL and BELON (1966). As stated by CRESSWELL (1968), it is really a fast wave-like propagation of faint arc fragments usually elongated in east-west direction, with horizontal scale several tens of km in length and about 10 km in width. The propagation speed is in a range from several tens of km/sec to 200 km/sec, and the repetition time is in a range from 1 to several seconds. The horizontal distance of the traverse of one wave is roughly the same to the poleward propagating mode in the previous section, but the mode of the propagation of the wave here is always smooth with a constant speed during its traverse, in contrast to the stepwise propagation of the poleward propagating mode. The occurrence of this mode seems to be limited to the region below geomagnetic latitude 65° . No evidence of this mode has been found at Syowa Station whereas it is often found at College (e.g., CRESSWELL and BELON, 1966; CRESSWELL, 1968) and at Cold Lake (SCOURFIELD and PARSONS, 1971a). The wave appears to be generated near the low-latitude boundary of or sometimes within the core of pulsating aurora.

3.8 Flash mode

Toward dawn, the pulsating aurora and the expanding aurora, seem to be gradually replaced by the flash mode. The temporal characteristics of this mode are similar to the pulsating aurora but the faster modulation dominates over the 10-sec band in the flash mode. The structure of the aurora involved is diffuse, faint and widespread within surfaces over 100 km in horizontal scale, in contrast to the fairly discrete, small-scale structure of the pulsating patches.

A strong impression exists that the fast modulation of pulsating patches origi-

nally is the nature of the flash mode and that it modulates the switch on level of the pulsating patches by overlapping the patches, whereas the 10-sec band is characteristic of the pulsating patches themselves. This mode is usually associated with VLF chorus emissions, but one to one correspondence between each flash and each riser of chorus is quite hard to find because so many chorus risers and flash auroras usually occur at the same time. The aurora studied by UNGSTRUP (1966), although he called it flickering, must be this mode. Satellite observations of precipitation associated with chorus (e.g., OLIVEN and GURNETT, 1968; HOLZER *et al.*, 1974) likely relate also to this. Most of X-ray bursts so far studied (e.g., ANDERSON *et al.*, 1966; PARKS *et al.*, 1968; VENKATESAN *et al.*, 1968; FOSTER and ROSENBERG, 1976) are likely related to this mode too. Because the change in luminosity of the flash mode is usually so coherent over a fairly widespread area, coherent pulses of X-rays are to be expected. By contrast pulsating patches are smaller and temporally independent so would possibly result in noise-like X-ray variations due to overlapping of effects from many patches (SCOURFIELD *et al.*, 1970).

3.9 *Propagation along an arc (streaming)*

This mode likely appears associated with the expansion mode or poleward propagation mode, as mentioned before, and hence could be classified into either category. Its period is in a range roughly the same as the poleward propagation mode.

3.10 *Propagation along the magnetic field (flaming)*

Although quite many have reported on this mode (e.g., CRESSWELL, 1969; SCOURFIELD and PARSONS, 1969b) it could not be found during the observations of several years at Syowa Station, Antarctica and in Canada, except for an appearance of flaming due to a perspective effect of horizontal motions. It seems to be a very rare condition of aurora and its distribution is not as clear as the other modes.

4. Summary and Discussion

The rapid fluctuation of aurora shows typical modes of variation according to location with respect to the auroral oval. The characteristics of auroral modes in the expansion region and in the wake just behind the expansion front are a combination of a right hand side splitting (or right hand side curving extension of an arm) and of clockwise rotation viewed along the magnetic field. The variations can occur in a wide range of horizontal scale sizes. The smallest and fastest of them are the splitting-unfolding process in the expansion front and flickering in the wake, whereas the largest and the slowest may be the auroral bulge. The middle scale and less active examples are the curl or trailing rotations where the rotational motion dominates, and shear folding where the right hand side splitting dominates.

In the region of recovery (or in the region of after effects), namely in low-latitude part of auroral oval, especially near dawn, the typical condition of the auroral display is pulsation. Here several cores of pulsating aurora that consist of a

number of pulsating or expanding patches appear in an alignment along the oval. The structure of the core gradually becomes diffuse, or from another point of view, the patches are replaced or overlapped by diffuse surfaces that flash with period from 0.1–1 sec near dawn. Much to envelop the core regions to the poleward side of them, the poleward propagating auroras appear. The equatorward propagating mode (auroral wave), on the other hand, appears on the equatorward side of the core region. Thus, the distribution of each mode along an oval can be shown as in Fig. 6.

One of the most important characteristics of the auroral rapid variation is that the energy of the corresponding electrons is higher than that in more common aurora (EVANS, 1976; WHALEN *et al.*, 1971; BRYANT *et al.*, 1971; COURTIER *et al.*, 1974; BRYANT *et al.*, 1975; BROWN *et al.*, 1976). The energy range of the corresponding electrons, in conjunction with the association of VLF chorus emissions and the association of geomagnetic pulsation as well, suggests that the coupling between energetic auroral electrons, VLF waves and HM waves in the magnetosphere plays an essential role in the region of after-effect. Since the core region of pulsating aurora is the most active region of pulsation, this region possibly links to the inner edge of the injected energetic electrons where an inward steep gradient of the plasma pressure is expected, and hence where instabilities such as due to drift-Alfvén and drift-bounce coupling tend to occur. Hence it is possible that the HM waves are first generated in this region and then modulate the pitch angle of the electrons, which subsequently results in an enhancement of cyclotron coupling between energetic electrons and VLF waves under a possible influence of cold plasma. Another possibility that the coupling between VLF waves and energetic electrons, resulting in the rapid escape of electrons in some regions in the magnetosphere, subsequently produces HM waves in this region, could not be overlooked. The propagating modes are understood to be due to compressional HM-waves generated here. HM waves could also be produced near the ionosphere as a result of q - p precipitation of electrons through q - p change in ionospheric electric current, possibly being fed back to the magnetosphere.

On the other hand, the precipitation in the expansion region must be understood in another way. The clockwise rotation suggest that a charge excess is one of the main reason of the precipitation here. Here, the relation between auroral hiss emissions and splitting-unfolding and flickering (the latter appears to relate to stationary hiss emission although this is not confirmed yet) is understood in terms of coherent Cerenkov emission that might be generated from some corrective motions of electrons at the steep boundary of small structures in the electron beams such as that corresponding to the splitting-unfolding and flickering. The fact that auroral hiss is mostly observed only when the corresponding aurora shows rapid variation supports this mechanism. The possibility of a wave conversion mechanism from electrostatic to electromagnetic near or in the auroral precipitations also can not be excluded. In any case, the close relation between auroral rapid variation and burst-like enhancement of auroral his emission not only in time (OGUTI, 1975b) but also in space (HAYASHI *et al.*, 1978 (this issue)) suggests that the two phenomena are

coupled, not in a wide-spread region in the magnetosphere, but in a space relatively near to the electron precipitation.

REFERENCES

- AKASOFU, S.-I., The development of the auroral substorm, *Planet. Space Sci.*, **12**, 273-282, 1964.
- AKASOFU, S.-I., *Polar and Magnetospheric Substorms*, Reidel Publ. Co., Dordrecht, Holland, 1968.
- ANDERSON, K.A., L.M. CHASE, H.S. HUDSON, M. LAMPTON, D.W. MILTON, and G.K. PARKS, Ballon and rocket observations of auroral-zone microbursts, *J. Geophys. Res.*, **71**, 4617-4629, 1966.
- BEACH, R., G.R. CRESSWELL, T.N. DAVIS, T.J. HALLINAN, and L.R. SWEET, Flickering, a 10-cps fluctuation within bright auroras, *Planet. Space Sci.*, **16**, 1525-1529, 1968.
- BROWN, N.B., T.N. DAVIS, T.J. HALLINAN, and H.C. STENBAEK-NIELSEN, Altitude of pulsating aurora determined by a new instrumental technique, *Geophys. Res. Lett.*, **3**, 403-404, 1976.
- BRYANT, D.A., G.M. COURTIER, and G. BENNETT, Equatorial modulation in a pulsating aurora, in *The Radiating Atmosphere*, edited by B.M. McCormac, pp. 170-175, Reidel Publ. Co., Dordrecht, Holland, 1971.
- CAMPBELL, W.H., Rapid auroral luminosity fluctuations and geomagnetic field pulsations, *J. Geophys. Res.*, **75**, 6182-6208, 1970.
- CAMPBELL, W.H. and M.H. REES, A study of auroral coruscations, *J. Geophys. Res.*, **66**, 41-55, 1961.
- COURTIER, G.M., M.J. SMITH, and D.A. BRYANT, Auroral electrons observed using a mother-daughter rockets, in *Magnetospheric Physics*, edited by B.M. McCormac, pp. 207-211, Reidel Publ. Co., Dordrecht, Holland, 1974.
- CRESSWELL, G.R., Fast auroral waves, *Planet. Space Sci.*, **16**, 1453-1464, 1968.
- CRESSWELL, G.R., Flaming auroras, *J. Atmos. Terr. Phys.*, **31**, 179-183, 1969.
- CRESSWELL, G.R., The morphology of displays of pulsating auroras, *J. Atmos. Terr. Phys.*, **34**, 549-554, 1972.
- CRESSWELL, G.R. and A.E. BELON, Observations of fast auroral waves, *Planet. Space Sci.*, **14**, 299-301, 1966.
- CRESSWELL, G.R. and T.N. DAVIS, Observations on pulsating auroras, *J. Geophys. Res.*, **71**, 3155-3163, 1966.
- DAVIS, T.N., The application of image orthicon techniques to auroral observation, *Space Sci. Rev.*, **6**, 222-247, 1966.
- DAVIS, T.N., Cinematographic observations of fast auroral variations, in *Aurora and Airglow*, edited by B.M. McCormac, pp. 133-141, Reinhold Publ. Co., New York, 1967.
- DAVIS, T.N., Auroral morphology, *Planet. Space Sci.*, **20**, 1369-1373, 1972.
- EVANS, D.S., A 10-cps periodicity in the precipitation of auroral-zone electrons, *J. Geophys. Res.*, **72**, 4281-4291, 1967.
- EVANS, W.F.J. and A. VALLANCE-JONES, Some observations of type B aurora with a multichannel photometer, *Can. J. Phys.*, **43**, 697-704, 1965.
- FOSTER, J.C. and T.J. ROSENBERG, Electron precipitation and VLF emissions associated with cyclotron resonance interactions near the plasmapause, *J. Geophys. Res.*, **81**, 2183-2192, 1976.
- HALLINAN, T.J. and T.N. DAVIS, Small-scale auroral arc distortions, *Planet. Space Sci.*, **18**, 1735-1744, 1970.
- HAYASHI, K., K. TSURUDA, S. KOKUBUN, T. OGUTI, R.E. HORITA, and T. WATANABE, Arrival directions of VLF emissions and corresponding activities in aurora and geomagnetic pulsations, *J. Geomag. Geoelectr.*, **30**, 361-362, 1978.
- HEPPNER, J.P., Time sequences and spatial relations in auroral activity during magnetic bays at College, Alaska, *J. Geophys. Res.*, **59**, 329-338, 1954.
- HILLIARD, R.I. and G.G. SHEPHERD, Upper atmospheric temperatures from Doppler line width-IV, A detailed study using the OI 5577 Å auroral and nightglow emission, *Planet. Space Sci.*, **14**, 383-406, 1966.

- HOLZER, R.E., T.A. FARLEY, R.K. BURTON, and M.C. CHAPMAN, A correlated study of ELF waves and electron precipitation on Ogo 6, *J. Geophys. Res.*, **79**, 1007–1013, 1974.
- JOHANSEN, O.E. and A. OMHOLT, A study of pulsating aurora, *Planet. Space Sci.*, **14**, 207–215, 1966.
- KANEDA, E., S. KOKUBUN, T. OGUTI, and T. NAGATA, Auroral radar echoes associated with Pc-5, *Rep. Ionosph. Space Res. Japan*, **18**, 165–172, 1964.
- KVIFTE, G.J. and H. PETERSEN, Morphology of the pulsating aurora, *Planet. Space Sci.*, **17**, 1599–1607, 1969.
- MCDIARMID, D.R. and A.G. McNAMARA, Periodically varying radio aurora, *Ann. Geophys.*, **28**, 433–441, 1972.
- OGUTI, T., Rotational deformations and related drift motions of auroral arc, *J. Geophys. Res.*, **79**, 3861–3865, 1974.
- OGUTI, T., Metamorphoses of aurora, *Mem. Natl. Inst. Polar Res., Ser. A, No. 12*, 1–101, 1975a.
- OGUTI, T., Hiss emitting auroral activity, *J. Atmos. Terr. Phys.*, **37**, 761–768, 1975b.
- OGUTI, T., Similarity between global auroral deformations in DAPP photographs and small scale deformations observed by a TV camera, *J. Atmos. Terr. Phys.*, **37**, 1413–1418, 1975c.
- OGUTI, T., Recurrent auroral patterns, *J. Geophys. Res.*, **81**, 1782–1786, 1976.
- OGUTI, T. and T. WATANABE, Quasi-periodic poleward propagation of on-off switching aurora and associated geomagnetic pulsations in the dawn, *J. Atmos. Terr. Phys.*, **38**, 543–551, 1976.
- OLIVEN, M.N. and D.A. GURNETT, Microburst phenomena, 3. An association between microbursts and VLF chorus, *J. Geophys. Res.*, **73**, 2355–2362, 1968.
- OMHOLT, A., *The Optical Aurora*, Springer-Verlag, New York, 1971.
- OMHOLT, A. and S. BERGER, The occurrence of auroral pulsations in the frequency range 0.01–0.1 c/s over Tromsø, *Planet. Space Sci.*, **15**, 1075–1080, 1967.
- OMHOLT, A., G.J. KVIFTE, and H. PETERSEN, Pulsating aurora, in *Atmospheric Emissions*, edited by B.M. McCormac and A. Omholt, pp. 131–143, Van Nostrand Reinhold Co., New York, 1969.
- PARKS, G.K., F.V. CORONITI, R.L. MCPHERRON, and K.A. ANDERSON, Studies of magnetospheric substorm, 1. Characteristics of modulated electron precipitation occurring during auroral substorms, *J. Geophys. Res.*, **73**, 1685–1696, 1968.
- PAULSON, L.V. and G.G. SHEPHERD, A cross spectral method for determining the mean lifetime of metastable oxygen atoms from photometric observations of quiet auroras, *J. Atmos. Terr. Phys.*, **27**, 831–841, 1965.
- PAULSON, K.V. and G.G. SHEPHERD, Short lived brightness oscillations in active aurora, *Can. J. Phys.*, **44**, 921–924, 1966.
- PAULSON, K.V., G.G. SHEPHERD, and P. GRAYTONE, A note on auroral-type fluctuations in the earth's electromagnetic field, *Can. J. Phys.*, **45**, 2813–2821, 1967.
- SCOURFIELD, M.W.J. and N.R. PARSONS, An image intensifier-vidicon system for auroral cinematography, *Planet. Space Sci.*, **17**, 75–81, 1969a.
- SCOURFIELD, M.W.J. and N.R. PARSONS, Auroral pulsations and flaming, *Planet. Space Sci.*, **17**, 1141–1147, 1969b.
- SCOURFIELD, M.W.J. and N.R. PARSONS, Television imaging of fast auroral waves, *Planet. Space Sci.*, **19**, 437–442, 1971a.
- SCOURFIELD, M.W.J. and N.R. PARSONS, Pulsating auroral patches exhibiting sudden intensity-dependent spatial expansion, *J. Geophys. Res.*, **76**, 4518–4524, 1971b.
- SCOURFIELD, M.W.J., W.F. INNES, and N.R. PARSONS, Spatial coherency in pulsating aurora, *Planet. Space Sci.*, **20**, 1843–1848, 1972.
- SCOURFIELD, M.W.J., G.R. CRESSWELL, G.R. PILKINGTON, and N.R. PARSONS, Auroral pulsations-television image and X-ray correlations, *Planet. Space Sci.*, **18**, 495–499, 1970.
- SHEPHERD, G.G. and E.V. PEMBERTON, Characteristics of auroral brightness fluctuations, *Radio Sci.*, **3**, 650–658, 1968.
- THOMAS, I.L., M.W.J. SCOURFIELD, and N.R. PARSONS, Classification of optical auroral pulsations, *Can. J. Phys.*, **51**, 2209–2215, 1973.
- TSURUTANI, B.T. and E.J. SMITH, Post midnight chorus: A substorm phenomenon, *J. Geophys. Res.*, **79**, 118–127, 1974.

- UNGSTRUP, E., Association between VLF emissions and flickering aurora, *J. Geophys. Res.*, **71**, 2395–2396, 1966.
- VALLANCE-JONES, A., *Aurora*, D. Reidel Publ. Co., Dordrecht, Holland, 1974.
- VENKATESAN, D., M.N. OLIVEN, P.J. EDWARDS, K.G. MCCracken, and M. STEINBOCK, Microburst phenomena, 1. Auroral zone X-rays, *J. Geophys. Res.*, **73**, 2333–2343, 1968.
- VICTOR, L.J., Correlated auroral and geomagnetic micropulsations in the period range 5 to 40 seconds, *J. Geophys. Res.*, **70**, 3123–3130, 1965.
- WHALEN, B.B., J.R. MILLER, and I.B. MCDIARMID, Energetic particle measurements in a pulsating aurora, *J. Geophys. Res.*, **76**, 978–986, 1971.