Difference between Fresh and Sea Water Fishes in the Accumulation of Environmental Chemical Pollutants. II Uptake and Elimination of Chlorobenzenes by Killifish (*Oryzias latipes*)

MARIKO TACHIKAWA,^a RYOJI SAWAMURA,^a SHOJI OKADA,^b ATSUSHI TAKEDA,^b YUTAKA KAWAZOE,^c and MASATO NARA^d

Department of Pharmacy, College of Science & Technology, Nihon University, Tokyo 101, Japan,^a School of Pharmaceutical Sciences, University of Shizuoka, Shizuoka 422, Japan,^b Faculty of Pharmaceutical Sciences, Nagoya City University, Nagoya 467, Japan^c and Shizuoka Prefectural Fisheries Experiment Station, Yaizu 425, Japan^d

Since fresh water fish and seawater fish are well-known to have different physiological mechanisms for the regulation of their osmotic pressure, the accumulation of environmental chemical pollutants in fish might differ between fresh and seawater areas. To evaluate this difference, we have investigated the differences of accumulation and excretion of environmental chemical pollutants such as pentachlorophenol (PCP), by utilizing euryhaline fishes that can be acclimated to seawater or fresh water. In the investigation of PCP accumulation and elimination by fresh and seawater killifish, the following information was obtained. (1) Fresh water killifish accumulated PCP faster and to a greater extent than seawater killifish. (2) PCP was eliminated from seawater fish faster than from fresh water fish. (3) The proportion of PCP metabolites excreted, PCP sulfate and PCP glucuronide, was different in fresh water and in seawater. Therefore, we investigated the effect of the salinity of surrounding water to elucidate further the relationships between the regulation of osmotic pressure and

the behavior of PCP in accumulation and excretion, and examined the accumulation and elimination of 1,2,3-trichlorobenzene (TCB) in fresh water and seawater killifishes to see the effect of chemical properties.

Fresh water killifish (0.3–0.4 g in body weight) were transferred into seawater, with various levels of salinity (0.1–1.9‰) and which contained 100 µg/l of PCP, and were maintained in it for 24 h. The accumulation of PCP in killifish decreased as the salinity of the seawater was increased from 0.1‰ to 0.47‰. When the salinity was increased to 0.47‰, the PCP accumulated in fish decreased to about 60% of that in fresh water. The accumulation of PCP in the fish began to decline about 8 h after the transfer. At the salinity of 0.23‰, although the accumulation of PCP did not decrease significantly, an increase of excretion of PCP conjugates into the surrounding water was observed. Slight changes of salinity (0.1–0.5‰) in the surrounding water altered the accumulation and excretion of PCP rapidly and effectively.

When the killifish were exposed to 1 mg/l of 1,2,3-TCB, the concentrations of 1,2,3-TCB in fresh water and seawater fish were *ca.* 250 and 200 µg/g fish, respectively. As regards the elimination of 1,2,3-TCB, fresh water killifish eliminated it faster than seawater killifish. The elimination half times for fresh water and seawater killifish were 13 h and 31 h, respectively. These results were different from those obtained on PCP and indicated that the differences of accumulation and elimination of chemical pollutants in fresh water and seawater fish were dependent on the properties of the chemicals.

These differences between fresh water and seawater fish in accumulation and elimination might be important for evaluation of the ecological hazard posed by chemical pollutants.