The Role of Zinc in Fatty Acid Metabolism in Rat Liver: Effects of Zinc Deficiency and Cadmium Administration

NAOMI KUDO,^a YASUHIITO NAKAGAWA,^a YOICHI KAWASHIMA,^b HIROSHI KOZUKA^b and KEIZO WAKU^a
Faculty of Pharmaceutical Sciences, Teikyo university,^a Sagamiko, Kanagawa 199-01, Japan and Faculty of Pharmaceutical Sciences, Toyama Medical and Pharmaceutical University,^b 2636 Sugitani, Toyama 930-01, Japan

Zinc is known to be an essential metal, and its deficiency is characterized by loss of appetite, growth retardation, skin defects, hair loss and impaired wound healing. Recent studies have demonstrated that dietary zinc deficiency can affect the metabolism of fatty acids such as the desaturation reaction by which polyunsaturated fatty acids are produced. Cadmium, an environmental pollutant, is thought to have an antagonistic effect on zinc in some biological reactions. However, little is known of how cadmium affects fatty acid metabolism. In this study, we compared the effects of zinc deficiency with those of cadmium administration on fatty acid composition and metabolism in the liver of rats fed a fat-free diet. In addition, we estimated microsomal desaturation activity *in vitro* in zinc-deficient and cadmium-treated rats.

Three-week-old male Wistar rats were fed a fat and zinc free diet for two weeks (ZD-rats). ZA-rats were supplemented with zinc (50 ppm) and were pair-fed to ZD-rats. Half of the ZA- and ZD-rats were injected with CdCl₂ (0.01 or 0.03 mmol/kg body weight) subcutaneously 2 or 48 h before sacrifice. Fatty acid analysis of liver phospholipids revealed that the levels of 18:1(*n*-9) and 20:3(*n*-9) were decreased and those of 18:0, 20:4(*n*-6) and 22:6(*n*-3) were increased in ZD-rats compared with ZA-rats. Exposure to cadmium

caused changes in the fatty acid composition of liver phospholipids, which were similar to those caused by zinc deficiency. The changes caused by cadmium were greater in ZD-rats than in ZA-rats.

Next, [¹⁴C]18:0 was injected intravenously 24 h before sacrifice in order to estimate the metabolism of (*n*-9) fatty acids. The radioactivity was recovered in 18:0, 18:1(*n*-9) and 20:3(*n*-9) in liver phospholipids. A decrease in the rate of conversion of [¹⁴C]18:0 to [¹⁴C]18:1 was observed in liver phospholipids in ZD-rats. When exposed to cadmium, the decrease was significant in ZD-rats but not in ZA-rats. These results suggested that the activity of Δ9 desaturase, which converts 18:0 to 18:1, was reduced in ZD-rats, especially when cadmium was administered.

Δ9 Desaturase activity can be recovered in the microsomal fraction in rat liver. This enzyme activity requires cytochrome b₅, NADH-cytochrome b₅ reductase and terminal desaturase. We estimated the individual enzymes of liver microsomal fraction *in vitro*. Terminal desaturase activity in ZD-rats exposed to cadmium was reduced to 1/10 of that in ZA-rats without significant changes in the amount of cytochrome b₅ or the activities of NADH-cytochrome c reductase and NADH-ferricyanide reductase. However, there was no significant change in the activity of terminal desaturase between ZA- and ZD-rats. The activity of lysophosphatidylcholine(lysoPC): 18:1-CoA acyltransferase, which catalyzes the transfer of 18:1 to lysoPC, was not changed in all groups. Terminal desaturase seemed very sensitive to cadmium toxicity and the decreases in the levels of 18:1(*n*-9) and 20:3(*n*-9) appeared to result mainly from the reduced activity of terminal desaturase when ZD-rats were treated with cadmium.

These results suggested that cadmium can affect the metabolism of fatty acid, in particular Δ9 desaturation of 18:0, in relation to zinc status.