Induction by Cadmium of DNA Fragmentation in Porcine Kidney LLC-PK₁ Cells

MASAMI ISHIDO, SHINO T. HOMMA, PO S. LEUNG and CHIHARU TOHYAMA

*National Institute for Environmental Studies, 16–2 Onogawa, Tsukuba,
Ibaraki 305, Japan*

Introduction

Cadmium is a nephrotoxic metal. The development of cadmium-induced lesions in the kidney is characterized by proteinuria and excessive urinary excretion of other substances such as enzymes, amino acids, and glucose. Biochemical characterization of the cellular mechanism involved in cadmium cytotoxicity has been hampered by difficulties in identifying target molecules, both on the cell surface and in the interior, and by lack of inhibitors which could identify a specific pathway. In cultured cells, cadmium cytotoxicity has been observed as perturbation of the energy producing system, cellular membrane defects and increased influx of calcium ions at the end-point of toxicity of the metal. New biochemical markers with high sensitivity for detecting and identifying cadmium cytotoxicity is needed to analyze the mechanism of the action of the metal at the cellular level.

In this study, we demonstrate for the first time that a low dose of cadmium can induce DNA fragmentation which is characteristic of apoptosis in the early stage of cytotoxicity in porcine kidney LLC-PK₁ cells.

Experimental Procedures

LLC-PK₁ cells were grown in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum, penicillin (100 U/ml), and streptomycin (100 µg/ml) in a humidified atmosphere of 95% air and 5% CO₂ at 37°C. To examine cadmium-induced DNA fragmentation, cells (4×10^6) were subcultured at 70% confluency

and maintained in serum-free medium for 16 h prior to the addition of cadmium and other metals tested. For treatment with TPEN, *N,N,N',N'*-tetrakis(2-pyridyl-methyl)-ethylenediamine, cells were incubated at 37°C for 40 min before exposure to cadmium. Treated cells were then washed twice with PBS, and lysed in 5 mM Tris buffer and 0.5% Triton X-100, pH 7.4, containing 20 mM EDTA at 4°C for 20 min. After centrifugation at $28000 \times g$ for 20 min, supernatants were extracted with phenol-chloroform and precipitated in ethanol. One half of each sample was treated with 20 µg/ml RNase A and electrophoresed on a 1.2% agarose gel.

Results

DNA fragmentation was induced by the addition of cadmium to cultured LLC-PK₁ cells, resulting in cell death. LLC-PK₁ cells were able to survive exposures to 10 µM cadmium without change in morphology, but most had rounded by 40 µM. DNA fragmentation induced by 10 µM cadmium was completely inhibited using an equimolar concentration of a cell-permeant chelator specific for heavy metals, TPEN. Furthermore, the apoptotic mechanism evoked by the metal was blocked by a five molar excess of Zn²⁺ which, when used alone, has no capacity for DNA fragmentation. Other metals tested, Cu²⁺, Co²⁺, Ni²⁺, and Pb²⁺ had much lower ability to induce DNA fragmentation in LLC-PK₁ cells under the same conditions used for cadmium, indicating that the fragmentation seen in kidney cells is cadmium-specific.