THE JOURNAL OF THE OTO-RHINO-LARYNGOLOGICAL SOCIETY OF JAPAN

EDITOR: C. SATTA

Published by
The Oto-Rhino-Laryngological Society of Japan.
University of Tokyo, Bunkyo-ku, Tokyo.

Abstract of Original Papers

EXPERIMENTAL STUDIES PERTAINING TO THE RESISTANCE OF EACH PART OF THE EAR DRUM AGAINST INFLAMMATION

By

K. KUSUOKA

*(The Otorhino-laryngology Section, Medical School,
University of Osaka.)*

Classifying the inflammations into bacteriogenic and non-bacteriogenic inflammations, three kind of bacteria: streptococcus haemolyticus, staphylococcus aureus, and pneumococcus (I form) were used as agencies of the former type, while 1% solution of carbonic acid and 5% solution of sodium carbonate were employed for causing the latter type.

By cutting the skin open at the back part of the ear of a guinea-pig until the bulla was exposed, one of the noxious material above mentioned was poured therein. The length of terms let by after pouring the material was 6 hours, 12 hours, 24 hours, 2 days, 4 days, 7 days and 10 days in each experimental group respectively. Thus, when the pouring-in process of the inflammation material was all over, the guillotine-fixation was made, the tissue block was inbedded in celloidin and serial sections were made parallel with the malleus handle and stained in haematoxylin-eosin for microscopical study.

CONCLUSIONS: 1. The connection of the mucous layer and the proper layer, compared with that of the cutaneous layer and the proper layer, was coarser and rougher.

2. The cutaneous layer was the strongest in the resistance against the inflammation, followed by the proper layer and the mucous layer in the order of strength.

3. The lower part of the ear drum was stronger in resistance than the

upper part.

4. Posterior upper part was more resistant than anterior upper part, while no differentiation of strength in resistance was recognizable between the posterior lower and anterior lower parts.

CASE OF PANMYELOPHTHISIS.

By

Y. HIROKAMI.

*(From the Oto-rhino-laryngological Department, Medical School,
University of Niigata.)*

Author reported a case of panmyelophthisis, 53 aged female, and discussed the origin of the name of this disease and the difference between two definitions, panmyelophthisis and agranulocytosis. In conclusion he stated that, there is only some quantitative difference of pathological changes in bone-marrow between these two diseases.

HEARING IMPAIRMENT AND THE RESIDUAL PERFORATION OF THE TYMPANIC MEMBRANE.

By

S. SOENO.

*(Department of Oto-Rhino-Laryngology, Medical School,
University of Tokyo.)*

Author classified the residual perforation of the tympanic membrane into five types according to the size of perforations (fig 1.) In the fifth type tympanic membrane is absent, as is the case after the radical mastoidectomy. When there is some adhesion or ankylosis of the membrane and ossicles, it is described as (type I+adhesion).