nearly the same as those of the middle ear catarrh cases.

3) In the ruptured tympanic membrane cases, the sound did not diminish within the limits of ± 30 mmHg.

4) In most of the cases the sound was perceived less distinctly when the pressure was raised or lowered. On the other hand, in a few cases hearing acuity became better under the abnormal pressure condition in the tympanic cavity.

THREE CASES OF ACUTE SINUITIS WITH OSTEOMYELITIS IN INFANTS.

By

T. KAMIO

(From Kamio Oto-rhino-laryngological Hospital, Tokyo.)

Case 1: Boy, four year old, Maxillary empyema and osteomyelitis derived from acute rhinitis. Cured operatively in a month.

Case 2: Girl, two year old. Sudden unaccountable attack of fever reaching 40°C, followed by swelling on the right side of the face. Also cured operatively in 2 months.

Case 3: Girl, ten months old. Sudden unaccountable fever reaching 40°C, then swelling on the left part of the frontal region, which spread gradually to left maxillary region, while purulent nasal discharge appeared. Cured by repeated radical operations on the maxillary as well as frontal bones in 5 months.

Statistics of all 78 cases reported to the Oto-rhino-laryngological Society of Japan from 1908 to 1948 are as follows:

Etiology: nasal 52 %, dental 18 %.

Age: under 3 months 36 %, 3~6 months 19 %.

Exit: recovery 63 %, deaths 27 %.

ON THE PATH OF SOUND THAT CAUSES INNER EAR DEAFNESS IN REPEATED APPLICATION

By

H. FUJINO

(From the Oto-rhino-laryngological Department of Setagaya National Hospital)

The author examined acoustic and vestibular function of 270 labcrers em-