OBSERVATIONS ON THE PROGRESSIVE NASAL GANGRENE

By

T. KIKUTI, T. TAKAFUJI

*(From the Otorhinolaryngological Department, Medical
School University of Tokyo.)*

and

Y. HOMMA

*(From the institute of Infections Disease, Medical School,
University of Tokyo.)*

Six cases of progressive nasal gangrene were reported. Various forms of treatment were tried without effect. All of them died eventually. Concerning the cause of this disease results of haematologic, histologic, bacteriologic autoptic and therapeutic studies were considered. Here are some:

(1) From the haematologic findings:

All these cases are free from any sign of blood disease. The anaemia was of aplastic or infectious, toxic type and secondarily caused by the primary disease.

(2) From the histological observation:

All cases showed the picture of chronic nonspecific inflammation without any sign of malignant tumor.

(3) From the bacteriological findings:

As to the cause of progressive necrosis, symbiosis of the bacillus proteus with several other bacteria were found most important. But it was uncertain whether there was relationship between this disease and the pleuropneumonia group which were found on the lesion.

(4) From the autoptic findings:

Two cases were examined. We observed that these patients had exerted very little resistance and reaction to this disease. Chronic, infections and toxic changes were manifest in the liver and the spleen in both cases.

(5) From the therapeutic observations:

Medication with chemical, immunological and antibiotic remedies did not secure any effect. Also X-ray or radium irradiation was without effect.