Precise and Statistical Studies of Morphology,
Physiology and Taxonomy on *Absidia*. Part I Morphology,
No. 1 Sporangial head

Kenji Yamashita

(Kyoto City)

ABSTRACT

The author advocated two words;

1) "sporangium proper" for only the sac full of sporangiospores and 2) "sporangial head" for a new entity, lumping sporangium proper apophysis and columella that is a whole thing produced at the top of sporangiophore. In this report 1 the sporangial head was accurately measured and examined by slide culture, as well as slant and dish culture. The slide culture made the examination easy and accurate.

The materials used were authentic strains obtained from CBS, ATCC and IFO. The data were expressed as numbers or figures as far as possible and the numerical data were analyzed statistically by including nonparametric statistical methods. The breadth and height of *Absidia* were examined. Statistically 62 distributions (31 for breadth, and 31 for height) were tested for normality. Only nine distributions were normal and the other 53 were not normal. The nine normal distributions were treated by the usual parametric methods and the 53 nonnormal by nonparametric methods.

Three ranges were presented: the common range included 80% of samples, the larger 96% and the largest, Min.-Max. The first two ranges were calculated statistically, especially when the distribution was normal, by parametric methods.

Many authors have reported simply and uniformly that the form of the sporangium (accurately sporangial head) is pyriform. The natural pear has a depression at the tip and at the base. The author named this form "eupyriform". However, an eupyriform head has never been found on sporangial head by the author, who has examined over 3000 *Absidia*. He classified sporangial heads into four categories: 1. almost pyriform 40 (1.29%), 2. subpyriform 600 (10.49%), 3. pseudopyriform 1859 (59.97%), 4. never pyriform 597 (10.25%). The pseudopyriform category was subclassified into: **z**. having a cone-formed neck of apophysis and **s**. having a concave funnel-like or trumpet-like neck.

Absidia causes otomycosis and rhinocerebral mucormycosis. The latter is a lethal disease, from sinusitis of the nose directly invades the brain or indirectly first makes a raid on the orbit then advances to the brain. Therefore this fungus is an important one for medical doctors, especially for ENT specialists as the author.

About forty-five years ago the author got in Taiwan very many fungi from the ears of otomycosis. Almost all of them were *Aspergilli* but four strains were Mucoraceae.

The fungi looked like *Absidia*. However all mycological books and literatures described that in *Absidia* sporangia were pyriform and at the nodes of stolon produce peculiar

*Former professor, Kansai Med. Univ. Taipe Med. College, Pekin Univ. and vice president National Kyoto Hosp.

rhizoids. Nevertheless the author's strains had neither pyriform sporangia nor rhizoids which easily found in *Rhizopus*. Therefore the author was much perplexed in identification of these strains.

At that time the author found accidentally the following description “*Lichtheimia* 'avec ou sans rhizoid'” in a French book (Brumpt 1949). Then they were published provisionally as *Lichtheimia* sp. Soon after Vuillemin (who created the genus *Lichtheimia*) the genus was considered generally as a synonym of *Absidia*. Then “*Lichtheimia* sp. “should be corrected to “*Absidia*” sp.” After the first experience on the strains from the ear, the author had often got such fungi later; from nose, mouth, throat, skin, house dust and air in the room. Is there a species of *Absidia* having neither peculiar rhizoid nor eupyriiform sporangium? For the author this question had not been solved till the beginning of this research. Because the retirement from official service and duty, life environment of the author changed to be able to take hours for studies and researches of *Absidia*, in order to solve the questions for long years.

Then authentic strains of *Absidia* were collected from the world famous organization of collection. As it was a good chance, the author has performed and is performing precise and statistic morphologic, physiologic and taxonomic studies of this genus. The results would be serially published in a few years. At first here Part I. Morphology, No. 1. Sporangial heads would be described.

“Sporangial head” is a new technical term proposed by the author (Fig. 1). In Latin sg. “Capitulum sporangialis” “pl. Capitula sporangiales”.

Fig.1 Schematic figure of sporangial head

As in *Aspergillus*, a unit including vesicle, sterigmata and conidial chains is called “Conidial Head”, Apophysis, Columella and sporangium proper (full of sporangiospores) as a whole is called “Sporangial head (pl. -s)”. Till now either one sporangium proper (a sac) or a unit (including sporangium proper, columella and apophysis as a whole) has been called “Sporangium”. Such nomenclature would cause some errors. The author's new nomenclature is right and does not cause any error.

MATERIALS AND METHODS

The Strains examined were as follows (in alphabetic order) They are all authentic

strains got from collectionsorgans. In the bracket optimal temperature for growth was added.

1. *Absidia anomala* Hesseltine et Ellis 1964. (30°C) CBS 125.68
2. *Ab. atropura* Naganishi et Hirahara (30°C) CBS 518.71
3. *Ab. blakesleana* Lendner. 1923. (35°C) 1970. ATCC 10148a
4. *Ab. californica* Ellis et Hesseltine (30°C) CBS 126.68
5. *Ab. coerulea* Bainier. 1988. (25°C) 1965. IFO 4011 (M)
6. *Ab. corymbifera* (Cohn) (30°C) 1912. Sacc. et Trotte IFO 4009 (+)
7. *Ab. cuneospora* Ore et Prunkett 1959. (35/C) ATCC 13693
8. *Ab. cylindrospora* Hagem 1908. (37°C) IFO 4000 (+)
9. *Ab. fassatia* Vanova 1871. (25°C) ATCC 24445
10. *Ab. fusca* Linneman 1936. (25°C) CBS 102.25
11. *Ab. ginsan* Kominami et al. 1952. (37°C) CBS 100.15
12. *Ab. glauca* Hagem 1908. (37°C) IFO 4002 (+)
13. *Ab. glauca* var. *paradoxa* Namyslowski 1906. (25°C) IFO 4007
14. *Ab. griseola* Naganishi et Hirahara 1970 (30°C) IFO 9472
15. *Ab. grisrola* var. *iguchii* Naganishi et Hirahara 1970. (35°C) ATCC 20431
16. *Ab. hesseltinii* Mehorthra et Naud 1963. (30°C) ATCC 24263
17. *Ab. heterospora* Ling-Young 1930. (30°C) ATCC 24264
18. *Ab. hyalospora* (Saito) Lendner 1908. (35°C) IFO 8082
19. *Ab. macrospora* Vanova 1968. (25°C) CBS 696.68
20. *Ab. orchidis* (Vuillemin) Hagem 1908. (35°C) IFO 4423
21. *Ab. parricida* Renner et Muscat ex Hesseltine et Ellis 1961. (27°C) CBS 174.67
22. *Ab. psrudocylindrospora* Hesseltine et Ellis 1958. (27°C) CBS 100.67
23. *Ab. psychrophila* Hesseltine et Ellis 1964. (40°C) CBS 128.16
24. *Ab. ramosa* (Lindt) Lendner 1908. (30°C) IFO 8083
25. *Ab. reflexa* van Tieghem 1876 27°C IFO 5875
26. *Ab. regnierii* (Lucet et Constantin) Lendner 1908. (32°C) IFO 8084
27. *Ab. repens* van Tieghem 1876. (35°C) CBS 100.19
28. *Ab. spinosa* Lendner 1907. (30°C) IFO 5873
29. *Ab. spinosa* var. *biappendiculata* Rall et Solheim 1964. (35°C) ATCC el5465
30. *Ab. sporophora-variabilis* Subrahmanyam (30°C) ATCC 36019
31. *Ab. zychnae* Hesseltin & Ellis 1966. (30°C) CBS 104. 35

Abbreviations of the organs from which the authentic strains derived are as follows.

IFO (Institution Fermentation Osaka Japan)

CBS (Centraalbureau voor Schimmelculture, Baarn Netherlands)

ATCC (American Type Culture Collection, Rockville U. S. A.)

The temperatures noted in bracket are optimal temperature for the growth, that had been experimentally decided in part II Physiology

Culture medium used chiefly was synthetic Mucor medium (SMA) (Hesseltine, 1954) and often also yeast extract Agar (Hesseltine 1954) was additionally or profitably employed

The fungi were made grown in slant tube, petri-dish and on slide.

The methods of the slide culture was as follows. From the culture medium solidified in a petri-dish a thin small and oblong piece (about 1-1.5 × 9 × 12 mm) was cut out and placed on a slide-glass sterilized by flame of Bunsen burner while it is still warm at about 45°-50°C. On the middle of the four sides the smallest piece of the fungus to be grown was inoculated, then a sterilized and cooled to about 45°-50°C cover-glass of 18 × 24 mm was put on the inoculated piece of medium. The specimen thus prepared was laid on

a U-form glass rod in a petri-dish and about one ml of sterilized water dropped into the dish and incubated in thermostat at the optimal temperature for the growth noted in bracket in the list above, that would be described perisely in part II physiology. The growth of the *Absisia* strain in slant tube, in petri-dish and on slide were examined at 24 hours, 48 hours 72 hours, 5 days, one week, two weeks and four weeks after inoculation. They were at first examined macroscopically, then directly microscopically, in the order from slant tube, petri dish to slide culture. The growth in the slant tube was examined from outside of the tube through glass at the upper and outer edge of the growth with light from down through Abbe's condenser and that in petri dish with the light falling from up and a little obliquely. In these two cases the object lens used had to be of weak power (4x, 5x at times 10x). On the contrary in the case of the slide culture, high power object lens (20x, 40x, 100x for oil immersion) could be used; therefore the enlargement from 20x to 2000x was able to be gotten, at measurement of size as well as observation. The objective organs sporangial heads and sporangiophora, should be in the position perpendicularly to the direction of vision through microscope. This condition could be got hardly both in the cases of slant and dish but easily in the case of slide culture. As far as the condition was satisfied, the objects were examined always at random. The last examination was usual microscopy: pick up a small piece of fungus and mount in lactophenol without any staining matter on a slide glass.

The findings of the examination were expressed in numerals (continuance and discrete variables) as far as possible and the numerals were treated by modern statistical methods (Masuyama, M. 1952, Assoc. Research, Statistics ed. 1952, Siegel, S. 1956, Fisher, R.A. 1970, Bishop, O.N. 1971, Kishine, T. 1974, Campbell, R.C. 1974, Fisher, R.A. et Yates, F. 1975, Yamashita, K. 1976, Ishii, S. 1977, Siegel S. 1956, Snedecor, G.S. et Cochran, W.G. 1978)

The number of samples per each item was always one hundred, except a few cases of which the number was specially noted.

For continuous variables at first the normality of each distribution was tested with goodness of fit test, and tests with skewness and kurtosis (e.g. Snedecor G.S. et Cochran W.G. 1978, 84-89).

Because normal distribution was found in a few cases or rarely, the nonparametric statistical methods were applied for all thirty one strains. As the value representing each of the distributions the median (Me.) calculated from frequency table was adopted in the place of arithmetical mean for normal distribution and other statistical treatments of data were performed by the methods of nonparametric statistics.

For range of sample, four kinds were determined (1) the common range including 80% of samples, (2) the larger range including 95% of samples, (3) the largest range including minimum-maximum of all 100 samples, per each strain and (4) the absolute range min-max of all 3100 samples (100 samples per each of 31 strains). When a distribution was normal, the common range was determined by calculation of $(\bar{x}-1.28 u)$ - $(\bar{x}+1.28 u)$ and the larger one by $(\bar{x}-1.95 u)$ - $(\bar{x}+1.95 u)$, where $\bar{x}$ was arithmetic mean and u =unbiased variance=unbiased sample standard deviation. When distribution is not normal, the formulae described above could not be utilized. The lower and upper border of the common and larger range were determined with the method as follows. Of the common range including eighty percent of samples the lower border was the eleventh in a table of 100 samples arranged in the order of number on each item of each strain and the higher border the ninetieth in the table. As for the larger range including ninety-five percent of samples, the lower border was determined by the average of the second and third values and the higher border that of ninety seven and ninety-eight.

When around a border of the common or larger range there were two or more same values, the following process was taken. The unavoidable error of micrometre for the author was one tenth of one degree of ocular micrometer. For example in the case of 40x15, as one degree of micrometer corresponds to $2.5 \mu\text{m}$, the unavoidable error was $0.25 \mu\text{m}$. Then the least value of a group having the same size was considered to be $-0.25 \mu\text{m}$ and the most $+0.25 \mu\text{m}$. The value between the least and the most were determined also by interpolation respectively.

Comparison of two median (see Table, 1, 2) was performed by Median test (e. q. Siegel, s. 1956, pp. iii-116) and Kolmogorov-Smirnov's two sample test (e. q. Ibid. pp 127-136). At first the highest value of median (Or. Me. 1 in Table 1 and 2) was compared to the second one (Or. Me. 2 in Table 1, 2). If there was no significant difference between them (company 1 vs. 2), the highest median was compared to that of the third (Or. Me. 3) namely (company 1 vs. 3). Such comparison repeated to the following one, 1 vs. 2, 1 vs. 3, 1 vs. 4, 1 vs. 5.....successively. When the significant difference was gotten for the first time (e.g. company 1 vs. 4 in Table 1), the species or variety (Or. Me. 4) was admitted to the top of the second group, and the value just before it (Or. Me. 3) was admitted the last of the first group. These process repeated successively to make several groups. If the values to be compared were dependent each other, the method of comparison of ratio utilizing F-table (Musumoto, M. 1952 pp 23-24, Yamashita, K. 1926 pp 1733-1734) and one sample test (Siegel, S. 1956, pp 42-47) was taken. On the rare cases with normal distribution, the comparison was performed also by usual method utilizing arithmetical mean ($\bar{x}$), unbiased standard deviation (u) and t-distribution table or F-distribution table.

The coefficient of variance (v.c.) is $\frac{s}{\bar{x}} 100$ where s =sample standard deviation, and $\bar{x}$ =sample arithmetic mean.

RESULTS A SIZE

The results of quantitative (height and breadth) examinations of on sporangial heads in twenty-eight species and three varieties were shown in Table 1, 2. The abbreviations in the tables meant as follows. Or. Me.: order of median, C.V.: coefficient of variance, Me.: median, Norm.: normality of distribution, common range: the range including 80% of samples, larger range: that including 95% of samples, Min.-Max.: minimum to maximum. Comp.: company of comparison, vs.: versus (=against), Me. -test: median test, K.S. two: Kolmogorov-Smirnov's two-sample test, gr.: group rank, ind.: individual rank. P: probability, Or. $\bar{x}$ u order by arithmetical mean u: unbiased sample standard deviation.

Measurement of the height of sporangial head was often not easy, because the under border of apophysis was not rarely uncertain. Nevertheless, when measuring of the breadth of sporangiophore was progressed gradually to the direction to the head, the point could be determined from where the breadth seemed to begin to increase. The point was considered to be the under border of the apophysis, i. e. the under border of the sporangial head.

The table 1. shows that the height of sporangial head was variable as much as eight groups could be divided with significant difference each other.

The coefficient of variance (C.V.) also very variable; ranging from 12.31% (Or. Me. 27 *Ab. anomala*)-58.73% (Or. Me. 22 *Ab. paricida*).

Table. I. Height of Sporangial Head
A. nonparametric data

Or. Me.	Species or Variety	C.V. (%)	No- rm.	Me. (μm)	Range			Difference			Rank	
					Common (μm)	larger (μm)	largest (μm)	comp.	Me.- test	K.S. two	Gr.	Ind.
1	Ab. coerulea	31.63	—	47.19	25.50-64.93	20.13-77.50	18.00-85.00	1 vs. 4	p0.001	p0.001	I	1
2	Ab.blakesleana	22.98	+	45.73	26.25-60.18	26.25-71.25	22.50-75.00					1
3	Ab. griseola	55.71	—	39.13	13.15-70.15	4.95-100.12	2.00-115.00					1
4	Ab. heterospora	30.94	—	38.75	24.91-55.00	19.79-65.00	18.00-65.00	4 vs. 10	p0.001	p 0.0005	II	4
5	Ab. ramosa	37.84	—	37.78	23.50-59.90	19.00-70.16	12.60-70.00					4
6	Ab. griseola var. igudhii	34.63	—	36.32	20.08-53.25	16.25-77.50	15.00-80.00					4
7	Ab. macrospora	36.05	—	35.89	25.05-50.08	18.75-61.75	17.50-87.50					4
8	Ab. corymbifera	27.92	++	32.83	21.99-48.01	16.51-59.95	13.00-62.00					4
9	Ab. hyalospora	41.87	—	29.24	21.50-55.50	16.50-73.50	11.00-110.00	10 vs. 20	p0.05	p0.005	III	4
10	Ab. regnierii	32.44	—	28.39	17.17-42.41	10.50-51.25	7.00-52.50					10
11	Ab. glauca var. paradoxa	31.08	—	27.85	19.00-49.96	15.08-53.80	6.00-60.00					10
12	Ab. atrospora	21.88	—	27.06	19.83-37.65	11.25-51.00	9.00-57.50					10
13	Ab. orchidis	19.55	—	26.75	19.95-24.61	12.75-33.13	13.00-50.00					10
14	Ab.cylindrospora	30.51	—	26.73	18.21-40.16	15.18-50.05	15.00-50.00					10
15	Ab. californica	23.09	—	25.81	9.15-35.04	7.49-40.06	11.20-47.56					10
16	Ab. zychae	42.13	—	25.25	19.89-52.34	14.75-64.00	12.50-70.00					10
17	Ab. repens	38.88	—	24.81	17.41-38.91	14.00-54.10	12.50-75.00					10
18	Ab. ginsan	26.21	—	23.67	18.28-35.04	15.05-40.31	11.00-35.00					10
19	Ab. sporangioph- oravariabilis	35.79	—	23.34	17.45-41.25	15.00-50.00	14.00-62.50	20 vs. 23	p0.01	p0.005	IV	10
20	Ab. fusca	49.09	—	22.88	13.25-47.60	8.00-55.00	7.50-61.5					20
21	Ab. glauca	47.60	++	22.53	13.50-33.75	10.50-46.25	9.00-60.00					20
22	Ab. parricida	58.73	—	21.84	14.89-41.25	10.25-56.24	5.00-70.00	23 vs. 27	p0.0001	p0.005	V	20
23	Ab. hesseltina	32.22	—	21.00	14.87-30.21	12.52-40.00	11.00-46.00					23
24	Ab. spinosa var. biappendicu- lata	30.04	—	20.92	12.77-25.09	10.08-36.55	10.00-37.50					23
25	Ab. psychrophila	36.42	—	19.43	12.20-30.21	9.88-40.19	8.50-43.00					23
26	Ab. reflexa	28.10	—	18.13	15.08-35.21	11.25-49.75	3.00-56.00	26 vs. 29	p0.001	p0.005	VI	26
27	Ab. anomala	12.31	—	17.92	12.41-30.75	10.25-34.00	10.00-40.00					26
28	Ab. cunesopora	28.96	—	17.00	12.67-27.62	12.15-35.13	10.00-35.50					26
29	Ab. spinosa	32.84	++	15.00	10.50-25.00	8.00-30.25	7.00-32.50	29 vs. 31	p0.0001	p0.01	VII	29
30	Ab. fassatiae	37.23	—	14.43	11.88-23.75	8.08-31.25	8.00-35.00					29
31	Ab. pseudo- cylindrospo- ra	36.42	—	13.00	12.27-30.21	9.88-40.17	6.50-45.00	30 vs. 31	p0.001	p0.01	VIII	31
	Average	33.09		26.50	14.80-31.14	11.81-37.80	10.75-58.44	absolute range 2.00-115.00				

B. parametric data

Or. $\bar{x}$.	Species	x (μm)	u (μm)	Range		Difference	
				common (μm)	larger (μm)	comp.	p
1	<i>Ab. blakesleana</i>	47.16	10.65	32.25 - 60.18	26.02 - 68.30	1vs. 2	0.01
2	<i>Ab. corymbifera</i>	34.48	9.82	22.31 - 47.37	15.76 - 53.93	2vs. 3	0.01
3	<i>Ab. glauca</i>	23.86	9.02	12.35 - 35.25	6.36 - 41.24	3vs. 4	0.01
4	<i>Ab. spinosa</i>	16.26	5.29	9.36 - 23.18	5.75 - 26.09		

In the samples with normal distribution, the ranges were calculated as follows : common range, $(\bar{x}-1.28u)-(\bar{x}+1.28u)$ and larger range $(\bar{x}-1.95u)-(\bar{x}+1.95u)$.

Median (Me.) was 26.5 μm in an average and also very variable ranging from 13.00 (Or. Me. 31) to 47.19 μm (Or. Me. 1), namely the largest median was about 3.6 folds of the smallest.

Common range including 80% of samples was 14.80-31.14 μm in an average.

Larger range including 95% of samples was 11.86-37.86 μm in an average.

The largest range including 100%, i. e. from min. to max. in each strain, was averagely 10.75-58.44 μm . The ratio 58.44: 10.75 was about 5.5: 1 or 0.85: 0.15.

Absolute range of the height of all 3100 sporangial heads, (100 samples per each of all 31 strains) was 3.00-115.0 μm .

The comparison of the median of each strain was performed nonparametrically by Me-test and Kolmogorov-Smirnov's Two sample test, because the normal distribution was found in only four strains (Tab. 1-B). The tests produced eight group. The first group includes *Absidia coerulea* (Or. Me. 1), *Ab. blakesleana* (Or. Me. 2) and *Ab. griseola* (Or. Me. 3). The median of *Ab. coerulea* had not significant differences from that of *Ab. blakesleana* and also from that of *Ab. griseola*, but showed a significant difference from that of *Ab. heterospora* (Or. Me. 4). Then *Ab. heterospora* was given the rank of the top of IInd group.

In turn the differences of the median of *Ab. heterospora* from those of the following species were tested successively in order of median (Or. Med.). When the tests arrived at *Ab. reenerii* (Or. Me. 10) where the difference of the median values was for the first time significant. Therefore *Ab. reenerii* got the rank of the top of IX1-Ard group.

Such comparisons and establishment of groups were performed successively in the same way in the order of median value (Or. Me.) and at last they reach VIIIth group that includes two fungus *Ab. sptnose* *Ab. pseudocylindrospora*.

The differences between the species or variety in the same group each other were not significant. Between two adjoining groups, only the top one in a larger group was exactly higher than any member of the next group. The other members of the larger group had not always larger value than the members of the next group, especially the one in lower orders of the larger group. Therefore it was safe to consider that one group had higher or lower value than the secondly adjoining group, for example, each member of the IIIrd group lower than any member of the Ist group and higher than that of the Vth group.

The data on the height of sporangial head of four species having normal distribution and treated parametrically were shown in Table I-B. Difference was determined by the standard method utilizing F-distribution table. Or. $\bar{x}$. 1, *Ab. blakesleana* belonged to the first group rank in Table 1-A. *Ab. corgmbifere* belonged to the II group rank in Table 1-A in hree Or $\bar{x}$ 2 in Table 1-B. *Ab. glauca* belonged to the IV group in Table 1

and here Or $\bar{x}^3$. in table 1-B. Ab. spinosa, belonged to VII group rank in Table 1-A was here in Table 1-B Or. $\bar{x}^4$ in Table 1-B. Therefore it would be presumed that nonparametrical order is I, II, IV, VII is almost agree with parametrical one, Or. $\bar{x}^1$ 1, 2, 3, 4.

Table 2 Breadth of Sporangial Head
A. nonparametric data

Or.	Species	C.V.	No- mr.	Me.	Range			Difference			Rank	
Me	or Variety	(%)		(μm)	common (μm)	larger (μm)	largest (μm)	comp.	Me.- test	K.S. two	Gr.	Ind.
1	Ab. griseola	35.75	—	41.28	24.51-60.06	20.50-80.13	18.90-90.00	1 vs. 4	p0.001	p0.001	I	1
2	Ab. coerulea	37.50	—	40.54	20.20-60.13	15.18-62.50	15.00-65.00					1
3	Ab. blakesleana	23.77	++	38.07	27.54-47.38	24.95-62.25	18.00-67.50					1
4	Ab. ramosa	32.20	—	31.65	19.00-48.13	15.40-57.50	12.50-62.00	4 vs. 7	p0.005	p0.001	II	4
5	Ab. corymbifera	29.67	++	30.06	20.01-40.00	11.99-50.01	10.00-58.00					4
6	Ab. hyalospora	32.33	—	29.51	20.00-48.00	15.00-55.13	12.00-58.00					4
7	Ab. griseola var iguchii	32.70	++	27.04	17.59-37.79	17.54-38.79	15.10-50.00	7 vs. 9	p0.005	p0.005	III	7
8	Ab. zychnae	30.15	+	26.00	16.50-36.50	14.50-41.30	13.00-45.00					7
9	Ab. heterospora	21.96	—	23.95	19.80-32.16	15.08-38.66	15.00-40.00	9 vs. 13	p0.005	p0.005	IV	9
10	Ab. californica	24.08	—	23.60	18.55-27.94	13.50-49.00	13.00-50.00					9
11	Ab. orchidis	21.06	—	23.32	17.50-27.00	15.00-30.00	12.50-37.50					9
12	Ab. macrospora	24.47	—	22.91	17.18-30.00	12.50-35.00	10.50-37.00					9
13	Ab. atropora	27.71	—	22.02	15.15-29.96	11.25-37.75	7.50-42.10	13 vs. 15	p0.001	p0.001	V	13
14	Ab. regnierii	27.66	—	21.10	14.43-27.70	9.25-33.75	7.50-47.50					13
15	Ab. cylindrospora	33.96	—	20.46	14.94-31.94	12.50-40.06	11.00-42.00	15 vs. 23	p0.05	p0.01	VI	15
16	Ab. ginsan	23.59	—	20.45	12.50-27.25	12.50-32.50	11.00-35.00					15
17	Ab. sporophora- variabilis	29.17	—	20.37	14.90-29.80	12.55-36.25	12.50-40.00					15
18	Ab. glauca var. paradoxa	16.46	—	19.56	13.01-27.75	13.01-27.75	10.50-30.17					15
19	Ab. reflexa	35.28	—	17.96	12.61-26.25	8.90-35.58	6.00-40.00					15
20	Ab. repens	27.42	—	17.73	14.00-29.56	11.25-32.62	10.00-35.00					15
21	Ab. parricida	37.79	—	17.50	11.20-30.19	9.00-36.00	5.00-40.00					15
22	Ab. cuneospora	24.89	—	17.35	12.33-23.88	10.50-27.84	10.00-28.50					15
23	Ab. fusca	46.89	—	16.33	9.15-35.63	7.47-40.00	7.50-42.50	23 vs. 28	p0.001	p0.001	VII	23
24	Ab. psychrophila	39.74	—	15.34	9.79-20.50	8.00-40.00	7.00-40.00					23
25	Ab. anomala	32.34	—	15.17	9.91-17.57	7.56-21.25	10.00-40.00					23
26	Ab. glauca	29.18	++	14.81	7.72-17.56	7.25-22.50	4.00-25.00					23
27	Ab. hesseltinii	11.09	—	14.51	12.51-22.50	10.18-24.50	10.00-27.50					23
28	Ab. fassatiae	28.17	—	12.43	10.03-17.60	7.50-20.00	7.00-25.50	28 vs. 30	p0.005	p0.01	VIII	28
29	Ab. spinosa var. biappendiculata	28.62	—	11.72	9.97-17.50	7.71-22.56	5.00-32.50					28
30	Ab. spinosa	33.12	—	11.65	7.95-19.75	7.15-21.00	5.00-23.00	30 vs. 31	p0.05	p0.05	IX	30
31	Ab. pseudocylindrospora	25.41	—	10.93	7.91-17.25	7.15-19.50	6.50-22.50					30
	average	28.91		22.32	14.80-30.69	11.86-37.84	10.75-58.44	absolute range for all 3100 heads 4.00-90.00				

B. parametric data							
Or. x.		x (μm)	u (μm)	Range		Difference	
				common (μm)	larger (μm)	comp.	p
1	Ab. blakesleana	39.04	9.37	27.05 - 51.03	20.77 - 57.39	1 vs. 2	0.01
2	Ab. corymbifera	30.60	9.17	18.86 - 43.34	12.72 - 48.48	2 vs. 3	0.01
3	Ab. griseola var. iguchii	27.52	9.14	15.82 - 38.22	9.70 - 45.34	3 vs. 4	0.01
4	Ab. zychae	24.94	7.60	15.21 - 56.67	10.12 - 39.76	4 vs. 5	
5	Ab. glauce	13.60	4.09	8.37 - 18.84	5.60 - 21.58		0.01

Pertaining to the breadth, the values of measurement and of statistical treatment were shown in Table 2. The meaning and significans of each values in the Table 2 were almost the same as those of Table 1 .

Coefficient of variance (C.V.) was 28.91% in an average. It varies from 11.09% (Or Me. 27, *Ab. hesseltinii*) to 46.89% (Or. Me. 23, *Ab. fusca*) and max.- min. wps about 4:1 or 0.8:0.2.

Median (Me.) was 22.32 μm in an average and it varies from 10.93 (Or. Me. 31, *Ab. pseudocylindrospora*) to 41.28 μm (Or. Me. 1, *Ab. griseola*).

Distribution of median was not normal in twenty-six species or varieties and normal in only five. Therefore nonparametric statistics was taken in all thirty-one species and varieties (Table 2, A) and the data of parametric statistics were also added for the five species and one variety having normal distribution (Table 2, B).

Common range including 80% of the samples was from 14.80 to 30.69 μm in an average.

The larger range including 95 percent of samples varies from 11.86 to 37.84 μm in an average.

The largest range was from 10.75 to 58.44 μm averagely; the ratio 58:44 10.75 was 0.78:0.22 or almost 5:4 times.

The absolute range in all 3100 heads was 4.0-90.0 μm . Also on the breadth median was compared in the same way as of the height, one group differd with quite reliability from two other groups adjoined secondly, e. g. the Vthh group was smaller than the IIIrd group and larger than the VIIth group.

The data of five cases with normal distribution treated by parametric methods were given in Table 2-B. The order of arithdetical mean (Or. $\bar{x}$) was not differed much from that of median (Or. Me).

RESULT B. FORM (Fig. 2. in report, 1 and Fig. 2 and Taebl III in this report)

Schematic drawings of sporangial heads were shown in Fig. 2, and frequencies of each types of sporangial head were presented in Table III.

- Sporangial head consists in general of body (B in Fig 2) and neck (N in Fig 2).
1. Eupyriform has several characterestiiis. (i) There are depression both at the top (da) and at the base (db), (ii) Transition from body to neck is so gradually (gt) that border line between body and neck can hardly be determind, (iii) Neck (apophysis) is like church bell (campaniformis).
 2. Almost pyriform. Generally like eupyriform but the depressions at the top and the base do not exist. Neck (apohysis) is like church bell.
 3. Subpyriform; Neck is lodger than that of eupyriform and of almost pyriform (α), or cylindrical (β). Neck of eupyriform of almost pyriform, has single and simple convection, but some of subpyriform have irregular convexion (γ), at times double

convexion (γ). Neck often is like cup (δ) or glpss (ϵ).

4. Pseudopyriform. Neck has not convection. Very often apophysis (neck) is like cone (α) or concave funnel or trumpet ($\beta_1, \beta_2, \beta_3$).

5. Never pyriform. Lack of typical N (apophysis) 5-I. Apophysis uncertain.

5. Never pyriform. Lack of typical N (apophysis)

5-I. Apophysis uncertain.

Between B and N there is a disk (α), thin saucer (β), (γ) or a nipple.

5-II. Apophysis (N) certainly none. B is spheric (α).

β) B is lying subpheric

γ) B is leaf form

δ) B is standing ellipsoid

ϵ) B is turbinate

From a century ago till now the form or shape of the sporangium (rightly “sporangial head”) of *Absidia* has been always considered and described as “pyriform”. Nevertheless, the sporangial head having the accurately the same form as that of real fruit of *pyrus communis* have never been found, at least by the author who have examined over five thousand sporangial heads of *Absidia*.

Typical fruits of *pyrus communis* has several characteristic form. (Eupyriform see schema of page Fig.1-2. 1. On a globose larger body attaches smaller neck. 2. The body transits to the neck so gradually that the border line between them can hardly be determined, gt in Fig.2 is abbreviation for gradual transition 3. There are depressions more or less remarkable at the top (da, da, da) and bottom (db, db, db) 4.

In the most typical case the neck looks like a church bell (campanulate). or (short trumpet)

There are rare sporangium looking like “Eupyriform” in *Absidia* but the lack of depressions at the top and bottom was grave and remarkable difference. Such form named by the author “Almost pyriform” (Fig. 2-2,3).

There are many sporangial heads of which transition of the body to the neck is not gradual but there some angle is formed and border line between body and neck can be traced. Fig. 2-4. An and ac an: angular transition producing some angular. ac: not gradual but acute transition with short radius of curvature in Fig. 2-4 Pseudopyriform. The neck is often tea-cup form (Fig. 2-4 δ) or form of glass for drinking water or beer glass (Fig. 2-4 ϵ) form (3 γ). The neck is convex but usually simply convex but at times it has double convexities (Fig. 2-3 γ). In the last one: body transits quite gradually to neck. Though the neck has a peculiar characters, two convexities, this is belonged in to subpyriform, because of important gradual transition from body to neck was hold clearly body and others (3.22%) 7. Almost pyriform (1.29%).

DISCUSSION

In the area of mycology the statistical methods have almost not been utilized yet or very few it has been used. As far as the author knows there has been only one example, i.e.. U-test of Mann and Whitney and Spearman's rank coefficient, both nonparametric statistical methods, were used in a paper (Notter, H., H.J. Scholer 1974). Subjective rough descriptions e. g. as many and rare or few, large and small, high and low or thin and thick etc. are used very often even now. Such evaluation and description have to be avoided to make the level of mycology up to modern precise science.

The first step for the purposes, the evaluation and description should be expressed by numerals or figures continuous and discrete variable, and the size of samples per each item has always to be concretely described as “The number of all items in this report is one hundred”. Secondly the numerals gotten have to be treated by modern

Fig. 2. Schematic drawings of oporangial heads.

Explanation of Fig 2

Fig. 1. Eupyriform: Typical form of natural fruit of *pyrus communis* (common pear) This form is that of true fruit of *pyrus communis* and can never be found in *Absidia* B: Body N: Neck d: depression db: depression at base or bottom da: depression at apex gt: gradual transition an: angular transition, Fig. 2. Almost pyriform Like Eupyriform but without depression at base and apex 3. Subpyriform: Apophysis show remarkably different forms than 2. Breadth of Neck near that of almost near that of almost pyriform Neck is a narrow at cylinder Apophysis has two convexity Apophysis cup-form, transitions from Body to Neck not gradual but angle formationc Apophysis glassform for drinking water or beer Fig. 3. Pseudo pyriform side wall straight, therefore Apophysis is cone-like Side wall concaved, like concave funnel or trumpet β_1 =short trumpet, β_2 =trumpet of moderate length. β =long trumpet Fig. 4. Neverpyriform 1. Apophysis uncertain :a δ disc between Body and Apophysis β nipple between Body and sporangophore thin saucer in the place of Apophysis thick saucer in the place of Apophysis. II. Apophysis certainly none : Body spheric : α Body subspheric : β Body leaf-form : Body standing ellipsoid : Body turbinate

Table III. From of Sporangial Heads

Alphabetical order species or variety	almost pyriform	subpyriform			pseudo- pyriform		never pyriform				sum
		α	β	$\gamma_{\delta\epsilon}$	α	β	α	β	γ	δ_{ϵ}	
1 Ab. anomala	0	5	6	10	40	22	2	13	1	1	100
2 Ab. atrospora	1	4	18	5	39	18	3	10	2	0	100
3 Ab. blakesleana	0	1	3	5	20	48	0	15	7	1	100
4 Ab. californica	6	8	8	4	44	18	0	10	7	1	100
5 Ab. coerulea	4	10	12	13	23	12	0	14	9	3	100
6 Ab. corymbifera	0	3	0	13	49	0	0	25	9	1	100
7 Ab. cuneospora	0	1	20	1	41	7	0	18	10	2	100
8 Ab. cylindrospora	1	0	0	5	18	50	0	5	7	14	100
9 Ab. fassatia	1	7	5	4	34	25	0	21	2	1	100
10 Ab. fusca	2	4	9	5	38	26	0	9	3	4	100
11 Ab. ginsan	2	1	16	7	40	11	0	18	5	0	100
12 Ab. glauca	0	3	14	17	39	18	1	5	2	1	100
13 Ab. glauca var. paradoxa	1	9	2	1	42	33	2	6	0	4	100
14 Ab. griseola	0	13	6	4	27	31	0	12	1	6	100
15 Ab. griseola var. iguchii	7	14	4	3	34	27	0	4	4	3	100
16 Ab. hesseltinii	1	7	7	3	48	19	0	14	1	0	100
17 Ab. heterospora	2	11	8	6	27	38	2	3	1	2	100
18 Ab. hyalospora	1	4	4	3	30	17	0	19	11	11	100
19 Ab. macrospora	3	3	15	10	45	21	1	1	1	0	100
20 Ab. orchidis	0	2	11	8	24	35	1	14	3	2	100
21 Ab. parricida	0	9	13	12	39	15	0	10	2	0	100
22 Ab. pseudo- cylindrospora	1	3	9	4	37	24	0	7	7	8	100
23 Ab. psychrophila	0	3	8	4	20	30	7	15	7	6	100
24 Ab. ramosa	0	0	1	0	15	12	0	16	16	0	100
25 Ab. reflexa	2	2	7	5	29	22	0	18	8	7	100
26 Ab. regnierii	0	1	6	4	24	43	0	12	3	7	100
27 Ab. repens	0	5	13	7	31	33	2	9	0	0	100
28 Ab. spinosa	2	13	8	5	30	26	0	2	4	10	100
29 Ab. spinosa var. biappendiculata	2	4	13	2	50	19	0	2	2	6	100
30 Ab. sporophoravariabilis	1	1	9	6	42	21	0	13	4	3	100
31 Ab. zychae	0	12	8	2	38	32	0	8	0	0	100
total	40	163	263	178	1096	763	21	339	134	103	3100
percent	1.29	5.25	8.48	5.74	35.35	24.61	0.67	10.93	4.32	3.22	100
order	9	5	4	5	1	2	10	3	5	8	
probability	0.01		0.01	0.01	0.01	0.01		0.01		0.01	
total	40	604			1859		597				3100
percent	1.29	19.49			59.97		19.25				100
order(p)	3	2(0.05)			1(0.01)		2(0.01)				
Total		644			1859		597				3100
percent		20.78			59.97		19.25				100
order(p)		2(0.05)			1(0.01)		2				

statistical methods.

Of course numerical expression have been often practiced. For example, the size of an organ almost always has been described a μm or b-c $\mu\text{m} \times$ d-e μm , but it was not certain that the figure "a" was arithmetical mean, median or mode and that in "b-c" $\times$ "d-e" μm the numerals are minimum to maximum or show other range including 70%, 80% or 90% of samples. The value "a" has to be exactly and mathematically noted what sort of value it is, i.e. arithmetical mean, median or mode, and the width of the range is to be "b-c" $\times$ "d-e" given concretely such as "including x% of samples" or from minimum to maximum. Perhaps it is for the first time in this report in mycological area that "a" is clearly described as median (Me.) or arithmetical mean ($\bar{x}$) and that concrete value of the width of range has been written as follows; common range includes 80% of samples, larger range 95% and largest covers min. – max. per each strain and absolute whole 3100 heads. (100 heads per each species or variety in all 31 strains).

Besides those matters, the numerical data have to be treated by the modern statistical methods as far as we can (refer to books and a paper such as 1. Japanese Association of Researches on Statistic ed. 1952. 2. Masumoto, M. 1952, 3. Siegel, S. 1956 4. Bishop, O.N. 1970, 5. Fisher, R.A. 1970, 6. Campbell, R.C. 1974, 7. Fisher R.A. and F. Yates 1975, 8. Kishine, T. 1975, 9. Ishii, S., 1975, 10. Yamashita, K. 1976, 11. Snedecor, G.S. et Cochran 1978).

All books and papers describe or at least impress the readers that in *Absidia* all or almost all heads are "pyriform". As described minutely in the chapter of results, it is true in no case. As eupyrimform, i.e. pyriform in very strict sense was never found. The form of sporangial heads of *Ab.sidia* would be classified at least in four classes: I. almost pyriform, II. subpyriform, III. pseudopyriform and IV. never pyriform.

Why such a great discordance has occurred? The author believes it would have been caused by three reasons. At first the mounted specimen in a solution, such as lactophenol, can hardly reveal the true form of mature sporangial heads, because in the solution the outer membrane of sporangium proper of mature head fuses easily and sporangiospores disperse in all directions. Then only young, immature heads are found in full shape which are almost pyriform. Secondly direct observation of sporangial heads of the growth in a Petri-dish and a slant tube can hardly reveal the exact form of the heads because it is difficult to find many heads with sporangiophore both of which lie in a perpendicular direction to the direction of vision through microscope. Besides that, the object lenses are restricted to that of low power (4x, 5x or at times 10x). On the contrary the slide culture utilized by the author (see the description of that in the chapter Materials and Methods in pp. 4-6) allows the easiness to search for the suitable heads with sporangiophores, being both in the perpendicular direction to the vision. Furthermore object lenses with strong power to the 100x for oil immersion can be used. After all the second reason is the fact that the slide culture has not ever applied to the study of *Absidia* before the author as far as he knows. Thirdly the idea of "pyriform" of the mycologists before the author of this report might be rough, loose and ambiguous and have too wide sense. Even if "pseudopyriform" were also thrown into a dust bin, the false pyriform" would be 90.75%, still not 100 or near 100% yet.

Who would consider or name a ball fixed with concave funnel (III Pseudopyriform) or with a trumpet (III Pseudopyriform) pyriform? Japanese pears have not neck are near apple and more different than western pears, then we Japanese know: the "Pyriform" from photos or drawings in atlas and books of the western horticultural of fruit trees. In this report in 3100 heads "Euptriform 0%, I "Almostpyriform" 1.29%

and their shapes II “Subpyriform” 19.49%, III “Pseudopyriform” 59.97% and “Neverpyriform” 19.25%.

The contributions of this report 1. (1) Recommendation and introduction of modern statistics into mycology, (2) Merit of slide culture to the examination in exact measuring and precise observation. In the air natural and right morphology can be determined, though in mount fluid the form especially of that of sporangium proper changes its original form (in the fluid mature sporangium proper fused or broken), but immature head can be seen in original form almost pyriform. It might be the greatest reason, the author estimates, that the examiners had seen this young immature heads and had been misled to the fault idea “head are pyriform” (3) The author proposed a new technical term “sg. Capitulum sporangialis, pl. Capitula sporangiales, sg. sporangial head pl. -s. It is an analogous nomination of “conidial head” of *Aspergillus* including, vesicle, sterigmata and conidial chains. Till now sporangium or sporangia have been used for either for the sac only, or adding only columella, and apophysis a whole with sporangiospores. To indicate sporangium (full of spores) alone should be called sporangium proper. A unit including sporangium proper, columella and apophysis should not be called or written “sporangium” but ought to be called and written, adding head, sporangial head.

ACKNOWLEDGEMENT

The author wishes to express his appreciation to Prof. M. Hinoki, M.D. and Mr. Y. Matsuda, M.D. for getting the strains of *Absidia* from ATCC, CBS and IFO, to Prof. M. Yamamoto, Ph. D. and Prof. K. Iwakuni, Ph. D. to get the copies of pertaining papers, Dr. Y. Mori for calculation with electronic computer, Prof. I. Uesaka, M.D. and Assit. Prof. B. Kuze and Mrs. T. Homma for preparing culture media and Miss S. Date for typewriting of the paper.

The acknowledgment is only here in this Part I, No. 1 described, though for the whole of this serial studies the author must say many thanks to the people mentioned above.

REFERENCES

(The books and literatures described here down are not only for the report 1 but also for all following reports of this serial work on *Absidia*.)

If in any following report, other books and literatures are necessary to be referred, they would be written at the end of the report as supplements)

A. MYCOLOGY

Ainsworth, G.C. ed. 1971. Ainsworth & Bisby's Dictionary of the Fungi. Commonwealths Mycology. Insit. Kew, England.

Ainsworth, G.C. and P.K.S., Austwick 1959. Fungal Diseases of Animals. Commonwealths Agricultural Bureaux, Farnham Royal Bucks England.

Ainsworth, G.C. 1952. Medical Mycology. Sir Isaak Pittman & Sons. L.T.D. London.

Brumpt, E. 1949. Précis de Parasitology. Vol. II, troisième partie, Etude systématique des Champignons parasites et mycoses. 1949, Masson et Cie, Paris.

Bainier, G. 1883a. Observation sur les Mucorinees. Ann. Sc. Nat. Bot. ser 6, 15, 70.

Bainier G. 1889. *Absidia coerulea*. Bull. Soc. Bot. France 36: 184.

Bainier, G. 1903. Sur quelques espèces de Mucorinees Nouvelles ou peu connues. Bull. Soc. Mycol. France 19: 153-172.

Campbell, M.E. 1938. An investigation of Mucorales in the soil. Trans. Roy. Soc. Edinburgh 59: 436-471.

Christenbreg, G.A. 1940. A taxonomic study of Mucorales in southeastern United States. Jour. Elisha Hist. Sci. Soc. 56(2): 333-360.

Chiba, K. 1960. Studies of *Absidia ramosa* isolated from a case of Mucormycosis of the Skin and Nail. Jpn. J. Med. Mycol. 22: 330-345.

Dodge, C.W. 1935. Medical Mycology.

Ellis, J.J. and C.W. Hesseltine 1965. The genus *Absidia*: Globose-sporal species. Mycolog. 57: 222-236.

Clements, F.E. and Shear, C.L. 1975. Genera of Fungi. Hafner Pub. Co. New York.

- Bessey, E.A. 1950. Morphology and Taxonomy of Fungi. Hafner Press. New York, London.
- Gilman, J.C. 1957. A Manual of Soil Fungi. Iowa State Uni. Press, Ames. Iowa.
- Hagem, O. 1908. Untersuchung über norwegischer Mucorineen. I. Christiania Vid. -Selsk. Skr. K.I. 4: 1-152.
- Halay, L.D. 1950. Etiology of Otomycosis. Arch. Otolaryngology 52: 202-204.
- Emmons, C.W., C.H. Benford, J.P. Utz and K.K. Kwon-Chung. Medical Mycology. 1977. Leaf & Febiger. Philadelphia.
- Helene Matterbok and H.F. Schjoler 1974. Taxonomy and identification of mucormycosis causing Fungi I Synonymy of *Absidia ramosa* with *Ab. corymbifera* Sabouraudia. 12: 64-76.
- Hesseltine, C.W. 1952. A survey of Mucorales. Trans. N.Y. Acad. Sci. Ser. II 14: 210-214.
- Hesseltine, C.W. 1955. Genera of Mucorales with notes on their synonymy. Mycologia. 47: 344-363.
- Hesseltine, C.W. and J.J. Ellis. 1961. Notes on Mucorales, especially *Absidia*. Mycologia. 53: 406-426.
- Hesseltine, C.W. and J.J. Ellis. 1964a. The genus *Absidia*: *Gongronella* and cylindero-spored species of *Absidia* Mycology 56: 568-601.
- Hesseltine, C.W. and Ellis, J.J. 1966. Species of *Absidia* with ovoid sporangiospores I. Mycologia. 58: 761-785.
- Hesseltine, C.W. 1954. The section *Genensis* of the *Mucor*. Mycologia. 46: 362.
- Higuchi, K. et H. Urabe 1964. Mycosis (Japanese) Kanehara Publ. Co. Tokyo.
- Klocker, A. 1906. Gärungsorganismen
- Kimura, M. 1959-1960. Horticulture of Fruits-Trees (in Japanese) Asakura-bookstore Tokyo.
- Kominami, K. et K. Tubaki 1952. Enumeration of molds of Japan II Mucorales. Nagaoa Inst. I (21) 192.
- Kominami, K.K. Tubaki, Y. Kobayashi. 1952. Mycology. Nagaoa I 14-25.
- Lafar, A. 1914. Handbuch der technischen Microorganismen
- Lendner, A. 1908. Les mucorinees de la Suisse, Material pour la flora cryptogamic Suisse fasc. 1, 1-180.
- Lindt, W. 1886. Mitteilungen über einige Schimmelpilze Naunyn-Schwiderberg's Arch. für experimentelle Pathologie. 21: 144-177.
- Muskat, J. 1955. Untersuchungen über bayrischer und bunesischer Bodm I. Floristisch-okologischer Teil. Arch. Microbiol. 22: 1-20.
- Ota, M. Champignons parasites de l'homme (Japanese) 1920. Journ. Dermatology u. Urolog. 26: 15.
- Ota, M. Human and animal mycosis and their pathogen fungi. Correspondence of the 74th lectures (Japanese) of Institute of infectious disease. Tokyo Imp. Univ. II 201-274.
- Takahashi, Y. 1952. Technics of examination of human-parasite fungi. Igaku-shoin Tokyo.
- Nakazawa, R. 1913. Über Gärungsorganismen in Formosa. Ber. Ver. Stat. Natw. Formosa. 2-525.
- Orr, G.F. and O.A. Punkett 1959. A New species of *Absidia* from California. Mycology 51: 203-209.
- Oudemans, C.A. et J. Klining 1902. Prodrome d'une Flore Mycologique de la terre humide du Spanderswood.
- Nammylowski, B. 1910. Studien über Mucorineen. Bull. internat. de l'Acad. Sci. de Cracovie, Class. Sci. math. et nat.
- Saccardo, P. 1888. Sylloge Fungorum t-III: 214-215.
- Sartory, A. 1920-1921. Champignon parasite de l'homme et des Animaux Fasc. 2.3. 72-180.
- Sartory, A. 1926. Guide pratique des principes manipulations mycologiques 4-20: 199-255.
- Singer, D.E. et E. Freeman, W.R. Hofferet, R.J. Keys, A.V. Hardy et R.B. Mitchell 1952. Otitis externa. Ann. Otol. etc. 61: 317-330.
- Skinner, C. C.W. Emmons et H.M. Tsuchiya 1947. Henrici's Molds, Yeasts, and Actinomyces. New York. John Wiley & Sons. Inc. London: Chapman & Hall Limited.
- Subramanian, C.V. 1952. Fungi isolated and recorded from Indian soils. J. Madras Univ., 22B: 206-222.
- Talbot, P.H.B. 1978. Principles of Fungal taxonomy. MACMILLAN PRES. LTD.
- Tieghem, P. van 1875. Nouvelles recherches sur les Mucorinees. Ann. Nat. 6 ser. 1: 5-174.
- Tieghem, P. van 1873. Recherches sur Mucorinees. Ann. Sci. Nat. 5 ser. 17: 261-399.
- Udagawa, S., K. Tubaki, S. Watanabe, K. Minoura etc. 1978. Manual and Atlas of Fungi (Japanese). Kodansha Co. Tokyo.
- Vuilleman, M.P. 1903a. Importants taxonomiques de l'appareil zygosporé des Mucorinees. Bull. Soc. Mycol. France 19: 106-118.
- Vuillemin, M.P. 1903b. Le genre Tieghemella et la série des Absidee Bull. Soc. Mycol. France 19: 119-127.
- Vuillemin, P. 1904. La *Lichtheimia remosa* (*Mucor ramosus* Lindt), Champignon pathogène, distinct du *L. corymbifera*. Arch. de Parasitologie 8: 562-572.

Wehmer, C. 1907. Morphologie und Systematik der Mucoraceen. Lafar's Handbook 4: 455-506.

Yamashita, K. 1941. Otomycosis in Formosa Report 1. A clinical study from a statistical standpoint. Acta Japonica Medicina. Tropicalis (Taipei Univ. at present)

Yamashita, K. Otomycosis in Formosa Report 2 *Lichtheimia maligna* (Japanese). Nettai Igaku I: 189-201.

Yamashita, K. 1941. Guide to Mycology (Japanese). Kanehara Publish. Co. Tokyo.

Yamashita, K. 1963. Fungal Flora in the Ear Nose, Throat and Mouth of man. Jpn. J. Med. Mycol. 4: 136-149.

Zycha, H. and R. Siepmann 1969. Mucorales. D-3301 LEHRE VERLAG VCN J. CRAMER

B. STATISTICS

Bishop, O. 1971. Statistics for Biology 2nd ed. O.N. Longman, London.

Campbell, R.C., 1974. Statistic for Biologist. Cambridge Univ. Press.

Fisher, R.A. 1970. Statistical Methods for Research Worker. 14th ed. Hafner Press New York. Collin-MacMillan Publ. London.

Fisher, Sir. R.A. and F. Yates 1975. Statistical tables for Biological Agricultural and Medical

Research, sixth ed. Longman Group Ltd. London.

Hajek, J. 1969. A course in Nonparametric Statistics. (Translated in Japanese by Okamoto M. Y. Miyamoto and K. Kogo, 1974. Nikkagiren Publ. Tokyo.

Ishii, S., 1977. An introduction to Biological Statistics (Japanese) Yokendo Publ. Co. Tokyo.

Kishine, T. 1975. Theories and Application of Statistics (Japanese) Yokendo Publ. Co. Tokyo.

Kitayama, T. and M. Masuyama 1952. New Statistical Tables. Kawadeshobo Publ. Co. Tokyo.

Masuyama, M. 1950. Statistical methods for small samples. (Japanese) Kawade Book Co. Tokyo.

Nakayama, I. ed. 1980. Large dictionary of present Statistics (Japanese) Toyokeizaishimpo. Co.

Siegel, S. 1956. Nonparametric Statistics for the Behavioral Sciences. Mc. Grow-Hill Co. New York, Toronto, London.

Snedecor, G.W. and Cochran, W.G. 1967. Statistical Methods. 6th ed. Iowa State Univ. Press Ames. Iowa.

Tan-naka, T. 1977. Theory and Application of Statistics Yokendo Pbl. Co. Tokyo.

Okugawa, K. 1959. Outline of mathematical statistics. Gakujutsutosho Pbl. Co. Tokyo.

58 5 25 606 58

Explanation of Photos of sporangial heads

1. Subpyriform. Sporangium proper standing subsphenic, transition from sporangium proper to apophysis quite gradual, apophysis irregular convexity, transition from apophysis to sporangiophore quite clear.

2. Almost pyriform. Sporangium proper lying subspherical, apophysis bell-like, transition from sporangium proper to apophysis gradual, transition from apophysis to sporangiophore in left side clear (almost angular near rightangle,) in right side angle-formation.

3. Subpyriform., Sporangium proper truncated spheric, transition from that to apophysis slightly angular, and from the latter to sporangiophore gradual.

4. Subpyriform. Sporangium proper oblique from (from right upper left lower) standing ovoid, right lowest portion of columella exposed, apophysis large cup-like, transition from that to sporangiophore clear.

5. Subpyriform. Sporangium proper lying short ellipsoid, apophysis drinking glass like, transition from that to sporangiophore almost clear.

6. Pseudopyriform. Sporangium proper lying short ellipsoid but prolapsed into apophysis (deep

depressed columella, see chapter three columella in the second report), apophysis long cone, transition from that to sporangiophore with large angle.

7. Pseudopyriform. Sporangium proper lying subspheric, transition to apophysis gradually, that from apophysis to sporangiophore quite gradual, apophysis trumpet-form.

8. Pseudopyriform. Sporangium proper truncated spheric, apophysis bending trumpet, transition from that to apophysis not clear.

9. Never pyriform. Sporangium proper spheric, apophysis uncertain but a nipple from sporangium proper into sporangiophore.

10. Never pyriform. Sporangium proper spheric, apophysis none.

11. Never pyriform L. Sporangium proper truncated subspheric. Transition sporangium proper to apophysis almost clear and that from apophysis to sporangiophore, apophysis moderately long cone. R.S. sporangium proper standing subspheric, apophysis none or uncertain, sporangiophore club-shaped, never pyriform

12. Never pyriform. Sporangium proper leaf-form, apophysis none