STOCHASTIC MODEL OF GLUCOSE TOLERANCE TEST

Motosaburo MASUYAMA

(Department of Applied Mathematics, Science University
of Tokyo)

SUMMARY

In the glucose tolerance test there holds an empirical law that $t_m < t_v$, where t_m and t_v denote respectively the times when the maxima of inter-person mean and variance of its concentration in blood are attained.

Let the stochastic version of classical equations of metabolic turnover be

$$\begin{aligned}dD_t &= -KD_t dt + B_1 D_t dW_1, \\dC_t &= (-kC_t + KD_t + a)dt + (B_2 C_t + b)dW_2 - B_1 D_t dW_1,\end{aligned}$$

where B_1 , B_2 and b are positive constants, and $W_i (i = 1, 2)$ are two mutually independent standard Wiener processes. We deduce the law under the plausible assumptions;

$$\begin{aligned}K &\gg B_1^2, \quad k \gg B_2^2, \quad K > k, \\ \sigma_{110} &\ll D_0^2, \quad \sigma_{120} = 0, \quad \sigma_{220} = \sigma_{22\infty},\end{aligned}$$

where (σ_{ij}) denotes the variance-covariance matrix of $(D_t, C_t)'$.

The proof is given for three disjoint cases, i.e. (i) $k < K < 2k$, (ii) $K = 2k$, and (iii) $2k < K$.

See the reference:

Masuyama, M. (1980): A stochastic model of glucose tolerance test, *TRU Mathematics*, 16-1, 83-88.