A REALISTIC MODEL FOR PREY-PREDATOR POPULATION INTERACTIONS AND THE KOLMOGOROFF-SAMUELSON CONDITIONS

K. OHTA
(Tokyo Keizai University, Kokubunji, Tokyo)

SUMMARY

A realistic model is proposed for prey-predator population interactions. Although giving a stable limit cycle in a certain case, the model does not necessarily meet one of the conditions given by Kolmogoroff. The condition is such that its necessity was doubted of by Samuelson.

1. Introduction: Lotka and Kolmogoroff

To my knowledge (Ohta, 1981b) Lotka (1923) was the first to suggest a possibility of a stable limit cycle for eaten-eater population interactions. Later Kolmogoroff (1936) proposed his famous conditions for either a stable equilibrium point or a stable limit cycle for a general prey-predator system. Since then a number of authors have accepted the Kolmogoroff conditions by adapting them to their specific models of prey-predator populations.

KEY WORDS: Prey-predator; Kolmogoroff-Samuelson conditions

2. Samuelson's Doubt

Samuelson (1967), however, has doubted of the necessity of one of the Kolmogoroff conditions. It is the condition

$$N_1 \frac{\partial f_2}{\partial N_1} + N_2 \frac{\partial f_2}{\partial N_2} \geq 0 \quad (1)$$

where N_1 and N_2 are the densities of the prey and predator populations respectively, and they are assumed to obey the differential equations

$$\dot{N}_1 = N_1 f_1(N_1, N_2) \quad (2)$$

$$\dot{N}_2 = N_2 f_2(N_1, N_2).$$

Mathematically speaking, (1) may be derived from the condition

$$\frac{df_2}{dr} \geq 0 \quad (3)$$

where r is the radius in polar coordinates of the point (N_1, N_2) . Biologically speaking, the condition (1) or (3) means that the per capita growth rate of the predator population does not decrease as both of the prey and predator populations increase in proportion.

3. A Realistic Model

As Samuelson (op. cit.) suggests, however,

the condition (1) or (3) seems not necessarily the case with some realistic systems. Actually, for the system

$$\begin{aligned}\dot{N}_1 &= [r_1(\frac{K_1-N_1}{K_1}) - \frac{c_{12}N_2}{N_1+\alpha_{12}}]N_1 \\ \dot{N}_2 &= [r_2(\frac{K_2-N_2}{K_2}) - \frac{c_{21}}{N_1+\alpha_{21}}]N_2\end{aligned}\tag{4}$$

(1) becomes

$$\frac{c_{21}N_1}{(N_1+\alpha_{21})^2} - \frac{r_2N_2}{K_2},$$

which is not necessarily positive, although (4) has a stable limit cycle for a certain case (Ohta, 1981a).

References

- Kolmogoroff, A. N. (1936). Giorn. Inst. Ital. Attuari 7:74-80 (Italian).
- Lotka, A. J. (1923). J. Wash. Acad. Sci. 13:152-158.
- Ohta, K. (1981a). J. Hum. Nat. Sci. (Tokyo Keizai Univ.) 58:129-146.
- Ohta, K. (1981b). Ibid. 59:65-71.
- Samuelson, P. A. (1967). Oper. Res.-ver. 3:307-320.