後頭・頭頂部の事象関連脳活動の変化と 心的イメージ操作パフォーマンスとの相関

○岩木 直, 須谷康一 (産業技術総合研究所ヒューマンライフテクノロジー研究部門)
稲川三千代 (京都大学経済研究所), 飛永芳一 ((株)エレガフィ)
西村和雄 (京都大学経済研究所)

Event-related changes in the gamma-band activity reflect individual performance in mental rotation
Sunao IWAKI, Kouichi SUTANI (National Institute of Advanced Industrial Science and Technology),
Michiyo INAGAWA (Kyoto Univ.), Yoshikazu TOBINAGA (Elegaphy, Ltd.),
Kazuo NISHIMURA (Kyoto Univ.)

1. はじめに

Mental image processing, including rotating and matching of a pair of 3-D objects, activates human parietal and extrastriate visual cortices [1][2]. It is also reported in the recent electrophysiological studies that the gamma-band spontaneous brain activities in these regions play a crucial role in manipulating mental representation of 3-D objects [3], however, it is not fully characterized whether individual changes in the gamma-band activity are correlated with the individual difference in the performance of mental image processing. In this study, we measured MEG responses to investigate neural correlates of the individual variability in the performance of 3-D mental image processing.

2. 方法

Fifteen subjects performed mental rotation task in which a pair of either (a) identical or (b) mirror-reversed 3-D objects with 4 different rotation angles (0, 30, 60, 120 deg) were presented (Fig. 1). Subjects' task was to discern whether presented pair of objects was either identical or mirror-reversed and to react by lifting their index of middle finger to measure percent correct of the task performance. MEG signals were measured using whole-cortex type SQUID system with 122 planar gradiometers. The stimulus related epochs of 2 s including 0.5 s pre-stimulus baseline, were recorded with a sampling rate of 600 Hz. More than 50 epochs were averaged for each condition.

Root-mean-square (RMS) values for 5 different MEG sensor groups covering the primary visual area, the left and right posterior inferotemporal areas, the left

and right superior parietal areas, each of which contains 4-8 sensors, were calculated to evaluate local signal power in each condition. Event-related time-frequency (TF) wavelet decomposition of the recorded MEG raw signals between 5 – 60 Hz was used to assess the event-related changes in the spontaneous brain activities (event-related synchronization (ERS)/ event-related desynchronization (ERD)) in response to the

Fig. 1 Examples of the mental rotation stimuli used in this study. (a) Identical or (b) mirror reversed pair with 4 different rotation angles were presented onset of mental rotation stimuli.

3. 結果

Based on the measured performance (percent correct of the subjects' response averaged over all conditions, subjects were divided into two groups, namely, (A) high-performers (HP) and (b) low-performers (LP).

Results of the MEG-RMS analysis (Fig. 2) showed that the activities in the primary visual area were significantly increased in the low-performer (LP: gray) compared to the high-performer (HP: black) at around 200 ms after the onset of the stimuli.

Figure 3 shows the results of the MEG time-frequency analysis, which indicate that the gamma-band activities around 30 Hz measured at the occipital and the parietal sensors were significantly increased in the HP compared to LP.

Fig. 2 MEG RMS waveforms averaged separately over HP (black lines) and LP (gray lines) subjects at the primary visual area.

4. 考察

Previous fMRI and MEG studies on mental rotation task indicated that activities in the superior parietal regions are crucial in mental image manipulation [1][2]. On the other hand, there is converging evidence that neural processes involved in binding visual features are accompanied by, or systematically related to, gamma-band spontaneous activities [4][5].

The current results show significant increase in gamma-band power at primary visual area and the superior parietal area in the high-performers, which suggests that the efficient binding of visual features for mental image manipulation in these regions leads to better performance in the mental rotation task.

Increase in the event-related responses observed in the primary visual area in the low-performers might indicate the subjects' dependence on the presented visual object in performing mental rotation task

参考文献

- 1) Cohen MS et al., Changes in cortical activity during mental rotation – A mapping study using functional MRI –, Brain, vol. 119, pp. 89-100, 1996.
- 2) Iwaki S et al., Dynamic cortical activation in

Fig. 3 Results of the TF group statistics of the event-related changes in the spontaneous brain activities at the occipital and superior parietal areas. Thick color indicates significant increase in the spontaneous activity in the HP compared to the LP

mental image processing revealed by biomagnetic measurement, Neuroreport, vol. 10, pp. 1793-1797, 1999.

- 3) Martinovic J et al, Induced gamma-band responses predict recognition delays during object identification, J Cogn Neurosci, vol. 19, pp. 921-934, 2008.
- 4) Engel AK et al, Temporal binding and the neural correlates of sensory awareness, Trends Cogn Sci, vol. 5, pp. 16-25, 2001.
- 5) Wyart et al, Neural dissociation between visual awareness and spatial attention, J Neurosci, vol. 28, pp. 2667-2679, 2008.