特別講演

The Relevance and Adaptations of the Neuromuscular System in Jaw Deformities

~The Role of Muscles in Reconstructive Surgery
Applied to the Craniomandibular Skeleton?~

Arthur J. Miller, Ph.D.

Professor, Department of Growth and Development, School of Dentistry,
Department of Physiology, School of Medicine, UCSF

Curriculum Vitae

Dr. Miller obtained his doctorate at the University of California at Los Angeles in 1975 with a specialty in neurophysiology. He was a trainee of the Brain Research Institute and his thesis involved studies in the brain stem control of laryngeal reflexes and deglutition. After five years as an assistant professor in the Department of Physiology, University of Illinois, he joined the School of Dentistry at the University of California at San Francisco. He is now a professor in the Department of Growth and Development and holds a joint appointment in the Department of Physiology, School of Medicine, UCSF.

Muscle is a unique tissue of the body in its ability to develop tension. The muscle force modifies bone by altering its shape (i.e., twisting and bending the mandible as measured by strain) as well as generating stress within the bone infrastructure. The force developed by the muscle depends upon its size (i.e., cross-sectional area). The mandibular jaw-closing muscles are multipennate muscles with multiple compartments of fibers recruited at different times during incising, swallowing, and chewing. The forces generated by the mandibular muscles effect the shape and size of the mandible and the craniofacial skeleton based on several studies. Lack of development of mandibular muscles in a particular mouse with a gene defect still results in development of the entire craniofacial skeleton and cartilage but the cartilage is smaller and the bone (i.e., mandible) are different shapes. The fetal craniomandibular skeleton needs normal intact muscle to develop its normal shape. During early postnatal development, altering the bilateral control of jaw muscles by partially destroying the motoneurons to the ipsilateral temporalis, masseter, medial pterygoid, and lateral pterygoid muscles, induces the development of an asymmetrical mandible and entire craniofacial skeleton. Imbalance in mandibular muscle recruitment in the rapidly growing craniomandibular skeleton will alter its shape and size.

Evaluating patients with jaw deformities and surgically correcting these deformities with long-term stability depends upon understanding the interaction of the jaw and facial muscles with the underlying bone. The surgically-corrected mandible must be able to meet two criteria: 1) provide functional forces through incising, chewing, and swallowing that prepare food for digestion; and 2) relate to the surrounding muscles so that forces generated on the mandible are harmonious with the bone shape and size. Bone will adapt its shape and size if its new position within the enveloping muscle means the muscles develop different directions of forces and different levels of force. Experimental work by one oral surgeon in which the origins of the temporalis and masseter muscles were removed from their original sites and placed along the posterior ramus in young growing monkeys altered the normal growth of the maxilla and mandible. Changing the vector and direction of force developed by the jaw-closing muscles in the early postnatal period modifies and reverses the normal maxillary and mandibular development. In humans developing mandibular asymmetries, the underlying cause may not initially be the muscle recruitment. However, the correction of that deformity will depend upon the muscle recruitment and their generated forces in the newly-positioned mandible. If the newly-positioned mandible undergoes forces not compatible with the shape and size of the mandible, the mandibular bone will alter. Finally, another major factor is involved and needs to be considered in reconstructive surgery. Work with several colleagues using quantitative computed tomography (QCT) has now established that the mineralization of the cortical bone in the mandible may serve as a diagnostic tool in evaluating the forces of muscles on bone. Work in over 30 subjects with asymmetrical mandibles shows that the highest mineralized bone is significantly different than subjects with symmetrical mandibles. The subjects with asymmetrical mandibles demonstrate patterns of mineralization suggestive of asymmetrical muscle forces that do not represent the bilateral symmetry of the normal jaw. Experimental studies in animals altering their development of muscle forces shifts the normal bilateral symmetry of mineralization in the cortical bone of the mandible. Mineralization may represent the long-term (i.e., many months) repeated uses of a muscle in a particular mode or recruitment pattern.