		毛の基部は黒色であるので、嘴で胸をかきまわした時には中央に巨大な暗色斑が現われる。(註 これは後に標本により確められた) 胸側には時々暗色横斑が現われ、印象的である。
飛 翔	稍波状と思える飛び方もするが、ひらひら飛ぶことが多い。翱翔をよくする。	常に波状に飛び (タヒバリの如き完全な波状ではない) ヒバリに似ていない。着陸直前の動作や形態 (風切羽外縁に淡色帯) はヒバリに似る。翔翔は一度も観察出来なかった。
地 鳴	ビュルル、ビュルルまたはチュリー、チュリーと鳴く、チー、チーまたはキー、キーと悲しそうに鳴くこともある。(一般に liquid)	チャッチャッチャッチャッまたはビュッビュッビュッビュッと聞える鳴き方をするのが常で (dry な撥音) 一波状毎に一言づつく鳴ようである。この他に弱いつぶやきを耳にしたが表記に至らなかった。

参照文献 山階芳麿 1939 鳥 10 (4): 472. 清棲幸保 1952 日本鳥類大図鑑 (I) pp. 113~114. (1958. 12. 18. 記)

A short-toed Lark, *Callandrella cinerea puii*, was collected at R. Sakawa, Kanagawa, 6 May, 1958. From Japan only two specimens are known from Sado I. Its behaviour and identification are compared with the Skylark, *Alauda*.

巣箱の自動食物記録写真装置 A device of an auto-cinematic food-recorder 蠟山 朋雄 Tomoo Royama

In the summer of 1959, attempts were made to assess the food brought to the young of birds which nest in nesting boxes by means of an automatic photography. The experiments were carried out with a pair of Great Tits (*Parus major*) at the Tokyo University Forest in Fuji, Yamanashi, Prefecture. The present paper deals only with the description of the apparatus illustrated by several specimens of the "photo-records." The results of detailed analysis of the full records will be presented in a separate paper (Royama, in preparation). I am much indebted to Professor M. Nitto who made these experiments possible.

A diagram of the apparatus and its circuit system is shown in Fig. 1. A 16 mm cine-camera with one-frame shot mechanism and a speed light are placed at the back of a nesting box. An electro-magnet E_1 is placed before the camera and releases the shutter first by pulling down on iron-bar I which in turn pushes the

shutter button *S* up through a cam *C*. A double set of perch contacts is attached to the entrance of the nesting box which is connected with the circuit of *E*₁. When a bird thus enters into the box and presses the inner perch down, the circuit of *E*₁ is closed and the shutter is released. The perch is, however,

Fig. 1. A diagram of the apparatus. *E*₁: electro-magnet (operated in 100V A.C.), *E*₂ and *E*₃: electro-magnets (operated in 12V D.C.), *A*: iron-arm with a silver contact, *Sp*: silver plate, *St*: solenoid trigger, *C*₁ and *C*₂: set of silver contacts, *F*: speed light, *S*: shutter button, *C*: cam, *I*: iron-bar, *G*₁ and *G*₂: glass plates, *G*₃ frosted glass plate *W*: wrist watch (for details see in the text).

pressed down not only at every arrival and departure but it is quite likely that while the bird is on the perch, there may be several contacts on account of a restless movement of the bird. Accordingly if the set of perch contacts is directly connected with *E*₁ the shutter would be released at every contact. This would simply mean waste of films, an over-loading for the shutter-releasing mechanism and the speed light, and probably disturb the bird. In the present apparatus the automatic shutter-releasing mechanism is so contrived that the shutter release is to be made only when the bird gets into the box but not when it gets out and, further, only the first contact at the inner perch is to release the shutter but the further contacts, if any within one visit, are not.

Between a pair of electro-magnets *E*₂ and *E*₃ an iron arm *A* is set which swings freely to either direction, to the top of which a silver contact is attached. When a bird arrives and presses the outer perch down, then the circuit of *E*₂ is closed and this causes *E*₂ to pull *A* toward it, thus bringing *A* into contact with a silver

plate Sp. Then the bird comes into the box and presses the inner perch down, closing the circuit of the solenoid trigger St (A and Sp consist of a part of the circuit). Now St pushes a silver contact C_1 toward another contact C_2 . The contact between C_1 and C_2 closes the circuit of E_1 which releases the shutter. At the same time, however, E_3 which is also connected with the inner perch contact pulls A toward it so that A is separated from Sp. Therefore the circuit of St is kept closed only during the moment at which A is sliding on, and keeping contact with, Sp. Thus St is not to operate again unless the bird presses the outer perch down to bring about a recontact between A and Sp, this being made when the bird leaves the nest or makes the next visit.

The speed light placed at the back of the camera is operated synchronously with the shutter release (X-contact is built into the shutter mechanism) and the light reflected at the back of the roof of the box (lined with a sheet of white paper) illuminates the bird at a right position on the inner perch. Thus a front portrait of the bird carrying food in its beak is taken through a glass window at the back wall of the box. The lens hood prevents the light from getting directly into the lens. The camera and the shutter-releasing mechanism are placed in a small wooden case which muffles a mechanical noise. The speed light illumination is defused by a sheet of thin white paper and a plate of frosted glass G_3 . 100V A.C. is used for the operation of the speed light and the shutter-releasing electro-magnet E_1 and 12V D.C. (dry batteries) for the other circuits. A wrist watch was also attached to the wall of the box just by the entrance to record the time of arrivals. The detailed photographic data are:

Camera: 16mm cine-camera with one-frame shot mechanism; Lens: $F=25\text{mm}$, $f=1:3.5$ Hexar lens; Lens stop: $f=1:8$; Film: Fuji Neopan 16mm reversal (ASA 40); Speed light: power output= 37 wat/sec at 100 V A.C., duration= $1/500\text{ sec}$, and interval= 5 sec .

The nesting box is detachable (see Photo 3 in Plate 8) so that several boxes of the same type were provided in the forest separately and to the one which happened to be occupied by a pair of the birds the recorder was attached just before the young hatched. The pair of Great Tits with which the present experiments were carried out seemed not to be disturbed at all by this recorder; they did not show any sign of being terrified at the noise and the light and its visiting frequency was quite a normal one as compared with that of the other pairs that were under observation at the same time by another type of nest-recorder (Royama, in press).

The birds, having got used to the apparatus, sometimes flew into the box without pressing down the perch and thus escaped recording, therefore a constant care was taken for the adjustment of the apparatus.

The identification of the food recorded on the films is not necessarily an easy matter, but assessment is possible to such an extent as to tell what kind of food they are, e.g., caterpillars, moths, spiders, or snail shells. More precise information will be obtained if, at the same time, direct sampling methods as practiced by Kluijver (1950) (collaring method) or by Betts (1955) (artificial gape method) are employed. A quantitative analysis is also possible from the "photo-records." One of the great advantages of this automatic recorder is that it provides us with a large amount of data with a very little field labour.

References Betts, M. M. 1954 Experiments with an artificial nesting. British Birds, 47: 229-231. Kluijver, H. M. 1950 Daily routines of the Great Tit. Ardea, 38: 99-135. Royama, T. (in press) Test of an automatic nest-recorder. British Birds.

Institute of Forest Zoology, Faculty of Agriculture, University of Tokyo.

要約 1959 年の夏、巣箱を利用する鳥のヒナに与える食物を記録する自動写真装置を試作した。実験は 1 番のシジュウカラ (*Parus major*) について、東京大学富士演習林 (山梨県) 内のカラマツ林中で行った。本報告は装置の記載および数枚の写真記録をかかげるにとどめ、全記録 (数千枚) の分析結果は、別文 (準備中) で述べる予定である。

装置のメカニズム (回路図を含む) を第一図に示す。巣箱の後側に 1 コマ撮影のできる 16 ミリ・シネカメラを置き、巣箱の後壁に設けられたガラス窓をとおして巣穴に首を差し入れた鳥を正面から撮影する。カメラの後にとりつけられたストロボは、シャッターが全開した瞬間に光を発し、その光は白紙で内張りされた巣箱の屋根裏に反射して鳥を照す。シャッターは電磁石 (E_1) によって切られる。巣箱の入口には 2 組の接点付トマリ木がとりつけられており、鳥が外側のトマリ木にとまると電磁石 (E_2) が鉄のうで (A) をひきつけ、銀板 (Sp) に接続する。次に鳥が中に入り内側のトマリ木にとまると、ソレノイド (St) の鉄芯が接点 (C_1) を押し、(C_2) に接触させる。そこで E_1 の回路 (100V A.C. を使用) が閉じ、シャッターが切られる。一方、同時に、内側トマリ木の接点は電磁石 (E_3) に接続しているのもので、 E_3 は A をひきつけ、St の回路はただちに切断される。このようにすれば鳥が外から中に入る時、ただ 1 回だけ、露出が行われることになり、鳥が外へ出る時やまた内側トマリ木の上でとびはねるたびに露出が行われるのを防ぐ。従って鳥が一度外に出て外側トマリ木を押し再び A を Sp と接触させない限り次の撮影は行われない。これによってフィルムの無駄とシャッター装置の過荷重を防ぐことができる。

Plate 9 に示した参考写真は 16 ミリ反転フィルムからの複製である。これだけでも食物の判定は、蛾の幼虫、成虫、クモ、カタツムリの殻程度の分類で行うことが出来るが、直接観察を併行すれば、更に細かい判定が可能である。量的な測定もまた可能である。

(東京大学農学部森林動物学教室)

留 鳥 の 群 性 (抄録と観察) Flocking behaviour of resident birds (Reviews and observations) 黒田 長 久 Nagahisa Kuroda

1. ワタリガラスの群性 G. A. J. Schmidt (Orn. Mitteilungen 9: 121-126, 1957) によると Schleswig-Holstein 地方で年を通じて群をなすのは第 1 年, 2 年若しくは第 3 年目の亜成鳥である。これらは頸部や雨覆の褐色なことで成鳥と区別がつくからである。そして番をなしていない。一方番をなす成鳥は周年なわばりをもっていて、それを出ないで暮す。亜成鳥群には秋から初冬にかけて求愛飛翔をみるが、多分第 2-3 年目の鳥であろう。亜成鳥群の採食場や葦の好条件の所には、特に厳冬は、成鳥も群に入ってくることもある。Schleswig-Holstein の三分の二は営巢地に適している。1953 年には 15,644 平方キロに 200 番が営巢した。1953~6 年の間狩猟家がその 25% を駆除した所、亜成鳥群は広く漂行するようになり、何年もワタリガラスの居なかった地方にも現われ、最近 Hamburg 及び Steinburg に侵入してきたもののような亜成鳥群であった。(Bender 氏抄録より)。

2. スズメの群性 日本に於けるスズメの群性がこれとよく似ていると思われる。(但しスズメでは多くは第 1 年の秋から繁殖のため pair formation にかかると思われる) 私はムクドリを観察の際にスズメがムクドリと合同 (両者は別群をなすが) の大群集団で眠る場所をいくつかみ、スズメもムクドリと同じように少なくとも 10 km は屋の採食地から時に飛ぶのを観察した。ところがその群が飛去った後にも附近の村には、そここの家の庭木に (多少は群れることもある) 分れて眠るものがあるのである。各々の例は何れまとめて報じるつもりであるが、面白いことに、愛知県林務課の池田技師は同県の所謂葦の「地獄網」で大量捕獲されるスズメは皆幼鳥で親は一羽もいないと報ぜられ (静岡県猟友会報, 第 11 号, 16 頁, 1958), 又中村司氏も同じ観察をされたという。つまり秋の稲田に群れるものは幼鳥群が主体で、附近の農家の成鳥は屋だけその採餌群に加わるのであろう。さすれば、これはドイツのワタリガラスと全く同じ型の留鳥の群性といふことができる。

なお、白附憲之氏は第 29 回日本動物学会 (松山) に於てスズメのなわばりにつき研究発表され、Summers-Smith (Ibis, 100: 190, 1958) もイエスズメのなわばりを研究した。又 Verheyen (Gerfaut 47: 161, 1957) の標識による移動性の研究などがある。

3. 群性となわばり性 Summers-Smith によると成鳥も含むイエスズメの巣立採食集団は冬までつづくが、成鳥は繁殖地 (なわばり) にもどり、若鳥はパーティをなして

Photo 1. The outside-view of the "Auto-cinematic food-recorder." Photo 2 & 3. The inside-views showing the mechanical construction of the recorder. Photo 4. The mechanism of the waste-exposure prevention (for details see in the text).

The specimens of the "photo-records". Photos 1-4: caterpillars, 5-8: moths, 9-11: spiders, 12-13: snail shells, 14-15: grits (the snail shells are seen included). The male and the female can be distinguished by their plumage patterns (photos 8, 13 and 14: the female). The wrist watch shows the time of arrivals (the photographs are not arranged in the order of succession).