如く他の地方にも相當に得らるゝ所より見れば此の體格の大なると云ふ事は二次的のもので標本となつた鳥は何か極めて良い條件の本に育つた鳥とも見る事が出来る。

猶採集者の言によれば二月も金北山麓の金澤村附近にてトキが二羽飛で居るのを見た。土地の人の話では其の附近で蕃殖する由である。又渡島の途中の海ではウ、カモメ(種不明)、ウミスズメ、アジサシ(種不明)等を見たと云ふ事である。

On the Specimens of Korean Birds, collected by Mr. Hyojiro Orii.

By

Marquis Y. Yamashina.

Mr. Hyojiro Orii stayed in Korea (including the Quelpart Is.) for one year following April, 1929, and was engaged in the collection of birds, collecting 1940 specimens which belong to 41 families and 279 forms and some eggs. This collection is one of the most perfect ever brought from Korea. We have gotten three new subspecies among this collection. Besides, we have been able to add twenty-six species to the list of Korean birds (six of them also to the Japanese list). I offer my hearty thanks to him in appreciation of his services.

The itinerary of his collecting trip is as follows. (See the attached map.)

He started from Tokyo in the last decade of March, 1929.

From April 2nd, 1929, to June 4th, 1929, he sojourned at Riugampo, Heian-hokudo, chiefly collecting birds passing there in spring.

From June 11th, 1929, to July 18th, 1929, he sojourned at Chōsen, Kogendo, engaged in the collection of birds in the forests around Mt. Kongo. He collected a new subspecies of Pigmy Woodpecker during this period.

Fig. 29. Korean Peninsula.

From July 23rd, 1929, to September 6th, 1929 he sojourned at Nojido, Kankyo-hokudo, engaged in the collection of birds in the whole neighbourhood of Mt. Hakuto. He collected new subspecies of Three-toed Woodpecker and Lesser Spotted Woodpecker (see the "Tori" No. 29), and in addition, a great number of birds which have some relations to ones in the Ussuri region, and illustrated the fact that Fauna in Kankyo-hokudo differ from those in the other parts of Korea.

After leaving Nojido on Sept. 6th, 1929, he made a collection of some

birds at Sanchō on his way, and then went to Seishin by way of Mozan, and farther northward to Yuki.

From September 12th, 1929, to November 21st, 1929, he sojourned at Bampo, Kankyo-hokudo, engaged in the collection of birds passing there in autumn. Jankowski's Bunting was collected during this period.

From November 25th, 1929, to December 6th, 1929, he occupied himself in the collection of birds at Kinkwa, Kogendo.

From the 7th of December, 1929 to the 7th of March, 1930, he sojourned at Moppo, Zenra-nando. He went to Chitō several times to collect birds.

On March 8th, 1930, he crossed over the Quelpart Is., staying there till March 31st, 1930, collecting birds.

He returned to Tokyo in the first decade of April, 1930.

About some of the specimens collected about this time, I have already described at the "Tori" No. 28, No. 29 and No. 30. And now again, I have given all the specimens, including them, in the table I have made, giving some explanations where I think, it is necessary to do so.

Family CORVIDÆ. 鴉 科

1. *Corvus corone corone* Linnaeus. ヘシボソガラス

Synonym: *C. c. interpositus* Laubmann.

C. c. takahashii Momiyama & Ishii.

2 ♂♂ ads. 10.-11. IV. 1929. Riuganpo.

1 ♂ ad., 1 ♀ ad. 3.-21. X. 1929. Banpo.

2 ♂♂ ads., 4 ♀♀ ads. 6. I.-25. II. 1930. Moppo.

The results of measurement of 5 ♂♂ ads. and 5 ♀♀ ads. above mentioned are as follows:

Wing ♂♂ ads. 312-345, ♀♀ ads. 302-328.

Tail ♂♂ ads. 172-200, ♀♀ ads. 166-213.

Culm. ♂♂ ads. 49-54, ♀♀ ads. 48-52.5.

The results of measurement above mentioned coincide with the description of *C. c. corone* in the "Practical Handbook of British Birds" by Mr. Witherby, wing ♂ 318-345, ♀ 310-325. And the results of measure-

ment of a great number of specimens of Hokkaido, Hondo and Kiusiu are as follows:

Wing ♂ 311-347. ♀ 309-340.

Thus, those in Hokkaido, Hondo and Kiusiu are undistinguishable from those in Europe. So the method used by Dr. Hartert and Dr. Laubmann to classify *C. corone* into the western and the eastern form is wrong. Those in Saghalin have the measurement, wing 12 ♂♂ ads. 330-367, 7 ♀♀ ads. 324-336 mm. and are of a larger size than *C. corone corone*. They belong to *C. c. orientalis* Eversmann and *C. corone saghalense* Kumagai is its synonym.

2. *Corvus coronoides mandshuricus* Buturlin.

テウセンハシブトガラス

Synonym: *C. c. quelpartis* Momiyama.

1 ♂ ad. 22. VIII. 1929. Nojidō.

1 ♀ imm. 3. X. 1929. Banpo.

1 ♀ ad. 28. XI. 1929. Kinkwa.

4 ♂♂ ads., 1 ♀ ad., 1 unsexed ad. 17.-30. III. 1930. Quelpart Is.

Those in the Quelpart Is. have a little longer bills than those in Korea, but the difference is not great. The results of measurement of *C. coronoides* in Korea and the Quelpart I. are as follows:

	Wing	Ex. c.
Quelpart Is. ♂ ♀	295-348	52.5-61
Korea ♂ ♀ ads.	317-344	56-64

3. *Corvus frugilegus pastinator* Gould. ミヤマガラス

1 ♂ juv. 21. X. 1929. Banpo.

5 ♂♂ ads., 3 ♀♀ ads. 2. I.-26. II. 1930. Moppo.

1 ♀ ad. 13. III. 1930. Quelpart I.

4. *Corvus dauricus dauricus* Pallas. コクマルガラス

1 ♂ ad. 21. X. 1929. Banpo.

3 ♂♂ ads., 7 ♀♀ ads. 25. XII. 1929-1. III. 1930. Moppo.

5. *Pica pica sericea* Gould. カササギ

Synonym: *Pica varia japonica* Schlegel.

2 ♂♂ ads. 19. IV.-17. V. 1929. Riuganpo.

- 1 ♂ ad., 3 ♀ ♀ ads. 28. VII.-12. VIII. 1929. Nojidō.
 2 ♂ ♂ ads., 6 ♀ ♀ ads. 24. IX.-23. X. 1929. Banpo.
 4 ♂ ♂ ads., 5 ♀ ♀ ads. 17. I.-27. II. 1930. Moppo.

According to the recent investigation of the specimens preserved in the collections of Takatsukasa, Kuroda and Yamashina, there is no distinction among the specimens of Formosa, Korea and Kiusiu, neither in measurement nor in coloring.

6. *Cyanopica cyanus interposita* Hartert. コマヲナガ
 7 ♂ ♂ ads., 3 ♀ ♀ ads. 21.-26. II. 1930. Moppo.
 7. *Garrulus glandarius taczanowskii* Lönnberg. カラフトミヤマカケス
 Synonym: *G. brandti okai* Momiyama.
 1 ♂ ad., 1 ♀ ad., 2 juvs. 9. VIII.-3. IX. 1929. Nojidō.

G. glandarius of Korea in many specimens are almost undistinguishable from *G. g. taczanowskii* of Saghalin. *G. g. pallidirostris* of Hokkaido, on the contrary, can be easily distinguished from *G. g. taczanowskii* of Saghalin. It is unable to determine whether *G. g. taczanowskii* are distinguishable from *G. g. brandti* or not, as there have never been found the specimens of the Altai region.

Family STURNIDÆ. 椋鳥科

8. *Spodiopsar cineraceus* (Temminck). ムクドリ
 3 ♂ ♂ ads., 3 ♀ ♀ ads. 11. IV.-28. V. 1929. Riuganpo.
 1 ♂ ad. 28. VII. 1929. Nojidō.
 9. *Sturnia sturnina* (Pallas). シベリヤムクドリ
 6 ♂ ♂ ads., 4 ♀ ♀ ads. 18.-27. V. 1929. Riuganpo.

Family ORIOLIDÆ. 黃鳥科

10. *Oriolus chinensis diffusus* Sharpe. カウライウグヒス
 Synonym: *O. i. ochroanthus* Oberholser.
O. chinensis swinhoii Momiyama & Ishii.
 3 ♂ ♂ ads., 2 ♀ ♀ ads. 11.-23. V. 1929. Riuganpo.

Taking many specimens, there is no distinction between those of Formosa and those of Korea neither in measurement nor in coloring.

Family FRINGILLIDÆ. 雀科

11. *Coccothraustes coccothraustes verticalis* T. & B.

カラフトシメ

2 ♂♂ ads., 1 ♀ ad. 27. X.-12. XI. 1929. Banpo.

3 ♀♀ ads. 25. XII. 1929-27. I. 1930. Moppo.

Those which live in Korea in winter belong to this subspecies. There is one more specimen beside those above mentioned.

1 ♂ juv. 28. VII. 1929. Nojidō.

This specimen is one which has just left its nest, so its subspecific name can not be ascertained. But, I have provisionally treated it as belonging to this subspecies.

12. *Coccothraustes coccothraustes japonicus* T. & S. シメ

1 ♀ ad. 17. III. 1930. Quelpart Is.

13. *Eophona migratoria migratoria* Hartert. コイカル

6 ♂♂ ads., 5 ♀♀ ads. 9.-20. V. 1929. Riuganpo.

2 ♀♀ ads. 24. II. 1930. Moppo.

14. *Chloris sinica ussuriensis* Hartert. テウセンカハラヒワ

6 ♂♂ ads. 4 ♀♀ ads., 1 juv. (27. V.) 4. IV.-29. V. 1929. Riuganpo.

1 ♀ ad., 2 juvs. 15. VI.-13. VII. 1929. Chōsen.

5 ♂♂ ads. 25.-26. XI. 1929. Kinkwa.

2 ♂♂ ads., 1 ♀ ad. 7.-27. I. 1930. Moppo.

C. s. ussuriensis is a common resident in Korea.

15. *Chloris sinica minor* (T. & S.) コカハラヒワ

Synonym: *C. s. affinis* Momiyama.

4 ♂♂ ads., 1 ♀ ad. 14.-30. III. 1930. Quelpart I.

Chloris sinica which is a resident in the Quelpart I. is given the name of *Chloris sinica affinis*, but it is undistinguishable from *C. s. minor* of Hondo.

16. *Carduelis spinus* (Linnaeus). マヒワ

3 ♂♂ ads., 2 ♀♀ ads. 22. X.-11. XI. 1929. Banpo.

2 ♂♂ ads., 3 ♀♀ ads. 17. I.-23. II. 1930. Moppo.

17. *Carduelis linaria linaria* (Linnaeus). ベニヒワ

1 juv. 17. XI. 1929. Banpo.

About this Specimen, I have described in the "Tori" Vol. VI. p. 257, that it belongs to *C. hornemanni exilipes* in accordance with the classifica-

tion of Dr. Hartert, but, following the classification of Dr. F. Solomonsen, it belongs to the subspecies above mentioned.

18. *Uragus sibiricus ussuriensis* Buturlin. ラナガベニマシコ

1 ♂ ad. 12. VIII. 1929. Nojidō.

1 ♂ ad. 27. X. 1929. Banpo.

1 ♀ ad. 2. III. 1930. Moppo.

The specimens above mentioned show the special feature of *ussuriensis* with the following measurement:

Wing ♂ ♂ 68, 68.5, ♀ 65. Tail ♂ ♂ 71, 73, ♀ 69.

19. *Pyrrhula pyrrhula kamtschatica* Taczanowski. ベニバラウソ

1 ♂ ad. 12. XI. 1929. Banpo.

(See the "Tori" Vol. VI. p. 257.)

20. *Pyrrhula pyrrhula griseiventris* Lafresnaye. ウソ

3 ♂ ♂ ads., 1 ♀ ad. 28. X.-9. XI. 1929. Banpo.

1 ♂ ad., 4 ♀ ♀ ads. 24. II.-2. III. 1930. Moppo.

21. *Pyrrhula cassini* Baird. ハイイロウソ

♀ ad. 6. XI. 1929. Banpo.

(See "Tori" Vol. VI. p. 257.)

22. *Erythrina rosea* (Pallas). オホマシロ

2 ♂ ♂ imms., 1 ♀ ad. 27. X. 1929. Banpo.

23. *Loxia curvirostra japonica* Ridgway. イスカ

1 ♂ ad., 3 ♀ ♀ ads. 24. II. 1930. Moppo.

It has long been believed that there live both *L. c. japonica* and *L. c. caucasica* in Japan. But it is wrong, because the difference of size which has hitherto been regarded as the main point of distinction between them is but a variation within one and the same subspecies. This variation is wing ♂ ♂ 92-102, ♀ ♀ 87-97. Thus, there is no difference between this variation and the variation of European *L. c. curvirostra*, wing ♂ ♂ 95-100, ♀ ♀ 91-98. It is, however, an unquestionable fact that those in Japan have a deeper and brighter color, so far as the plumage of adult male are concerned, than those in Europe, and only from this point, they are distinguishable from *L. c. curvirostra*. *L. c. caucasica* is a synonym of *L. c. japonica*, I believe.

24. *Fringilla montifringilla* Linnaeus. ア ト リ
 6 ♂♂ ads., 4 ♀♀ ads. 4.-17. IV. 1929. Riuganpo.
 1 ♂ ad. 29. I. 1930. Moppo.
 1 ♂ ad. 30. III. 1930. Quelpart I.
25. *Leucosticte arctoa brunneinucha* (Brandt). ハギマシコ
 2 ♂♂ ads., 2 ♀♀ ads. 15. XI. 1929. Banpo.
26. *Passer montanus saturatus* Stejneger. スズメ
 1 ♂ ad., 2 ♀♀ ads. 19.-21. III. 1930. Quelpart I.
27. *Passer montanus dybowskii* Domaniewski. テウセンズズメ
 2 ♂♂ ads., 2 ♀♀ ads. 4. IV.-4. VI. 1929. Riuganpo.
 2 ♂♂ ads., 1 ♀ ad. 20.-24. IV. 1929. Chōsen.
 2 ♂♂ ads. 22. VIII. 1929. Nojidō.
 3 ♂♂ ads., 1 ♀ ad. 16.-29. X. 1929. Banpo.
 2 ♂♂ ads., 1 ♀ ad. 20. XII. 1929. Moppo.
28. *Passer rutilaus rutilaus* (Temminck). ニウナイスズメ
 Synonym: *P. r. parvirostris* Momiyama.
 5 ♂♂ ads., 3 ♀♀ ads. 13.-27. VI. 1929. Chōsen.
 1 ♂ ad. 26. III. 1930. Quelpart I.

There is no distinction between those in Japan (Hondo) and those in the Quelpart Is.

29. *Emberiza rutila* Pallas. シマノジロ
 7 ♂♂ ads., 2 ♀♀ ads. 10. V.-4. VI. 1929. Riuganpo.
30. *Emberiza aureola* Pallas. シマアヲジ
 7 ♂♂ ads. 3 ♀♀ ads. 30. IV.-16. V. 1929. Riuganpo.
31. *Emberiza elegans elegans* Temminck. ミヤマホホジロ
 Synonym: *E. e. ticehursti* Sushkin.
 3 ♂♂ ads., 7 ♀♀ ads. 3.-13. IV. 1929. Riuganpo.
 1 ♂ ad. 30. VI. 1929. Chōsen.
 1 ♂ ad., 3 juvs. 27. VII.-21. VIII. 1929. Nojidō.
 7 ♂♂ ads., 1 ♀ ad. 15.-21. X. 1929. Banpo.
 1 ♂ ad. 19. I. 1930. Moppo.
 1 ♂ ad. 10. III. 1930. Quelpart I.

Those which pass the northern part of Korea in spring and autumn

(the so-called *ticehursti*) and those which breed in Korea are both undistinguishable from which breed in Hondo (that is, *E. c. elegans*).

32. *Emberiza spodocephala extremi-orientis* Shulpin. カラアマジ

9 ♂♂ ads., 10 ♀♀ ads. 13. IV.-6. V. 1929. Riuganpo.

1 juv. 2. VIII. 1929. Nojidō.

1 ♂ ad., 6 ♀♀ ads., 2 imms. 19. IX.-18. X. 1929. Banpo.

Twenty-nine specimens above mentioned all show the special feature of *E. s. extremi-orientis*, with a deep yellow in their throats and breasts. Sometimes, *E. s. spodocephala* which have not been found in the collection of this time are collected in Korea. (There are some specimens of them in the collection of Takatsukasa!)

33. *Emberiza cioides weigoldi* Jacobi. ジベリアホホジロ

5 ♂♂ ads., 2 ♀♀ ads. 7.-10. IV. 1929. Riuganpo.

5 ♂♂ ads., 3 ♀♀ ads. 15. IX.-19. XI. 1929. Banpo.

The results of measurement of the above-mentioned 15 specimens are as follows:

10 ♂♂ ads. Wing 74-82. Tail 73-81.

5 ♀♀ ads. Wing 72-75. Tail 69-78.

E. c. weigoldi breed in the most northern part of Korea, and in winter they migrate southward to the middle part (to the neighbourhood of Keijō). Those which breed in the neighbourhood of Chōsen (in the northern part of Kogendō) belong not to this subspecies, but to next one.

34. *Emberiza cioides castaneiceps* Moore. テウセンホホジロ

Synonym: *E. c. tyoosenica* Momiyama.

3 ♂♂ ads. 14.-15. VI. 1929. Chōsen.

1 ♀ ad. 6. I. 1930. Mōppo.

The results of measurement of the above-mentioned specimens are as follows:

Wing ♂♂ 77-80. ♀ 74.

Tail ♂♂ 72-77. ♀ 72.

Mr. Momiyama has given the name, *E. c. tyoosenica* to them, but there is no difference between them and *E. c. castaneiceps* which are

natives of the eastern part of China, neither in measurement nor in coloration.

35. *Emberiza cioides ijunae* Stejneger. イイジマホホジロ

6 ♂♂ ads., 3 ♀♀ ads. 10-30. III. 1930. Quelpart Is.

The result of measurement of the nine specimens above mentioned are as follows:

Wing ♂♂ 73-78, ♀♀ 71-73

Tail ♂♂ 69-75, ♀♀ 68-69

Those which live in the Dagelet I., the Quelpart I., Tsushima I. and the Iki I. belong to this subspecies. Those in Kiusiu also have hitherto been regarded as belonging to it. But that is not the case, but they are typical *E. c. ciopsis*.

36. *Emberiza jankowskii* Taczanowski. コマホホジロ

3 ♂♂ ads., 3 ♀♀ ads. 15. IX.-25. X. 1929. Banpo.

They are the first specimens collected in Korea. (See the "Tori" Vol. VI, p. 258.)

37. *Emberiza fucata fucata* Pallas. ホホアカ

Synonym: *E. f. laubmanni* Stachanow.

7 ♂♂ ads., 1 ♀ ad. 17. IV.-6. V. 1929. Riuganpo.

1 ♂ ad. 13. VI. 1929. Chōsen.

1 ♂ ad. 23. II. 1930. Moppo.

1 ♂ ad. 13. III. 1930. Quelpart I.

As the result of investigating a great number of specimens of the same season, it has become known that there is no distinction between those which pass the northern part of Korea (*E. f. fucata*) and those which breed in Hondo.

38. *Emberiza rustica* Pallas. カシラダカ

Synonym: *E. r. latifuscia* Portenko.

5 ♂♂ ads., 2 ♀♀ ads. 2.-27. X. 1929. Banpo.

2 ♂♂ ads., 1 ♀ ad. 25.-26. XI. 1929. Kinkwa.

1 ♂ ad. 2. I. 1930. Moppo.

39. *Emberiza pusilla* Pallas. コホホアカ

- 3 ♂♂ ads., 1 ♂ ? ad., 3 ♀♀ ads. 28. IV.-6. V. 1929. Riuganpo.
40. *Emberiza chrysophrys* Pallas. キマユホホジロ
2 ♂♂ ads. 2. V. 1929. Riuganpo.
41. *Emberiza tristrami* Swinhoe. シロハラホホジロ
7 ♂♂ ads., 5 ♀♀ ads. 24. IV.-6. V. 1929. Riuganpo.
42. *Emberiza yessoensis continentalis* Witherby.
テウセンコジュリン
1 juv., 1 ♀ ad., 1 unsexed ad. 15.-26. X. 1929. Banpo.
2 ♂♂ ads., 1 ♀ ad. 27. XII. 1929. Moppo.
- E. y. continentalis* is easily distinguishable from the other subspecies.
43. *Emberiza pallasi pallasi* (Cabanis). シベリアジュリン
1 ♂ ad., 6 ♀♀ ads., 3 ♀♀ imms. 26. XI.-1. XII. 1929. Kinkwa.
3 ♂♂ ads., 3 ♀♀ ads., 1 imm. 24. XII. 1929-6. I. 1930. Moppo.
44. *Calcarius lapponicus coloratus* Ridgway. ツメナガホホジロ
1 ♀ ad. 4. IV. 1929. Riuganpo.
3 ♂♂ ads., 1 ♂ imm., 6 ♀♀ ads. 13. XII. 1929-6. I. 1930. Moppo.
- Family ALAUDIDÆ. 雲雀科
45. *Calandrella brachydactyla dukhunensis* (Sykes).
カラフトコヒバリ
1 ♂ ad. 17. IV. 1929. Riuganpo.

This is the first specimen collected in Korea (see the "Tori" Vol. VI. p. 169).

46. *Galerida cristata coreensis* Taczanowski. カンムリヒバリ
8 ♂♂ ads., 3 ♀♀ ads. 7.-27. IV. 1929. Riuganpo.
1 ♂ ad. 7. I. 1930. Moppo.
47. *Alauda arvensis pekinensis* Swinhoe. オホヒバリ
2 ♂♂ ads., 3 ♀♀ ads. 13. XII. 1929. Moppo.
- A few of this subspecies also pass the winter in Korea.
48. *Alauda arvensis lönnerbergi* Hachisuka. カラフトチウヒバリ
3 ♂♂ ads., 4 ♀♀ ads. 8.-20. X. 1929. Banpo.
2 ♂♂ ads., 1 ♀ ad. 13. XII. 1929. Moppo.
1 unsexed ad. 14. III. 1930. Quelpart Is.

Many of this subspecies pass Korea in spring and autumn, and those

which pass the winter there are not so rare.

49. *Alauda arvensis intermedia* Swinhoe. ウチヒバリ

1 ♂ ad., 3 ♀ ♀ ads. 4.-18. IV. 1929. Riuganpo.

11 ♂ ♂ ads., 3 ♀ ♀ ads. 14. VI.-15. VII. 1929. Chōsen.

1 ♂ ad., 2 juvs., 1 unsexed ad. 30. VII.-15. VIII. 1929. Nojidō.

6 ♂ ♂ ads., 5 ♀ ♀ ads., 1 ♀ imm. 14. IX.; 9. XI. 1929. Banpo.

2 ♂ ♂ ads. 2.-23. I. 1930. Moppo.

Those which breed in Korea all belong to this subspecies. In winter most of them migrate away southward and a few remain in Korea. Therefore, in winter three subspecies, that is, *pekinensis*, *lönnerbergi* and *intermedia*, live together in Korea.

50. *Alauda arvensis japonica* T. & S. ヒバリ

Synonym: *A. a. quelpartae* Momiyama.

2 ♂ ♂ ads. 21.-30. III. 1930. Quelpart I.

A. arvensis in the Quelpart I. is undistinguishable from those in Japan.

Family MOTACILLIDÆ. 鵲 鵲 科

51. *Anthus richardi richardi* Vieillot. マミジロタヒバリ

5 ♂ ♂ ads., 1 unsexed ad. 4.-12. V. 1929. Riuganpo.

The result of measurement is as follows:

Wing 90-93.5. Ex. c. 14.5-16.

Claw of hind-toe 16.5-18.

52. *Anthus hodgsoni berezowskii* Sarudny. ビンズイ

8 ♂ ♂ ads., 3 ♀ ♀ ads. 29. IV.-11. V. 1929. Riuganpo.

1 ♀ ad. 9. X. 1929. Banpo.

1 ♂ ad. 17. I. 1930. Moppo.

53. *Anthus gustavi gustavi* Swinhoe. セジロタヒバリ

6 ♂ ♂ ads., 1 ♀ ad. 29. IV.-3. V. 1929. Riuganpo.

5 ♂ ♂ ads. 1.-11. X. 1929. Banpo.

The length of wing of the above-mentioned 12 specimens are as follows:

11 ♂ ♂ ads. 75-87, 1 ♀ ad. 77.

54. *Anthus gustavi menzbieri* Shulpin. コセジロタヒバリ

3 ♂♂ ads., 1 ♀ ? ad. 29. IV.-3. V. 1929. Riuganpo.

These 4 specimens coincide with the description of *Anthus g. menzbieri*. The length of wing of the above mentioned specimens is as follows:

4 ads. 76-82.

55. *Anthus rufogularis* Brehm. ムネアカタヒバリ

8 ♂♂ ads., 3 ♀♀ ads. 28. IV.-4. V. 1929. Riuganpo.

2 ♂♂ ads. 4.-8. X. 1929. Banpo.

1 ♀ ad. 11. III. 1930. Quelpart I.

56. *Anthus spinoletta blakistoni* Swinhoe. サメイロタヒバリ

1 ♂ ad. 1. XII. 1929. Kinkwa.

This is the first specimen collected in Korea. ("Tori" Vol. VI, p. 256.) According to Mr. La Touche, many of this subspecies live in the northern China in winter, so it is no cause for wonder that it was collected in Korea.

57. *Anthus spinoletta japonicus* T. & S. タヒバリ

Synonym: *A. borealis* Hesse.

1 ♂ ad., 5 ♀♀ ads. 15. IV.-3. V. 1929. Riuganpo.

2 ♂♂ ads., 9 ♀♀ ads. 26. IX.-29. X. 1929. Banpo.

1 ♂ ad. 19. II. 1930. Moppo.

I have already described in the "Journal für Ornithologie" 1931, p. 504, that *Anthus borealis* resembles closely *Anthus s. japonicus*. And, as the result of the further investigation of a great number of specimens, it has become known that both are quite undistinguishable from each other.

58. *Motacilla flava macronyx* (Stresemann).

キタツメナガセキレイ

8 ♂♂ ads., 2 ♀♀ ads. 7.-16. IV. 1929. Riuganpo.

1 ♂ ad. 20. IX. 1929. Banpo.

These specimens are typical *macronyx*. There is one more specimen, ♂ ad. 17. IX. 1929, Banpo. It has almost undiscernible grey patches in its supercilium. It is an intermediate specimen between *M. f. simillima* and *M. f. macronyx*, I believe.

59. *Motacilla flava simillima* Hartert. マミジロツメナガセキレイ

2 ♂♂ ads., 1 ♀ ad. 17.-18. IX. 1929. Banpo.

These specimens are typical *simillima*. There is one more specimen, ♂ ad., 8. V. 1929, Riuganpo. Its supercilium is not white, but grey. It is also an intermediate specimen between *macronyx* and *simillima*, I believe.

60. *Motacilla flava taiwanus* (Swinhoe). ツメナガセキレイ

1 ♂ ad., 1 ♀ ad. 2. V. 1929. Riuganpo.

These specimens are typical *taiwanus*. There is one more, ♀ ad., 13. IV. 1929, Riuganpo. Its supercilium is yellow, but the color is very faint. It is an intermediate specimen between *macronyx* and *taiwanus*, I believe.

61. *Motacilla cinerea caspica* (S. G. Gmelin). キセキレイ

2 ♀♀ ads. 9.-30. IV. 1929. Riuganpo.

1 ♂ ad., 2 ♀♀ ads. 16. VI.-12. VII. 1929. Chōsen.

2 ♀♀ ads. 26.-27. IX. 1929. Banpo.

1 ♀ ad. 11. III. 1930. Quelpart I.

62. *Motacilla alba leucopsis* Gould. ホホジロセキレイ

7 ♂♂ ads., 1 ♀ ad., 1 juv. 7.-28. IV. 1929. Riuganpo.

1 ♂ ad. 14. VI. 1929. Chōsen.

One young among the above-mentioned specimens is one which has just left its nest. That it left its nest in the last decade of April is rather an early example for that region.

63. *Motacilla alba ocularis* Swinhoe. タイワンハクセキレイ

3 ♂♂ ads., 5 ♀♀ ads. 28. IV.-2. V. 1929. Riuganpo.

3 ♂♂ ads., 1 ♀ ad., 4 imms. 17.-27. IX. 1929. Banpo.

64. *Motacilla alba lugens* Kittlitz. ハクセキレイ

1 ♂ ad., 1 ♀ ad. 11. III. 1930. Quelpart I.

? *Motacilla alba amurensis* Seeböhm.

1 ♂ ad. (No. 4012) 4. IV. 1929. Riuganpo.

This specimen is a full adult male and has a black back. Nevertheless, its remex is pale black as in *M. a. ocularis* and its chin and throat are also quite black as in *ocularis*. (The remex of a young bird of *M. a. lugens* is not pale black, but deep brownish black.) This fact wholly coincides with the description of *Motacilla amurensis* Seeböhm (1878—Gulf of Abrek, in the Sea of Japan). But it cannot be determined without

further investigation whether it is a good subspecies which lies between *lugens* and *ocularis*, or only a hybrid between them.

65. *Dendronanthus indicus* (Gmelin). イワミセキレイ

4 ♂♂ ads. 9.-17. V. 1929. Riuganpo.

4 ♂♂ ads., 1 ♀ ad., 1 unsexed ad. 13. VI.-4. VII. 1929. Chōsen.

Family ZOSTEROPIDÆ. 繡眼兒科

66. *Zosterops palpebrosa ijimae* Kuroda. イイジマメジロ

5 ♂♂ ads., 5 ♀♀ ads. 9.-18. III. 1930. Quelpart I.

67. *Zosterops erythropleura erythropleura* Swinhoe.

テウセンメジロ

1 ♂ ad., 1 ♀ ad. 16. V. 1929. Riuganpo.

Family CERTHIIDÆ. 木 走 科

68. *Certhia familiaris familiaris* Linnaeus. キタキバシリ

Synonym: *C. f. ernsti* Kuroda.

C. f. kawamurai Momiyama.

6 ♂♂ ads., 4 ♀♀ ads. 22.-26. X. 1929. Banpo.

Family SITTIDÆ. 五 十 雀 科

69. *Sitta europaea amurensis* Swinhoe. アムールゴジフカラ

1 ♂ ad., 1 ♀ ad. 11.-19. X. 1929. Banpo.

It has hitherto been believed that *S. europaea* which live in Korea all belong to *amurensis*. But, according to the recent study, those which have been collected in Korea are not *amurensis*, but *hondoensis*, and the two specimens collected this time are the first true *amurensis* collected there. *S. e. amurensis* do not breed in Korea but merely migrate there rarely in winter.

70. *Sitta europaea hondoensis* Buturlin. コジフカラ

7 ♂♂ ads., 3 ♀♀ ads. 20. VI.-12. VII. 1929. Chōsen.

9 ♂♂ ads., 1 ♀ ad., 2 juvs. 25. VII.-1. IX. 1929. Nojidō.

3 ♂♂ ads. 2.-11. X. 1929. Banpo.

Those which breed in Korea are undistinguishable from those in Hondo neither in measurement nor in coloring.

71. *Sitta canadensis corea* O.-Grant. テウセンゴジフカラ

4 ♂♂ ads., 1 ♀ ad. 30. VII.-29. VIII. 1929. Nojidō.

Family PARIDÆ 四十雀科

72. *Parus major takahashii* Momiyama. テウセンシジフカラ

2 ♂♂ ads., 2 ♀♀ ads. 4.-29. IV. 1929. Riuganpo.

4 ♂♂ ads., 3 ♀♀ ads., 3 juvs. 15. VI.-9. VII. 1929. Chōsen.

1 ♂ ad., 1 ♀ ad., 1 juv. 10. VIII.-1. IX. 1929. Nojidō.

1 ♂ subad., 2 ♀♀ subads., 1 unsexed ad. 30. IX.-19. X. 1929. Banpo.

P. major in Korea are easily distinguishable from those in Hondo.73. *Parus major minor* T. & S. シジフカラSynonym: *P. m. quelpartensis* Kuroda.

5 ♂♂ ads., 4 ♀♀ ads. 10.-30. III. 1930. Quelpart I.

Parus major of Quelpart I. are quite undistinguishable from *P. m. minor* of Hondo in a great number of specimens.74. *Parus varius varius* T. & S. ヤマガラSynonym: *P. v. saisiuensis* Kuroda & Mori.*P. v. koreensis* (Kuroda & Mori).

1 ♂ ad., 2 ♀♀ ads., 3 juvs., 1 unsexed ad.

13. VI.-12. VII. 1929. Chōsen.

2 ♂♂ ads., 2 ♀♀ ads. 25. II.-3. III. 1930. Moppo.

5 ♂♂ ads., 3 ♀♀ ads. 16.-30. III. 1930. Quelpart I.

75. *Parus ater amurensis* Buturlin. キタヒガラSynonym: *P. a. tyoosenensis* Momiyama.

2 ♂♂ juvs., 1 ♀ ad. 9 VII. 1929. Chōsen.

1 ♀ ad., 5 juvs. 26. VII.-29. VIII. 1929. Nojidō.

2 unsexed ad. 19. X. 1929. Banpo.

1 ♀ ad. 27. XI. 1929. Kinkwa.

1 ♂ ad. 11. I. 1930. Moppo.

All of *P. ater* in Korea and Saghalin belong to this subspecies76. *Parus ater insularis* Hellmayr. ヒガラSynonym: *P. a. takahashii* Momiyama.

1 ♂ ad. 16. III. 1930. Quelpart I.

As the result of investigating many specimens, it has become known that *P. ater* in the Quelpart I. are undistinguishable from those in Hondo.77. *Parus palustris hellmayri* Bianchi. テウセンコガラ

- 1 ♂ ad., 2 ♀ ♀ ads. 16.-30. VI. 1929. Chōsen.
 1 ♂ ad., 2 imms. 27. XI.-4. XII. 1929. Kinkwa.
 1 ♀ ad. 1. III. 1930. Moppo.

P. p. hellmayri distribute all over Korea, except the northernmost part. And the results of measurement are follows:

Wing ♂ ♂ 58-62, ♀ ♀ 58-62.
 Tail ♂ ♂ 51-55, ♀ ♀ 50-54.

78. *Parus palustris jeholicus* Kleinschmidt. カンキョウコガラ
 9 ♂ ♂ ads., 4 ♀ ♀ ads., 5 unsexed ad., 1 imm.

24. VII.-1. IX. 1929. Nojidō.

1 unsexed ad. 6. IX. 1929. Sanchō.

The results of measurement of the above-mentioned 9 ♂ ♂ ads., 4 ♀ ♀ ads., 6 unsexed ad. are as follows:

Wing: ♂ ♂ 62-68, ♀ ♀ 64-65, unsexed 62-69.
 Tail: ♂ ♂ 59-65, ♀ ♀ 60-63, unsexed 57-68.

P. p. jeholicus is the first collected in Korea. In N. E. Chihli, *P. p. jeholicus* distributes between the distribution of *P. p. crassirostris* and *P. p. hellmayri*. In Korea also it distributes in Kankyodo, North Korea, as an intermediate form between *crassirostris* in Ussuri and *hellmayri* in Korea.

79. *Aegithalos caudatus caudatus* (Linnaeus). カウライシマエナガ

Synonym: *A. c. magna* Clark.

1 ♂ ad., 2 ♀ ♀ ads., 1 unsexed ad., 1 ♂ juv.

26. VII.-3. IX. 1930. Nojidō.

1 ♂ ad., 1 unsexed ad. 27. I. 1930. Moppo.

The results of measurement of the above-mentioned 3 ♂ ♂, 2 ♀ ♀, 1 unsexed ad. are as follows:

Wing ♂ ♂ 62-64, ♀ ♀ 61, 61, unsexed 64, 64.
 Tail ♂ ♂ 88-105, ♀ ♀ 98, 98, unsexed 91-93.

Of the collection this time, the two specimens above mentioned were collected in South Korea. But *A. c. caudatus* migrate to South Korea only in winter, I believe.

80. *Aegithalos caudatus shimokoriyamae* Kuroda.

テウセンエナガ

9 ♂♂ ads., 1 ♀ ad., 1 subad., 1 unsexed ad.

11. XII. 1929-29. I. 1930. Moppo.

The results of measurement of 9 ♂♂ ads., 1 ♀ ad. above-mentioned are as follows:

Wing ♂♂ 61-63 (one=59) ♀ 61.

Tail ♂♂ 81-88 (one=73) ♀ 77.

81. *Aegithalos caudatus trivirgatus* (T. & S.) エナガ

Synonym: *A. c. tarihoe* Momiyama.

1 ♀ ad. 17. III. 1930. Quelpart I.

The measurement of this specimen is as follows:

Wing 60.5. Tail 73.

Mr. Momiyama has said that *A. caudatus* in the Quelpart I. have the following measurement:

Wing 56-59.5, tail 71.5-80.

and have a larger size than those of Hondo. But also in Hondo, there are not a few which have tails above 80 mm. long, and sometimes even a specimen which has a tail 84 mm. long is found there. Even in coloring, there is no distinction between those in Hondo and those in the Quelpart Island.

82. *Remiz pendulinus consobrinus* (Swinhoe) スキンホーガラ

6 ♂♂ ads., 2 ♀♀ ads. 25. IV.-2. V. 1929. Riuganpo.

83. *Regulus regulus japonensis* Blakiston. キクイタダキ

4 ♂♂ ads., 6 ♀♀ ads., 1 unsexed ad. 16.-21. X. 1929. Banpo.

1 ♀ ad. 9. I. 1930. Moppo.

1 ♀ ad. 18. III. 1930. Quelpart I.

84. *Suthora webbiana fulvicauda* Campell. ダルマエナガ

4 ♂♂ ads., 5 ♀♀ ads., 1 unsexed ad.

11. XII. 1929-9. I. 1930. Moppo.

Family LANIIDÆ. 鷓鴣科

85. *Lanius sphenocercus sphenocercus* Cabanis. オホカラモズ

1 ♀ ad. 20. VIII. 1929. Nojidō.

1 ♂ ad., 1 ♀ ad. 6.-7. IX. 1929. Sanchō.

1 ♂ ad. 18. IX. 1929. Banpo.

1 ♂ ad., 1 ♀ ad. 25. XI.-1. XII. 1929. Kinkwa.

86. *Lanius tigrinus* Drapiez. チゴモズ

5 ♂ ads., 2 ♀ ads., 2 unsexed ad. 17.-26. V. 1929. Riuganpo.

1 ♂ ad. 16. VI. 1929. Chōsen.

87. *Lanius cristatus confusus* Stegmann. ウスアカモズ

1 ♂ ad., 1 ♀ ad. 11.-21. V. 1929. Riuganpo.

These two specimens coincide with the description of *L. c. confusus*, with an intermediate coloring between *L. c. cristatus* and *L. c. lucionensis* (see the "Tori" Vol. VII. p. 2).

88. *Lanius cristatus lucionensis* Linnaeus. シマアカモズ

6 ♂ ads., 3 ♀ ads. 9.-24. V. 1929. Riuganpo.

1 ♂ ad. 26. VII. 1929. Nojidō.

89. *Lanius bucephalus* T. & S. モズ

1 ♂ ad., 1 ♀ vix ad. 5. IV. 1929. Riuganpo.

1 ♀ ad. 15. VI. 1929. Chōsen.

1 ♂ juv. 22. X. 1929. Banpo.

1 ♂ ad., 5 ♀ ads. 11. XII. 1929-24. II. 1930. Moppo.

1 ♂ ad., 1 ♂ subad. 9. III. 1930. Quelpart I.

Family BRACHYPODIDÆ. 鶇科

90. *Microscelies amaurotis hensoni* (Stejneger). エゾヒヨドリ

6 ♂ ads., 4 ♀ ads. 18. XII. 1929-26. II. 1930. Moppo

1 ♂ ad. 30. III. 1930. Quelpart I.

The above-mentioned specimens are undistinguishable from those collected in Hokkaido.

91. *Microscelies amaurotis amaurotis* (Temminck). ヒヨドリ

Synonym: *M. a. kanrasani* Momiyama.

6 ♂ ads., 1 ♀ ad. 17-30. III. 1930. Quelpart I.

These specimens are undistinguishable from the natives of Hondo.

Family CAMPEPHAGIDÆ. 山椒喰科

92. *Pericrocotus roseus intermedius* Clark. テウセンサンセウクヒ

5 ♂ ads., 5 ♀ ads., 1 unsexed ad.

24. IV.-16. V. 1929. Riuganpo.

Family MUSCICAPIDÆ. 鶇 科

93. *Tchitrea paradisea incei* (Gould). カワリサンコウテウ
3 ♂♂ ads., 3 ♀♀ ads. 25. V.-4. VI. 1929. Riuganpo.
94. *Alseonax latirostris latirostris* (Raffles). コサメビタキ
6 ♂♂ ads., 2 ♀♀ ads., 2 unsexed ad. 4.-12. V. 1929. Riuganpo.
95. *Hemichelidon griseisticta griseisticta* Swinhoe. エゾビタキ
2 ♂♂ ads., 2 ♀♀ ads. 20.-23. V. 1929. Riuganpo.
96. *Hemichelidon sibirica incerta* La Touche. ウスサメビタキ
Synonym: *H. s. opaca* Shulpin.

3 ♂♂ ads., 6 ♀♀ ads. 24.-28. V. 1929. Riuganpo.

The specimens collected in Korea, Saghalin and the Kuril Islands have feathers of a pale color than those in Hondo. This fact coincides with the description of *incerta* La Touche and with description of *opaca* Shulpin. It is, however, unable to determine whether those in Hondo belong to *H. s. sibirica* or not, as there has never been found a specimen from the type locality of *sibirica*.

97. *Siphia parva albicilla* (Pallas). フジロヒタキ
1 ♂ ad., 1 ♀ ad. 8.-19. A. 1929. Riuganpo.
98. *Siphia mugimaki* (Temminck). ムギマキ
2 ♂♂ ads., 5 ♀♀ ads., 3 imms. 7.-23. V. 1929. Riuganpo.
99. *Zanthopygia nacrissina zanthopygia* (Hay). マミジロキビタキ
8 ♂♂ ads., 4 ♀♀ ads. 28. IV.-12. V. 1929. Riuganpo.
100. *Cyanoptila cyanomelana cyanomelana* (Temminck). オホルリ
6 ♂♂ ads., 4 ♀♀ ads. 24. IV.-7. V. 1929. Riuganpo.
1 ♂ ad. 29. VII. 1929. Nojidō.

Those which breed in Korea all belong to *C. c. cyanomelani* and are undistinguishable from those in Hondo.

101. *Cyanoptila cyanomelana intermedia* (Weigold).

テウセンオホルリ

1 ♂ ad. 11. V. 1929. Riuganpo.

This specimen has a conspicuously pale color compared with those in Korea and Hondo. It corresponds to *C. c. intermedia*. I have already described that *C. c. intermedia* do not breed in Korea.

Family SYLVIIDÆ. 鶯 科

- 102.
- Acanthopneuste nitidus plumbeitarsus*
- (Swinhoe).

ヤナギムシクヒ

3 ♂♂ ads., 2 ♀♀ ads. 17.-27. V. 1929. Riuganpo.

- 103.
- Acanthopneuste tenellipes*
- (Swinhoe). エゾムシクヒ

3 ♂♂ ads., 4 ♀♀ ads., 1 unsexed ad. 14. V.-3. VI. 1929. Riuganpo.

- 104.
- Acanthopneuste borealis borealis*
- (Blasius). コムシクヒ

10 ♂♂ ads., 7 ♀♀ ads. 17. V. 3. VI. 1929. Riuganpo.

1 ♂ ad. 16. IX. 1929. Banpo.

The results of measurement of the 18 specimens above-mentioned are as follows:

Wing ♂♂ 63-69.5, ♀♀ 62.5-65.

Tail ♂♂ 44-49, ♀♀ 43-48.

- 105.
- Acanthopneuste borealis xanthodryas*
- (Swinhoe). メボン

3 ♀♀ ads. 28. V.-4. VI. 1929. Riuganpo.

These specimens are determined to belong to *A. b. xanthodryas*, owing to the fact that they are rich in yellow in the under surface and have a long 1st primary. Those which breed in Korea belong to this subspecies, I believe.

- 106.
- Acanthopneuste occipitalis coronata*
- (T. & S.).

センダイムシクヒ

7 ♂♂ ads., 2 ♀♀ ads. 16.-28. IV. 1929. Riuganpo.

- 107.
- Phylloscopus inornatus inornatus*
- (Blyth). キマユムシクヒ

7 ♂♂ ads., 7 ♀♀ ads. 16. IV.-18. V. 1929. Riuganpo.

- 108.
- Phylloscopus proregulus proregulus*
- (Pallas).

カラフトムシクヒ

5 ♂♂ ads., 7 ♀♀ ads. 5.-17. IV. 1929. Riuganpo.

- 109.
- Phylloscopus fuscatus fuscatus*
- (Blyth). ムヂセツカ

1 ♂ ad. 26. IV. 1929. Riuganpo.

1 ♂ ad. 1. X. 1929. Banpo.

- 110.
- Herbivocula schwarzi*
- (Radde). カラフトムヂセツカ

3 ♂♂ ads., 1 ♀? ad. 11.-20. V. 1929. Riuganpo.

1 ♀ ad. 27. VII. 1929. Nojidō.

According to the collector, they sing "hō-hokekyō," which resembling those of *Horeites cantans*.

111. *Horeites cantans borealis* (Campbell). テウセンウグヒス

9 ♂♂ ads. 13. VI.-16. VII. 1929. Chōsen.

1 ♀ juv. 28. VIII. 1929. Nojidō.

1 ♂ ad. 19. X. 1929. Banpo.

Besides, there is one nest (having 3 eggs) which was collected in Nojidō on July 25th, 1929. The collector asserts that it is the nest of one of this subspecies. But the specimen differs from the others in the point that the eggs have large brown blotches on the white ground and the nest is made of many grass stems. So I can not believe the nest belong to *H. cantans*. I shall wait for the result of further investigation.

112. *Horeites cantans takahashii* (Momiyama). サイシウウグヒス

6 ♂♂ ads., 4 ♀♀ ads. 10.-21. III. 1930. Quelpart I.

H. cantans in Hondo and those in Quelpart I. are easily distinguishable from each other by the fact that the latter are richer in olive than the former.

113. *Urosphena squameiceps ussuriensis* (Seeböhm).

テウセンヤブサメ

2 ♂♂ ads., 7 ♀♀ ads. 24. IV.-5. V. 1929. Riuganpo.

1 ♀ ad. 9. VII. 1929. Chōsen.

U. squameiceps which live in Korea have a conspicuously pale feather color compared with those in Hondo in the specimens of the same season. According to Mr. Seeböhm, the type specimen of *U. squameiceps* has dark feather like those in Japan. Therefore, by the fact that the type locality of *U. squameiceps* is Canton, I regard with some doubt those which live in Korea as belonging to *U. s. ussuriensis*.

114. *Locustella fasciolata* (Gray). エゾセンノウ

♂ ad. 4. VI. 1929. Riuganpo.

115. *Locustella certhiola minor* David & Oustalet.

シベリアセンノウ

2 ♂♂ ads., 3 ♀♀ ads. 24.-30. V. 1949. Riuganpo.

The results of measurement of the above-mentioned 5 specimens are as follows:

Wing ♂♂ 62-67, ♀♀ 60-65.

Tail ♂♂ 50-53, ♀♀ 48-52.

116. *Locustella neava lanceolata* (Temminck). マキノセンニウ

7 ♂♂ ads., 3 ♀♀ ads. 11.-23. V. 1929. Riuganpo.

117. *Phragamaticola aëdon rufescens* Stegmann.

ハシプトオホヨシキリ

3 ♂♂ ads., 4 ♀♀ ads. 20.-27. V. 1929. Riuganpo.

118. *Acrocephalus stentoreus orientalis* (T. & S.). オホヨシキリ

9 ♂♂ ads., 2 ♀♀ ads., 1 unsexed ad.

20. V.-3. VI. 1929. Riuganpo.

119. *Acrocephalus bistrigiceps* Swinhoe. コヨシキリ

3 ♂♂ ads., 2 ♀♀ ads. 25. V.-1. VI. 1929. Riuganpo.

1 ♂ ad., 2 ♀♀ ads. 1.-13. X. 1929. Banpo.

Family TURDIDÆ. 鶇科

120. *Turdus aureus aureus* Holandre. トラツグミ

3 ♀♀ ads. 26. IV.-6. V. 1929. Riuganpo.

121. *Turdus sibiricus sibiricus* Pallas. シベリヤマミジロ

7 ♂♂ ads., 3 ♀♀ ads. 16.-20. V. 1929. Riuganpo.

122. *Turdus hortulorum* Selater. カラアカハラ

8 ♂♂ ads., 2 ♀♀ ads. 3. IV.-4. V. 1929. Riuganpo.

123. *Turdus pallidus* Gmelin. シロハラ

1 ♂ ad., 2 ♀♀ ads. 27.-30. IV. 1929. Riuganpo.

1 ♀ ad. 17. III. 1929. Quelpart I.

124. *Turdus obscurus obscurus* Gmelin. マミチヤジナイ

6 ♂♂ ads., 3 ♀♀ ads. 11.-19. V. 1929. Riuganpo.

125. *Turdus naumanni* Temminck. ハチゼウツグミ

2 ♂♂ ads., 5 ♀♀ ads. 4.-28. IV. 1929. Riuganpo.

1 ♂ ad., 1 ♀ ad. 19.-25. I. 1930. Moppo.

126. *Turdus eunomus* Temminck. ツグミ

7 ♂♂ ads., 4 ♀♀ ads. 4. IV.-12. V. 1929. Riuganpo.

1 ♀ ad. 12. III. 1930. Quelpart I.

127. *Monticola gularis* (Swinhoe). ヒメインヒヨ
 8 ♂♂ ads., 3 ♀♀ ads. 11. 19. V. 1929. Riuganpo.
 2 juvs. 26. VIII. 1929. Nojidō.
128. *Monticola philippensis magna* (La Touche). インヒヨドリ
 4 ♂♂ ads., 1 ♂ subad., 5 ♀♀ ads. 14.-24. VI. 1929. Chōsen.
 1 ♂ ad. 7. I. 1930. Moppo.
 1 ♂ ad. 13. III. 1930. Quelpart I.
129. *Saxicola torquata stejnegeri* (Parrot) ノビタキ
 3 ♂♂ ads., 4 ♀♀ ads. 6.-29. IV. 1929. Riuganpo.
 1 ♂ ad. 19. VI. 1929. Chōsen.
130. *Ianthia cynura cynura* (Pallas). ルリビタキ
 2 ♂♂ ads., 2 ♂♂ subad., 5 ♀♀ ads. 2.-9. IV. 1929. Riuganpo.
 1 ♂ imm. 19. I. 1930. Moppo.
131. *Phoenicurus aureus aureus* (Pallas). ジャウビタキ
 2 ♀♀ ads. 19.-20. VI. 1929. Chōsen.
 1 ♂ ad., 1 ♀ ad., 1 juv. 28.-29. VII. 1929. Nojidō.
 1 ♀ ad. 24. X. 1929. Banpo.
 1 ♂ ad. 30. XI. 1929. Kinkwa.
 1 ♂ ad. 7. I. 1930. Moppo.

P. a. aureus breed in the mountain region of the middle and the northern part of Korea.

132. *Luscinia calliope calliope* (Pallas). ノゴマ
 1 ♂ ad., 1 ♀ ad. 15.-26. IV. 1929. Riuganpo.

The results of measurement of these specimens are as follows:

Wing ♂ 75, ♀ 73 Tail ♂ 59, ♀ 59

133. *Luscinia calliope camtschatkensis* Gmellin. オホノゴマ
 2 ♂♂ ads. 12.-17. V. 1929. Riuganpo.

The results of measurement of these specimens are as follows:

Wing 80, 80. Tail 54, 60

According to the recent investigation, *L. c. calliope* has wings 71-77.5 mm. long and *L. c. camtschatkensis* wings 77-85 mm. long. It is only in the migration seasons of spring and autumn that specimens which have wings above 78 mm. long are collected in Sagalin, Korea and Manchuria.

134. *Larvivora sibilans* Swinhoe. シマゴマ
7 ♂♂ ads., 2 ♀♀ ads., 1 unsexed ad. 7.-17. V. 1929. Riuganpo.

135. *Larvivora cyane* (Pallas). コ ル リ
7 ♂♂ ads., 3 ♀♀ ads. 10.-18. V. 1929. Riuganpo.

Family PRUNELLIDÆ. 岩 鷓 科

136. *Prunella montanella* (Pallas). ヤマヒバリ
1 ♂ ad. 27. X. 1929. Banpo.

Family TROGLODYTIDÆ. 鶇 鶇 科

137. *Troglodytes troglodytes peninsulæ* (Clark).

テウセンミンソサザイ

Synonym: *T. t. quelpartis* Kuroda & Mori.

T. t. morii Momiyama.

- 1 ♂ ad., 1 unsexed ad. 21.-22. X. 1929. Banpo.

- 3 ♂♂ ads., 2 ♀♀ ads., 1 unsexed ad.

26. XI.-2. XII. 1929. Kinkwa.

- 2 ♂♂ ads., 1 ♀ ad. 13.-27. XII. 1929. Moppo.

- 3 ♂♂ ads., 5 ♀♀ ads. 16.-30. III. 1930. Quelpart I.

T. troglodytes in the Quelpart I. and the Dagelet I. are both undistinguishable from those in Korea, in many specimens.

138. *Circulus pallasii pallasii* Temminck. キタカハガラス

- 1 juv. 2. VII. 1929. Chōsen.

- 1 unsexed ad. 2. XII. 1929. Kinkwa.

Family HIRUNDINIDÆ. 燕 科

139. *Hirundo rustica gutturalis* Scopoli. ツ バ メ

- 7 ♂♂ ads., 4 ♀♀ ads. 19. IV.-4. VI. 1929. Riuganpo.

- 1 ♀ juv. 13. VII. 1929. Chōsen.

- 1 ♂ ad. 13. III. 1930. Quelpart I.

1 juv. above mentioned is pretty rich in red on its under surface.

140. *Hirundo rustica tytleri* Jerdon. アカハラツバメ

- 5 ♂♂ ads., 1 ♀ ad. 3. V.-4. VI. 1929. Riuganpo.

- 3 ♂♂ ads., 1 ♀ ad. 17. IX. 1929. Banpo.

It is worth remarking that in spring *H. r. tytleri* remain in Korea till the beginning of June.

141. *Hirundo daurica nipalensis* Hodgson. コシアカツバメ
7 ♂♂ ads., 2 ♀♀ ads. 13. VI.-14. VII. 1929. Chōsen.
1 ♀ ads. 25. VII. 1929. Nojidō.

142. *Delichon urbica dasypus* (Bonaparte). イワツバメ
3 ♂♂ ads., 4 ♀♀ ads. 24. VI.-6. VII. 1929. Chōsen.

143. *Riparia riparia ijimae* (Lönnerberg). セウドウツバメ
2 ♂♂ ads. 29. V. 1929. Riuganpo.
6 ♂♂ ads., 3 ♀♀ ads. 13.-20. IX. 1929. Banpo.

Family MICROPODIDÆ. 雨燕科

144. *Micropus pacificus pacificus* (Latham). アマツバメ
1 ♂ ad. 29. V. 1929. Riuganpo.
4 ♂♂ ads., 3 ♀♀ ads. 3. VII. 1929. Chōsen.
145. *Hirundapus caudacutus caudacutus* (Latham).

ハリヲアマツバメ

- 1 ♀ ad. 16. V. 1929. Riuganpo.
2 ♂♂ ads., 2 ♀♀ ads. 29. VII. 1929. Nojidō.

Family CAPRIMULGIDÆ. 蚊母鳥科

146. *Caprimulgus indicus jotaka* T. & S. ヨタカ
2 ♂♂ ads., 1 ♀ ad. 12.-21. V. 1929. Riuganpo.

Family UPUPIDÆ. 戴勝科

147. *Upupa epops saturata* Lönnerberg. ヤツガシラ
3 ♂♂ ads., 2 ♀♀ ads. 3. IV.-30. V. 1929. Riuganpo.

Family CORACIIDÆ. 佛法僧科

148. *Eurystomus orientalis calonyx* Sharpe. ブツボウソウ
2 ♂♂ ads., 2 ♀♀ ads., 1 unsexed ad. 12.-20. V. 1929. Riuganpo.

Family ALCEDINIDÆ. 翡翠科

149. *Alcedo atthis japonica* Bonaparte. カハセミ
2 ♂♂ ads., 2 ♀♀ ads. 22. IV.-22. V. 1929. Riuganpo.
1 ♂ imm. 15. VII. 1929. Chōsen.

150. *Halcyon coromanda major* (T. & S.). アカセウビン
1 ♂ ad. 9. VII. 1929. Chōsen.

Family PICIDÆ. 啄木鳥科

151. *Picus canus jessoensis* Stejneger. ヤマゲラ

Synonym: *P. c. griseoviridis* (Clark).

1 ♀ ad., 3 juvs. 9.-10. VII. 1929. Chōsen.

5 ♂♂ juvs. 1.-10. VIII. 1929. Nojidō.

2 ♂♂ ads., 1 ♀ ad. 3.-11. XI. 1929. Banpo.

As the result of investigating many specimens, it has become known that there is no difference between *P. canus* in Korea and *P. c. jessoensis* in Hokkaido either in coloring or in measurement.

152. *Dryobates major japonicus* (Seebolm). エゾアカゲラ

3 ♂♂ ads., 2 ♀♀ ads., 2 juvs. 24. VII.-2. IX. 1929. Nojidō.

3 ♂♂ ads. 24. X.-17. XI. 1929. Banpo.

The results of measurement of 6 ♂♂ ads. and 2 ♀♀ ads. above-mentioned are as follows:

Wing 130-135. Tail 82-87. Culmen 26-28.5.

D. major which breed in the mountains of the northern part of Korea and migrate to the middle part in winter have the same coloring and measurement as those in Hokkaido.

153. *Dryobates major hondoensis* Kuroda. アカゲラ

Synonym: *D. m. seoulensis* Kuroda & Mori.

1 ♂ ad., 1 juv. 6.-9. VII. 1929. Chōsen.

1 ♂ ad., 1 ♀ ad. 25.-28. XI. 1929. Kinkwa.

1 ♂ ad. 28. II. 1930. Moppo.

D. major which live in the middle and the southern part of Korea differ from those which breed in the mountain region of Korea, and have the same coloring and measurement as those in Hondo.

154. *Dryobates leucotos uralensis* (Malherbe).

シベリアオホアカゲラ

6 ♂♂ ads., 2 ♀♀ ads., 4 juvs. 24. VII.-22. VIII. 1929. Nojidō.

1 ♂ ad., 2 ♀♀ ads. 3.-17. XI. 1929. Banpo.

I have already described in the "Tori," Vol. VII, p. 1, that *D. leucotos* which live in the mountains in Kankyohokudo, Korea, are identical with *D. l. uralensis* which live in the Ural region.

155. *Dryobates leucotos coreensis* Clark. テウセンオホアカゲラ

3 ♂♂ ads., 1 ♀ ad. 6.-16. VII. 1929. Chōsen.

D. l. coreensis distributes all over Korea, except Kankyohokudo.

156. *Dryobates leucotos quelpartensis* Kuroda & Mori.

サイシウオホアカゲラ

5 ♂♂ ads., 5 ♀♀ ads. 16.-26. III. 1929. Quelpart I.

157. *Dryobates minor nojidoensis* Yamashina. テウセンコアカゲラ

4 ♂♂ ads., 5 ♀♀ ads., 1 ♂ juv., 1 ♀ subad.

30. VII.-1. IX. 1929. Nojidō.

(See the "Tori," Vol. VI, p. 254.)

158. *Dryobates hyperythrus subrufinus* (Cab. & Heine).

チャベラアカゲラ

2 ♂♂ ads., 1 ♀ ad. 12. V.-3. VI. 1929. Riuganpo.

These are the first specimens collected in Korea. (See the "Tori," Vol. VI, p. 168.)

159. *Dryobates nanus doerriesi* (Hargitt). アムールコゲラ

1 ♂ ad., 1 ♀ ad. 9. VII. 1929. Chōsen.

1 ♀ ad. 26. VII. 1929. Nojidō.

2 ♀♀ ads. 16.-28. X. 1929. Banpo.

1 ♂ ad. 24. II. 1930. Moppo.

160. *Dryobates kizuki acutirostris* Yamashina. ハシナガコゲラ

2 ♂♂ ads., 2 ♀♀ ads., 5 juvs.

13. VI.-4. VII. 1929. Kongōsan, Chōsen.

(See the "Tori," Vol. VII, p. 111.)

161. *Dryobates kizuki nippon* (Kuroda). コゲラ

Synonym: *D. k. siragiensis* Momiyama.

1 ♂ ad., 2 ♀♀ ads. 24.-27. II. 1930. Moppo.

These specimens are quite undistinguishable from *D. k. nippon* of Hondo like the type specimens of *D. k. siragiensis* collected in Kōryo, Keikido near Keijō.

162. *Picoides tridactylus kurodai* Yamashina. テウセンミュビゲラ

2 ♂♂ ads., 3 ♀♀ ads. 19. VIII.-2. IX. 1929. Nojidō.

(See the "Tori," Vol. VI, p. 255.)

163. *Dryocopus martius martius* (Linnaeus). クマゲラ

Synonym: *D. m. morii* Kuroda.

2 ♂♂ ads., 8 ♀♀ ads. 24. VII.-19. VIII. 1929. Nojidō.

The results of measurement of the 10 ♂♀ above mentioned are as follows:

Wing	♂♂ 233-247	♀♀ 233-250.
Tail	♂♂ 155-162	♀♀ 155-174.
Culmen	♂♂ 58-60	♀♀ 53-59.

164. *Jynx torquilla intermedia* Stegmann. シベリアアリスヒ

6 ♂♂ ads., 3 ♀♀ ads. 8. IV.-2. V. 1929. Riuganpo.

The results of measurement of the nine specimens above mentioned are as follows:

Wing	♂♂ 81-85	♀♀ 81-83.
Tail	♂♂ 64-73	♀♀ 68-70.
Culmen	♂♂ 13.5-14	♀♀ 13.5-16.

Family CUCULIDÆ. 杜鵑科

165. *Cuculus optatus optatus* Gould. ツツドリ

5 ♂♂ ads., 5 ♀♀ ads. 8. V.-1. VI. 1929. Riuganpo.

166. *Cuculus poliocephalus poliocephalus* Latham. ホトトギス

1 ♂ ad. 24. V. 1929. Riuganpo.

167. *Cuculus micropterus micropterus* Gould. セグロカクコウ

1 ♂ ad. 24. V. 1929. Riuganpo.

168. *Hierococcyx fugax hyperythrus* (Gould). ジウイチ

2 ♂♂ ads., 3 ♀♀ ads. 21.-25. V. 1929. Riuganpo.

Family STRIGIDÆ. 梟鵂科

169. *Bubo bubo tenuipes* Clark. ワシミミズク

1 ♂ ad., 1 ♀ ad. 12. I.-12. II. 1930. Moppo.

170. *Otus bakkamoena ussuriensis* (Buturlin).

サメイロオホコノハヅク

8 ♂♂ ads., 9 ♀♀ ads. 15.-27. X. 1929. Banpo.

1 unsexed ad. 8. II. 1930. Moppo.

The results of measurement of the specimens above mentioned are as follows:

Wing	♂♂ 162-171	♀♀ 176-189.
Tail	♂♂ 70-84	♀♀ 82-89.

A great number of this subspecies pass Korea in spring and autumn and only few pass the winter there.

171. *Otus scops stictonotus* (Sharpe). テウセンコノハヅク

2 ♂♂ ads., 3 ♀♀ ads. 28. IV.-12. V. 1929. Riuganpo.

1 ♂ ad. 28. VIII. 1929. Nōjido.

1 ♂ ad. 5. X. 1929. Banpo.

The measurement of 4 ♂♂ and 3 ♀♀ above mentioned is as follows:

Wing ♂♂ 137-147 ♀♀ 143-147.

Tail ♂♂ 59-68 ♀♀ 63-67.

Those which breed in North Korea belong no doubt to this subspecies. But, sometimes, *O. s. japonicus* are collected in the middle part of Korea, but they are not found in the collection of this time. It is not clear as yet what subspecies those which breed in the middle and the southern part of Korea belong to.

172. *Asio otus otus* (Linnaeus). トラフヅク

1 ♂ ad. 4. IV. 1929. Riuganpo.

1 ♂ ad., 3 ♀♀ ads. 19.-22. X. 1929. Banpo.

173. *Asio flammeus flammeus* (Pontoppidan). コミミヅク

1 ♀ ad. 14. IV. 1929. Riuganpo.

1 ♂ ad., 1 ♀ ad. 12. I.-12. II. 1930. Moppo.

174. *Ninox scutulata ussuriensis* Buturlin. テウセンアラバヅク

3 ♀♀ ads. 8.-21. V. 1929. Riuganpo.

1 ♂ ad. 16. X. 1929. Banpo.

The measurement of 4 specimens above mentioned is as follows:

Wing ♂ 233 ♀♀ 229-232.

Tail ♂ 128 ♀♀ 117-121.

Ninox scutulata ussuriensis merely pass Korea and do not breed there. Those which breed in Korea belong to *N. s. scutulata*. But the latter were not collected this time.

Family FALCONIDÆ. 鷲鷹科

175. *Falco peregrinus calidus* Latham. ハヤブサ

2 imms. 30. IX.-4. X. 1929. Banpo.

176. *Falco subbuteo subbuteo* Linnaeus. チゴハヤブサ

- 4 ♀ ♀ ads. 11.-28. V. 1929. Riuganpo.
 177. *Falco columbarius insignis* (Clark). コ チ ヨ ウ ゲ ン ボ ウ
 1 ♀ ad. 18. X. 1929. Banpo.
 1 ♂ juv. 22. 1. 1930. Chitō.
 178. *Falco vespertinus amurensis* Radde. ア カ シ チ ヨ ウ ゲ ン ボ ウ
 2 juvs. 6. IX. 1929. Sanchō.
 1 juv. 11. X. 1929. Banpo.
 179. *Falco tinnunculus japonensis* Ticehurst. チ ヨ ウ ゲ ン ボ ウ
 1 ♂ ad. 29. XI. 1929. Kinkwa.
 2 ♀ ♀ ads. 24. XII. 1929. Moppo.

The measurement of the 3 specimens above mentioned is as follows:

Wing ♂ 244 ♀ ♀ 251-261.
 Tail ♂ 174 ♀ ♀ 175-183.

180. *Falco tinnunculus tinnunculus* Linnaeus.

テ ウ セ ン チ ヨ ウ ゲ ン ボ ウ

♀ ad. 21. X. 1929. Banpo.

The measurement of this specimen is as follows:

Wing 266. Tail 189.

This specimen is regarded as belonging to *F. t. tinnunculus* owing to its pale feather color. *F. t. tinnunculus* rarely migrate to Korea in winter.

181. *Aquila chrysaetos japonica* Severtzow. イ ス フ シ

1 juv. 2. XII. 1929. Kinkwa.

182. *Buteo ferox hemilasius* T. & S. オ ホ ノ ス リ

1 ♂ ad., 2 ♀ ♀ ads. 17. I.-17. II. 1929. Moppo.

The measurement of the 3 specimens above mentioned is as follows:

Wing ♂ 429 ♀ ♀ 447, 464.
 Ex. culmen. ♂ 26 ♀ ♀ 26, 27.5.

183. *Buteo buteo burmanicus* Hume. ノ ス リ

2 ♂ ♂ ads., 1 ♀ ad. 26.-28. XI. 1929. Kinkwa.

1 ♂ ad., 2 ♀ ♀ ads., 29. I.-19. II. 1930. Moppo.

1 ♀ ad. 30. III. 1930. Quelpart I.

184. *Circus spilonotus* Kaup. シ ベ リ ア チ ウ ヒ

1 ♂ ad. 21. IV. 1929. Riuganpo.

- 1 ♀ ad. 24. VIII. 1929. Nojidō.
185. *Circus aeruginosus aeruginosus* (Linnaeus). チ ウ ヒ
3 ♀ ♀ ads. 13. IX.-12. X. 1929. Banpo.
186. *Circus melanoleucus* (Forster). マダラチウヒ
1 ♂ ad., 3 juvs. 25. VII.-29. VIII. 1929. Nojidō.
1 ♂ ad. 15. IX. 1929. Banpo.
187. *Circus cyaneus cyaneus* (Linnaeus). ハイイロチウヒ
1 ♂ imm. 7. XI. 1929. Banpo.
188. *Accipiter nisus nisosimilis* (Tickell). ハイタカ
1 ♀ ad. 10. IV. 1929. Riuganpo.
1 ♀ ad. 24. VIII. 1929. Nojidō.
1 ♀ juv. 10. X. 1929. Banpo.
2 ♂ ♂ imms., 1 ♀ ad. 11. XII. 1929-20. II. 1930. Moppo.
189. *Accipiter virgatus gularis* (T. & S.). ツ ミ
1 ♀ ad. 14. V. 1929. Riuganpo.
1 ♀ juv. 6. IX. 1929. Nojidō.
190. *Milvus migrans lineatus* (Gray). ト ビ
2 ♀ ♀ ads. 12. I.-19. II. 1930. Moppo.
191. *Haliaeetus pelagicus pelagicus* (Pallas). オホワシ
1 ♂ juv. 24. XII. 1929. Moppo
1 ♀ juv. 22. III. 1930. Quelpart I.
192. *Pandion haliaetus haliaetus* (Linnaeus). ミ サ ゴ
1 ♀ ad., 1 unsexed ad. 2.-25. X. 1929. Banpo.
Family PLEGADIDÆ. 朱 鷲 科
193. *Platalea minor* T. & S. クロツラヘラサギ
1 ♂ ad. 23. IV. 1929. Riuganpo.
Family ARDEIDÆ. 鷺 科
194. *Ardea cinerea rectirostris* Gould. アヲサギ
1 ♂ ad., 1 ♀ ad. 9.-20. IV. 1929. Riuganpo.
2 juvs. 24.-29. IX. 1929. Banpo.
1 ♂ ad. 23. I. 1930. Chitō.
195. *Ardea purpurea manillensis* Meyen. ムラサキサギ
5 juvs. 24. IX.-IX. 1929. Banpo.

196. *Egretta alba alba* (Linnaeus). ダイサギ

1 ad. 31. I. 1930. Moppo.

Measurement: Wing 410, culmen 113.

197. *Egretta alba modesta* (Gray). チウダイサギ [コモ、ジロ]

1 ♀ ad. 22. IV. 1929. Riuganpo.

Measurement: Wing 350, culmen 105.

198. *Hemigarzetta eulophotes* (Swinhoe). カラシラサギ

1 ♂ ad. 30. IV. 1929. Riuganpo.

199. *Demigretta sacra ringeri* Stejneger. クロサギ

1 ♂ ad. 22. III. 1930. Quelpart I.

Measurement: Wing 303, culmen 89.

200. *Ixobrychus eurhythmus* (Swinhoe). オホヨシゴイ

3 ♂ ♂ ads., 3 ♀ ♀ ads. 20.-24. V. 1929. Riuganpo.

Family ANATIDÆ. 雁 鴨 科

201. *Cygnopsis cygnoid* (Linnaeus). サカツラガン

1 ♂ ad., 1 ♀ ad. 7.-10. IV. 1929. Riuganpo.

1 ♂ ad., 1 ♀ ad. 29. IX. 1929. Banpo.

202. *Anser albifrons albifrons* (Scopoli). マ ガ ン

2 ♂ ♂ ads., 1 ♀ ad. 4.-15. IV. 1929. Riuganpo.

1 ♂ ad., 1 ♀ ad., 2 imms. 13. XII. 1929-10. II. 1930. Moppo.

203. *Anser fabalis serratirostris* Swinhoe. ヒシクヒ

1 ♂ ad. 10. X. 1929. Banpo.

1 ♂ ad. 12. II. 1930. Moppo.

The measurement of the 2 specimens above mentioned is as follows:

Wing 455-468, culmen 63-71.

They belong, no doubt, to *serratirostris*, but the following specimens certainly an intermediate form between *serratirostris* and *curtus*.

1 imm. 12. X. 1929. Banpo.

Measurement: Wing 420, culmen 62.

204. *Anser fabalis curtus* Lönnerberg. ヒメヒシクヒ

1 ♂ ad., 1 ♂ juv., 2 ♀ ♀ imms. 20. IX.-12. X. 1923. Banpo.

The measurement of the 4 specimens above mentioned is as follows:

	Wing	Culmen
1 ♂ ad.	421	59
1 ♂ juv.	418	55
2 ♀ ♀ imm.	413-428	54-59
205. <i>Branta bernicla nigricans</i> (Lawrence). コグガン		
1 ♂ ad., 1 ♀ ad.	11. I. 1929.	Moppo.
206. <i>Tadorna tadorna</i> (Linnaeus). ツクシガモ		
2 ♂ ♂ ads.	17. IV. 1929.	Riuganpo.
207. <i>Anas platyrhynchos</i> Linnaeus. マガモ		
1 ♂ ad.	4. IV. 1929.	Riuganpo.
1 ♂ ad., 1 ♀ ad.	8.-12. X. 1929.	Banpo.
208. <i>Anas superciliosa zonorhynchos</i> Swinhoe. カルガモ		
1 ♀ ad.	6. IV. 1929.	Riuganpo.
1 ♀ ad.	13. VII. 1929.	Chōsen.
1 ♂ ad.	10. X. 1929.	Banpo.
1 ♂ ad.	27. XII. 1929.	Moppo.
2 juv. (downy).	25. VI. 1929.	Chōsen.
209. <i>Querquedula crecca crecca</i> (Linnaeus). コガモ		
1 ♂ ad., 2 ♀ ♀ ads.	11.-21. IV. 1929.	Riuganpo.
1 ♂ ad., 1 ♂ ad.	16. IX. 1929.	Banpo.
210. <i>Querquedula querquedula</i> (Linnaeus). シマアザ		
1 ♂ ad., 1 ♀ ad.	17.-20. IV. 1929.	Riuganpo.
211. <i>Mareca penelope</i> (Linnaeus). ヒドリガモ		
1 ♂ ad.	22. IV. 1929.	Riuganpo.
212. <i>Eunetta falcata</i> (Georgi). ヨシガモ		
1 ♂ ad.	22. IV. 1929.	Riuganpo.
1 ♂ imm., 1 ♀ imm.	24. IX. 1929.	Banpo.
213. <i>Spatula clypeata</i> (Linnaeus). ハシビロガモ		
♀ ad.	23. IV. 1929.	Riuganpo.
1 ♂ imm., 1 ♀ ad.	12.-23. X. 1929.	Banpo.
214. <i>Aix galericulata</i> (Linnaeus). ラシドリ		
3 ♂ ♂ ads., 1 ♀ ad.	5.-14. IV. 1929.	Riuganpo.
215. <i>Nyroca fuligula</i> (Linnaeus). キンクロハジロ		

- 1 ♂ imm., 1 ♀ ad., 2 ♀ ♀ imms. 14. X.-12. XI. 1929. Banpo.
 216. *Nyroca marila mariloides* (Vigors). スズガモ
 1 ♂ imm., 1 ♀ ad. 12. XI. 1929. Banpo.
 217. *Bucephala clangula clangula* (Linnaeus). ホホジロガモ
 1 ♀ ad. 28. III. 1930. Quelpart I.
 218. *Melanitta fusca stejnegeri* (Ridgway). ビロウドキンクロ
 1 ♂ ad. 24. X. 1929. Banpo.
 219. *Melanitta nigra americana* (Swainson). クロガモ
 2 ♂ ♂ ads. 24. X. 1929. Banpo.
 220. *Histrionicus histrionicus pacificus* Brooks. シノリガモ
 4 ♂ ♂ ads., 1 ♀ ad. 24. X. 1929. Banpo.
 221. *Mergus merganser merganser* Linnaeus. カハアイサ
 1 ♀ ad. 28. XI. 1929. Kinkwa.
 1 ♀ ad. 18. XII. 1929. Moppo.

The measurement of 2 ♀ ♀ ads. above mentioned is as follows:

Wing 260-267. Culmen 52-56.

222. *Mergellus albellus* (Linnaeus). ミコアイサ
 1 ♂ imm., 4 ♀ ♀ ads. 9.-15. IV. 1929. Riuganpo.
 Family PHALACROCORACIDÆ. 鷗 鷺 科
 223. *Phalacrocorax capillatus* (T. & S.). ウミウ
 2 ♀ ♀ imms. 2.-20. X. 1929. Banpo.
 1 ♂ ad., 1 ♀ ad., 1 unsexed ad. 13.-22. III. 1930. Quelpart I.
 224. *Phalacrocorax pelagicus pelagicus* Pallas. ヒメウ
 1 ♂ imm., 1 ♀ ad., 1 ♀ imm. 13.-22. III. 1930. Quelpart I.
 Family PODICIPIDÆ. 鷺 鷗 科
 225. *Podiceps auritus* (Linnaeus). ミミカイツブリ
 ♀ ad. 6. X. 1929. Banpo.

Family COLUMBIDÆ. 鳩 鴿 科

226. *Columba rupestris rupestris* Pallas. カウライバト
 2 ♂ ♂ ads., 3 ♀ ♀ ads. 15.-18. IV. 1929. Riuganpo.

1 ♀ ad. collected on April 18th has an egg in its oviduct, whose egg-shell has been already formed. The measurement of this egg is 35×26 mm.

227. *Streptopelia orientalis orientalis* (Latham). キジバト
 3 ♂♂ ads., 12-16. IV. 1929. Riuganpo.
 1 ♂ imm., 1 ♀ ad. 15-VII. 1929. Ohōsen.
 1 ♂ ad. 24. II. 1930. Moppo.
228. *Streptopelia decaocto decaocto* (Frivaldsky). シラコバト
 1 unsexed ad. VIII. 1929. Banpo.

Family CHARADRIIDÆ. 千鳥科

229. *Charadrius dubius curonicus* Gmelin. コチドリ
 1 ♂ ad., 3 ♀♀ ads., 1 unsexed ad. 6.-12. IV. 1929. Riuganpo.
230. *Charadrius placidus* Gray. イカルチドリ
 4 ♂♂ ads., 1 ♀ ad. 30. XI.-4. XII. 1929. Kinkwa.
231. *Charadrius mongolus mongolus* Pallas. メダイチドリ
 1 ♀ ad. 8. V. 1929. Riuganpo.
 3 ♂♂ ads., 2 ♀♀ ads. 17.-19. IX. 1929. Banpo.
232. *Pluvialis apricarius fulvus* (Gmelin). ムナグロ
 1 unsexed ad. 14. IV. 1929. Riuganpo.
 2 ♂♂ ads., 2 ♀♀ ads. 17.-18. IX. 1929. Banpo.
233. *Squatarola squatarola hypomelaena* (Pallas). ダイゼン
 3 ♀♀ ads. 15.-29. IV. 1929. Riuganpo.
 1 ♂ ad., 2 ♀♀ ads. 17.-26. IX. 1929. Banpo.
234. *Vanellus vanellus* (Linnaeus). タゲリ
 1 ♂ imm., 1 ♀ ad. 13. X-1. XI. 1929. Banpo.
235. *Haematopus ostralegus osculans* Swinhoe. ミヤコドリ
 1 ♂ ad. 23. IV. 1929. Riuganpo.
 1 ♀ ad. 7. II. 1930. Moppo.
 2 ♂♂ ads. 15. II. 1930. Chitō.
236. *Arenaria interpres interpres* (Linnaeus). キョウジョシギ
 1 ♂ imm., 2 ♀♀ ads. 19.-22. IX. 1929. Banpo.

Family SCOLOPACIDÆ. 鷸科

237. *Erolia alpina sakhalina* (Vieillot). ハマシギ
 3 ♂♂ ads., 2 ♀♀ ads. 25. IV. 1929. Riuganpo.
 2 ♂♂ ads., 3 ♀♀ ads. 16.-19. IX. 1929. Banpo.
 1 ♀ ad. 3. I. 1930. Moppo.

Among the above specimens ♀ ad. (19. IX. 1929, Banpo) is a full albino.

238. *Pisobia minuta ruficollis* (Pallas). トウネン
 3 ♂♂ ads., 2 ♀♀ ads. 25. IV.-13. V. 1929. Riuganpo.
 239. *Pisobia minutilla subminuta* (Middendorff). ヒバリシギ
 3 ♂♂ ads., 2 ♀♀ ads. 9.-19. V. 1929. Riuganpo.
 240. *Pisobia maculata* (Vieillot). アメリカカウヅラシギ
 1 ♂ imm., 2 ♀♀ imms. 19.-25. IX. 1929. Banpo.

About the specimen, ♀ imm., 25. IX. 1929, Banpo, among the above-mentioned specimens, I have not described in the "Tori," Vol. VI, p. 252.

241. *Pisobia acuminata* (Horsfield). ウヅラシギ
 4 ♀♀ ads. 19.-25. V. 1929. Riuganpo.
 242. *Calidris tenuirostris* (Horsfield). ラバシギ
 1 ♂ imm., 1 ♀ ad., 1 ♀ imm. 19.-23. IX. 1929. Banpo.
 243. *Crocethia alba* (Pallas). ミユビシギ
 2 ♂♂ ads., 3 ♀♀ ads. 13. IX. 1929. Banpo.
 244. *Limicola falcinellus sibirica* Dresser. キリアイ
 1 ♀ ad. 16. IX. 1929. Banpo.
 245. *Glottis nebularius* (Gunnerus). アラアシシギ
 3 ♂♂ ads., 1 unsexed ad. 2.-14. V. 1929. Riuganpo.
 1 ♂ ad., 3 ♀♀ ads., 1 ♀ imm. 16. IX.-9. X. 1929. Banpo.
 246. *Tringa erythropus* (Pallas). ツルシギ
 1 ♀ ad. 11. IV. 1929. Riuganpo.
 3 ♀♀ imms. 21. IX.-18. X. 1929. Banpo.
 247. *Tringa totanus eurhinus* (Oberholser). アカアシシギ
 3 ♂♂ imms., 1 ♀ ad., 1 ♀ imm. 17.-19. IX. 1929. Banpo.
 248. *Tringa ochropus* Linnaeus. クサシギ
 1 ♂ ad. 1. X. 1929. Banpo.
 1 ♀ ad. 27. XII. 1929. Moppo.
 249. *Tringa glareola* Linnaeus. タカブシギ
 4 ♂♂ ads., 1 ♀ ad. 24. IV.-10. V. 1929. Riuganpo.
 1 ♀ ad. 9. X. 1929. Banpo.
 250. *Tringa incana brevipes* (Vieillot). キアシシギ

- 1 ♀ ad. 4. X. 1929. Banpo.
251. *Tringa hypoleucos* Linnaeus. イノシギ
1 ♀ imm. 19. IX. 1929. Banpo.
252. *Lobipes lobatus* (Linnaeus). アカエリヒレアシシギ
2 ♂♂ ads., 3 ♀♀ ads. 15.-17. IX. 1929. Banpo.
253. *Limosa limosa melanuroides* Gould. ラグロシギ
3 ♂♂ ads., 2 ♀♀ ads. 13. IX. 1929. Banpo.
254. *Limosa lapponica baueri* Naumann. オホソリハシシギ
5 ♂♂ imm. 17. IX. 1929. Banpo.
255. *Numenius arquata orientalis* Brehm. ダイシャクシギ
2 ♂♂ ads., 1 unsexed ad. 17. I.-14. II. 1930. Moppo.
256. *Numenius cyanopus* Vieillot. ホウロクシギ
1 ♂ ad., 1 ♀ ad. 14.-19. V. 1929. Riuganpo.
257. *Numenius phaeopus variegatus* (Scopoli). チウシャクシギ
3 ♂♂ ads., 1 ♀ ad. 14. IV.-2. V. 1929. Riuganpo.
258. *Numenius minutus* Gould. コシャクシギ
1 ♂ ad., 1 ♀ ad. 29.-30. IV. 1929. Riuganpo.
- The measurement of ♂♀ above mentioned is as follows:
Wing ♂ 176 ♀ 185, culmen ♂ 42 ♀ 47.
259. *Scolopax rusticola rusticola* Linnaeus. ヤマシギ
1 ♂ ad. 9. IV. 1929. Riuganpo.
- The measurement of this specimen is, wing 185, culmen 72.
260. *Capella gallinago raddei* (Buturlin). タシギ
3 ♂♂ ads., 1 ♀ ad. 12.-18. IV. 1929. Riuganpo.
1 ♀ ad. 14. III. 1930. Quelpart I.
261. *Capella stenura* (Bonaparte). ハリヲシギ
1 ♂ ad., 4 ♀♀ ads. 10.-11. V. 1929. Riuganpo.
1 ♀ ad. 6. VIII. 1929. Nōjidō.
- Family LARIDÆ. 鷗科
262. *Sterna hirundo longipennis* Nordmann. アジサシ
2 ♂♂ juvs., 1 ♀ juv. 25. IX. 1929. Banpo.
263. *Larus argentatus vegae* Palmén. セグロカモメ
1 unsexed ad., 2 subad., 1 imm. 13. III. 1930. Quelpart I.

264. *Larus canus major* Middendorff. カモメ

1 ♀ ad. 19. II. 1930. Moppo.

265. *Larus crassirostris* Vieillot. ウミネコ

1 unsexed ad. 14. IV. 1929. Riuganpo.

266. *Larus ridibundus sibiricus* Buturlin. ユリカモメ

1 ♂ ad., 3 ♀ ♀ ads., 2 ♀ ♀ 2nd years. 15.-27, IX, 1929 Banpo

I have described in the "Tori," Vol. VI, p. 253, that *L. r. ridibundus* live in Korea, too. But according to the recent study, it is wrong to classify *L. ridibundus* which come to Japan into two subspecies, the difference of size, which has hitherto been regarded as the main point of the classification, being only an individual variation. Further investigation, however, is required to determine which name we can rightly apply to them, *L. r. ridibundus* or *L. r. sibiricus*.

Family BALEARICIDÆ. 鶴科

267. *Grus monachus* Temminck. ナベヅル

3 ♀ ♀ ads., 2 ♀ ♀ imms. 15. I.-11. II. 1930. Moppo.

268. *Grus vipio* Pallas. マナヅル

1 ♀ imm. 29. I. 1930. Moppo.

Family RALLIDÆ. 秧鶴科

269. *Porzana pusilla pusilla* (Pallas). ヒメクヒナ

1 ♀ ad. 30. V. 1929. Riuganpo.

1 ♂ ad. 16. X. 1929. Banpo.

270. *Porzana fusca erythrothorax* (T. & S.). ヒクヒナ

1 ♂ ad. 13. VII. 1929. Chōsen.

271. *Porzana paykullii* (Ljungh). カウライヒクヒナ

♀ ad. 26. V. 1929. Riuganpo.

272. *Gallinula cinerea* (Gmelin). ツルクヒナ

2 ♂ ♂ ads. 31. V.-3. VI. 1929. Riuganpo.

2 ♂ ♂ ads., 1 ♀ ad. 25.-29. VI. 1929. Chōsen.

273. *Fulica atra atra* Linnaeus. オホバン

2 ♂ ♂ ads., 1 ♀ ad. 15.-20. IX. 1929. Banpo.

Family TURNICIDÆ. 三班鶉科

274. *Turnix tanki blanfordi* Blyth. テウセンミフウヅラ
2 ♀ ♀ ads. 23.-24. V. 1929. Riuganpo.

Family TETRAONIDÆ. 松 鶉 科

275. *Lyrurus tetrix ussuriensis* (Kohte). クロライテウ

Synonym: *L. t. koreensis* Mori.

- 4 ♂ ♂ ads., 3 ♀ ♀ ads., 4 ♀ ♀ imms. 11.-26. VIII. 1929. Nōjidō.
276. *Tetrastes bonasia coreensis* Kuroda & Mori.

テウセンエゾライテウ

- 1 ♂ ad., 3 juvs. 6.-9. VII. 1929. Chōsen.
1 ♂ ad., 1 juv. 25. VIII-1. IX. 1929. Nōjidō.

Family PHASIANIDÆ. 雉 科

277. *Coturnix coturnix japonica* T. & S. ウ ズ ラ

- 1 ♂ ad., 1 ♀ ad. 4.-11. V. 1929. Riuganpo.
2 ♂ ♂ ads. 15.-18. VIII. 1929. Nōjidō.
1 ♂ ad., 1 ♀ ad. 22. IX.-22. X. 1929. Banpo.
1 ♀ ad. 17. I. 1930. Moppo.

278. *Phasianus colchicus karpowi* Buturlin. カウライキジ

Synonym: *P. c. quelpartis* Momiyama.

- 2 ♂ ♂ ads., 2 ♀ ♀ ads. 18. XII. 1929-31. I. 1930. Moppo.
2 ♂ ♂ ads., 1 ♀ ad. 9-30. III. 1930. Quelpart I.
279. *Phasianus colchicus pallasi* Rothschild. ホクマンキジ
1 ♀ ad., 6 juvs. 27. VII-11. VIII. 1929. Nōjidō.
3 ♂ ♂ ads., 1 ♀ ad., 1 ♀ imm. 27. IX-23. X. 1929. Banpo.