

Effects of the Thermal Environment on Articular Chondrocyte Metabolism: A Fundamental Study to Facilitate Establishment of an Effective Thermotherapy for Osteoarthritis

Akira ITO, PT^{1,2}, Tomoki AOYAMA, MD, PhD³, Junichi TAJINO, PT¹, Momoko NAGAI, PT¹,
Shoki YAMAGUCHI, PT¹, Hirotaka IJIMA, PT¹, Xiangkai ZHANG¹, Haruhiko AKIYAMA, MD, PhD⁴,
Hiroshi KUROKI, PT, PhD¹

¹⁾ Department of Motor Function Analysis, Human Health Sciences, Graduate School of Medicine, Kyoto University, Kyoto 606-8507, Japan

²⁾ Research Fellow of Japan Society for the Promotion of Science, Tokyo 102-0083, Japan

³⁾ Department of Development and Rehabilitation of Motor Function, Human Health Sciences, Graduate School of Medicine, Kyoto University, Kyoto 606-8507, Japan

⁴⁾ Department of Orthopaedic Surgery, Graduate School of Medicine, Gifu University, Gifu 501-1194, Japan

ABSTRACT. Aim: To facilitate establishment of an effective thermotherapy for osteoarthritis (OA), we investigated the effects of the thermal environment on articular chondrocyte metabolism *in vitro*. Methods: Chondrocytes were isolated from porcine knee joints, and cultured at 32°C, 37°C and 41°C. Cell proliferation and viability were assessed at Days 2, 4 and 8. In addition, TdT-mediated dUTP nick end labeling (TUNEL) assay was performed at Day 3 to determine the proportion of apoptotic chondrocytes. Analysis of genes specific for factors related to the cartilage extracellular matrix (ECM), cartilage destruction, and cartilage protection was performed at Day 2. Furthermore, evaluation of heat stress tolerance, and heat shock protein 70 (HSP70) mRNA expression and protein synthesis was performed at Day 2 and 3, respectively. Results: Cell proliferation was more at 37°C than at 32°C and 41°C. Cell viability and the number of TUNEL-positive cells were not affected until Day 8 and 3, respectively. The expression of the ECM-related genes was up-regulated at higher temperature. The expression of MMP13, a type II collagen destructive enzyme, and that of TIMP1 and TIMP2, which are MMP inhibitors, were up-regulated at higher temperatures. Finally, the chondrocytes cultured at 41°C may acquire heat stress tolerance, in part, due to the up-regulation of HSP70, and may inhibit apoptosis induced by various stresses, which is observed in OA. Conclusions: The thermal environment affects articular chondrocyte metabolism, and a heat stimulus of approximately 41°C could enhance chondrocyte anabolism and induce heat stress tolerance.

Key words: Thermal environment, articular chondrocyte metabolism, thermotherapy

(*J Jpn Phys Ther Assoc* 17: 14–21, 2014)

Osteoarthritis (OA) is a highly prevalent degenerative disease¹⁾, characterized by progressive articular cartilage destruction, which results in pain and decreased physical

function. While many approaches for treating various aspects of OA have been studied, there exists no established effective treatment, because severely degenerated articular cartilage cannot be regenerated, and the specific factors related to OA development are complex^{2–4)}.

Physical therapy is often used as a first-line treatment in the management of OA symptoms. Since it has been reported that exercise intervention has a therapeutic effect

Received: January 15, 2014

Accepted: April 7, 2014

Advance Publication by J-STAGE: May 28, 2014

Correspondence to: Hiroshi Kuroki, 53 Kawahara-cho, Shogoin, Sakyo-ku, Kyoto 606-8507, Japan

e-mail: kuroki.hiroshi.6s@kyoto-u.ac.jp

on pain and physical function^{5,6}, exercise therapy has been recommended in OA treatment guidelines^{7,8}. On the other hand, the therapeutic effect of physical therapy modalities for OA treatment is still under debate^{7,8}. In particular, clear evidence of the therapeutic effect of thermotherapy has not been shown^{9,10}. However, in a previous study, application of a heat stimulus to treat OA within an experimental animal model, showed inhibition of OA development^{11,12}. In addition, when women with chronic knee pain were treated with long-term thermotherapy in a randomized controlled trial, their pain was found to decrease; this reduction in pain and improvement of physical function was attributed to the combination of thermotherapy and exercise therapy, rather than to the exercise therapy alone¹³. These reports suggest that thermotherapy could be an effective treatment for OA if the temperature of the heat stimulus and the duration of exposure are optimized and if the target region for heat stimulus application is selected appropriately. However, the knowledge used as the basis for setting up each condition, which in many cases is based on empirical rules and not on objective data, makes it difficult to properly setup conditions. Furthermore, existing thermotherapy research for OA treatment has considered the effects on pain, blood flow, and physical function, but, there is little research which has taken into consideration the influence on the articular cartilage, which is a primary point of pathological OA processes, or the articular chondrocytes which play a key role in the metabolic activities of the articular cartilage.

Therefore, the purpose of this study was to facilitate the establishment of an effective thermotherapy for OA by clarifying the effects of the thermal environment on articular chondrocyte metabolism. In particular, in this study we focused on the effects of the thermal environment on cell proliferation, cell death, gene expression, and heat stress tolerance for articular chondrocytes.

Materials and Methods

The protocols described for the following experiments did not require approval of the animal research committee.

Chondrocyte isolation and expansion

Knee joints of 6-month-old pigs were purchased from a meat processor. Articular cartilage plugs were aseptically harvested from the femoral condyle using a biopsy punch, and chondrocytes were isolated as previously described¹⁴. The isolated cells were suspended in Dulbecco's modified Eagle medium/Ham's F12 (Nacalai Tesque Inc., Kyoto, Japan) containing 10% fetal bovine serum (Hyclone, Logan, USA), 50 U/mL penicillin (Nacalai Tesque Inc.), and 50 μ g/mL streptomycin (Nacalai Tesque Inc.), were seeded in culture dishes. The chondrocytes were expanded in a CO₂ incubator (5% CO₂, 37°C, 95% humidity) to obtain a sufficient quantity of cells (2–3 passages). To investigate the

effects of the thermal environment on chondrocyte metabolism, the chondrocytes were cultured at the following three conditions: 32°C as normal intra-articular temperature^{15,16}, 37°C as internal body temperature, and 41°C as the threshold temperature for mammalian cell survival^{17,18}.

Cell proliferation, viability, and apoptosis

To assess cell proliferation and viability at the three different culturing temperatures, the expanded chondrocytes were divided into 27 culture dishes at 8.5×10^3 cells/cm², and then pre-cultured overnight at 37°C in a CO₂ incubator. After pre-culture, the dishes were transferred into three distinct CO₂ incubators set at 32°C, 37°C, and 41°C, 9 dishes for each. The cells were trypsinized from the culture dishes at Days 2, 4, and 8 (3 dishes for each day; n = 3). The number of live chondrocytes was counted using a hemocytometer, and cell viability was measured by using a trypan blue dye exclusion test. This experiment was repeated again using cells harvested from another pig to confirm the accuracy and reproducibility of the results. Additionally, a TdT-mediated dUTP nick end labeling (TUNEL) assay was performed to analyze the number of apoptotic chondrocytes according to the manufacturer's protocol (In Situ Apoptosis Detection Kit; Takara Bio Inc., Shiga, Japan). Briefly, the expanded chondrocytes were sub-cultured into 2-well Lab-Tek chamber slides (Nalge Nunc International, Naperville, USA) at 7.2×10^3 cells/cm². After pre-culture as described above, the cells were cultured at 32°C, 37°C, and 41°C for 3 days. Subsequently, the cells were fixed in 4% paraformaldehyde for 15 min. After digestion with a permeabilization buffer, the slides were treated with TdT enzyme and incubated at 37°C for 60 min, then further incubated at 37°C for 30 min with an anti-FITC HRP Conjugate and propidium iodide, which were used as counter stains. More than 3,000 TUNEL-positive and TUNEL-negative cells were counted from more than 6 views ($1272.8 \mu\text{m} \times 1272.8 \mu\text{m}/\text{view}$) that were randomly captured using a confocal laser scanning microscope system (FluoView FV10i; OLYMPUS CO., Tokyo, Japan). The proportion of TUNEL-positive cells/view was then calculated ($n \geq 6$).

Gene expression analysis

Analysis of genes related to cartilage extracellular matrix (ECM) and cartilage-destroying/protecting factors was performed. The 9 culture dishes of expanded chondrocytes were transferred into incubators at 32°C, 37°C, and 41°C (3 dishes for each; n = 3), and incubated for 2 days. Total RNA was extracted using the RNeasy Mini Kit according to the manufacturer's protocol (Qiagen Inc., Valencia, USA). Extracted total RNA was dissolved in water and tested for purity using a NanoDrop2000 (Thermo Scientific, Wilmington, USA). The gene expression level was quantified by real-time reverse transcription-polymerase chain reaction (RT-PCR). Total RNA (300 ng) was reverse-transcribed to

Table 1. Primer sequences for real-time RT-PCR

	Sense (5'–3')	Antisense (5'–3')	Length (bp)
COL2A1	GCTATGGAGATGACAACCTGGCTC	CACTTACCGGTGTGTTTCGTGCAG	256
COL1A1	CAGAACGGCCTCAGGTACCA	CAGATCACGTCATCGCACAAAC	101
aggrecan	GAATTTCTTGGCGTGAGAAC	GGGGATGTTGCGTAAAAGAC	107
SOX9	GTACCCGCACCTGCACAAC	GACTGCTGAATGAGAGCGAGA	68
MMP13	CACCCGTGACCTTATCTTCATC	GCTGCGCTTATCCTTTTAACC	132
MMP1	GCCAAATGGACTTCAAGCTG	AGCCAAAGGATCTGTGGATG	137
TIMP1	ACCACCTGCAGTTTTGTGG	AGTTTGCAGGGGATGGATG	134
TIMP2	GAACGACATCTACGGCAACC	TTCTTTCCTCCGATGTCCAG	150
HSP70	TCACCATCACCAACGACAAG	TCATGTTGAAGGCGTACGAC	147
Beta-actin	AAGCCAACCGTGAGAAGATG	TCCATCACGATGCCAGTG	124

synthesize cDNA, and then real-time PCR was performed using the Applied Biosystems 7500 Real-Time PCR System (Life Technologies Corporation, Carlsbad, USA). cDNA templates were amplified with PowerSYBR Green PCR Master Mix (Life Technologies Corporation) in 25- μ L reactions with 0.2 μ M gene-specific primers (Table 1) and de-ionized water. The PCR reaction was performed for 10 min at 95°C, followed by 40 amplification cycles (15 s at 95°C, 60 s at 60°C). The following target genes were examined: the ECM-related markers collagen type IIA1 (COL2A1), aggrecan, SRY (sex-determining region Y)-box 9 (SOX9), and collagen type IA1 (COL1A1); the cartilage-degrading factors matrix metalloproteinase (MMP)-1 and MMP-13; and the MMP-inhibitory factors tissue inhibitor of metalloproteinase (TIMP)-1 and TIMP2. Beta-actin was used as the control housekeeping gene. This experiment was repeated again using cells harvested from another pig to confirm the accuracy and reproducibility of the results.

The data obtained by real-time PCR were analyzed using the comparative threshold cycle method. Briefly, the amount of target was normalized to that of beta-actin, the value of the calibration sample (the cells cultured at 32°C) was set to 1, and the values for each of the other conditions were shown relative to that of the calibration sample. Before using the comparative threshold cycle method for quantitation, we performed a validation experiment, and the obtained absolute value of which the slope of log input amount versus delta threshold cycle was less than 0.1.

Heat shock protein 70 synthesis

The 18 culture dishes of expanded chondrocytes were transferred into incubators maintained at temperatures of 32°C, 37°C, and 41°C, and incubated for 2 or 3 days for the analysis of heat shock protein 70 (HSP70) mRNA expression and HSP70 protein synthesis, respectively (3 dishes for each: $n = 3$). The HSP70 mRNA expression was analyzed by the method described above. HSP70 protein synthesis was evaluated using western blotting. Cell lysates were

prepared in SDS-sample buffer (70 mM Tris-HCl, pH 6.8, 11.2% glycerol, 3% SDS, 0.01% bromophenol blue, and 5% 2-mercaptoethanol). Equal amounts of protein (2 μ g) were loaded on 10% SDS-polyacrylamide gels and blotted on a polyvinylidene fluoride microporous membrane using a semi-dry transfer system (Bio-Rad, Hercules, USA). The primary antibodies used were as follows: rabbit anti-HSP70 polyclonal antibody (StressGen Biotech, Victoria, Canada), and mouse anti-beta-actin monoclonal antibody (BioVision, Mountain View, USA). Blots were probed with HRP-conjugated donkey anti-rabbit IgG (GE Healthcare, Little Chalfont, England) for HSP70, and with sheep anti-mouse IgG (GE Healthcare) for beta-actin, and then visualized using an ECL method (GE Healthcare). Protein levels were quantified using Image Lab software (Bio-Lad). The expression level of HSP70 was normalized to that of beta-actin, and was calculated by comparing to the expression level of 32°C samples.

Heat stress tolerance

Heat stress tolerance of the cells cultured at 32°C and 41°C was analyzed. 37°C condition was excluded in this experiment, since the expression of the HSP70 on the mRNA and the protein level didn't differ from 32°C. The chondrocytes cultured at 32°C or 41°C for 7 days were sub-cultured into 2-well Lab-Tek chamber slides at 1.2×10^4 cells/cm². After pre-culture as previously described, the cells were exposed to excessive heat stress at 48°C for 1 h on a hot plate and then incubated at 37°C for 1 day. Subsequently, a TUNEL assay was performed as described above. The cells were counted from 20 random views (1272.8 μ m \times 1272.8 μ m/view), and then the cell number/view and the proportion of the TUNEL-positive cell/view were calculated ($n = 20$).

Statistical analysis

All values are reported as means \pm standard deviation (SD). Statistical significance was determined using

unpaired Student's *t* test or one-way analysis of variance (ANOVA) with the post-hoc multiple comparison Tukey-Kramer test. The differences observed were considered to be significant if the *p* value was less than 0.05.

Results

The effects of thermal environment on cell proliferation, viability, and apoptosis

The cell proliferation, viability, and apoptosis induction at three different culturing temperatures were assessed. The cell number increased at each temperature, however, the cells were more proliferative at 37°C than at 32°C and 41°C (Fig. 1A). There were no significant differences between 32°C and 41°C at Day 4 and 8, although 41°C showed a lower number of cells than 32°C at Day 2. There were no significant differences in the cell viabilities (Fig. 1B) and the proportion of TUNEL-positive cells (Fig. 1C) among the three examined temperatures. Under these experimental conditions, the viability remained high (more than 94.5%), and the proportion of TUNEL-positive cells remained low (less than 1%).

Gene expression analysis

The expression of genes related to the ECM, cartilage-degrading factors, and cartilage-protecting factors were analyzed at each temperature by real time PCR. The expression of genes related to the ECM tended to increase in a temperature-dependent manner. Specifically, COL2A1 mRNA expression at 41°C was approximately 28 times the expression at 32°C (Fig. 2A). COL1A1 mRNA expression was also higher at higher temperatures, although increase in expression was apparently less than that of COL2A1 (Fig. 2B). Additionally, aggrecan mRNA expression was up-regulated at 37°C and 41°C (Fig. 2C), and SOX9 mRNA expression at 41°C was up-regulated approximately 6 times that of the 32°C samples (Fig. 2D). Interestingly, MMP1 and MMP13, which are cartilage-degrading factors, showed different trends with respect to each other. The expression of MMP1 mRNA was higher at lower temperatures (Fig. 2E), while expression of MMP13 mRNA was higher at higher temperatures (Fig. 2F). The mRNA expression of both TIMP1 (Fig. 2G) and TIMP2 (Fig. 2H), which are MMP-inhibitory factors, was up-regulated at 41°C and was up-regulated in a temperature-dependent manner.

HSP70 synthesis and heat stress tolerance

The HSP70 mRNA and the HSP70 protein were analyzed by real-time RT-PCR and Western blotting, respectively. A significant increase in mRNA (Fig. 3A) and protein levels (Fig. 3B) at 41°C was observed, although a significant difference was not detected between 32°C and 37°C.

In order to clarify articular chondrocyte heat stress tolerance, the cell death after exposure to an excessive heat




Fig. 1. Effects of the thermal environment on cell proliferation, viability, and apoptosis. (A) Cell number and (B) cell viability when cultured at 32°C, 37°C, and 41°C for 2, 4 and 8 days (*n* = 3). (C) The proportion of TdT-mediated dUTP nick end labeling (TUNEL)-positive cells. The TUNEL assay was performed to determine the proportion of apoptotic chondrocytes cultured at 32°C, 37°C, and 41°C for 3 days (*n* ≥ 6). All values represent the means and standard deviations (SD). ***P* < 0.01. N.S.; not significant.

stimulus (48°C) was evaluated. There was a significantly larger number of cells/view at 41°C (Fig. 4A). Moreover, the proportion of TUNEL-positive cells was significantly higher at 32°C (Fig. 4B).

Discussion

We investigated the effects of thermal environment on chondrocyte metabolism from a viewpoint of cell prolifera-




Fig. 2. Gene expression analysis. Relative mRNA expression of (A) COL2A1, (B) COL1A1, (C) aggrecan, (D) SOX9, (E) MMP1, (F) MMP13, (G) TIMP1, and (H) TIMP2 cultured at 32°C, 37°C, and 41°C for 2 days are shown ($n = 3$). Values represent the means and SD. $*P < 0.05$. $**P < 0.01$.

tion, cell death, gene expression, and heat stress tolerance.

Our results indicated that approximately 37°C (near internal body temperature) is the most suitable temperature for cell proliferation (Fig. 1A). The more the temperature deviates from the internal body temperature, the more the cellular proliferative potential is inhibited. To the best of our knowledge, there are few published reports demonstrating that mammalian cells cultured at temperatures below 37°C proliferate more than those cultured at 37°C, with all other conditions being identical. There is only one report indicating that oligodendrocyte precursor cells cultured at 31.5°C resulted in an increase of the cell number relative to the cells cultured at 37°C¹⁹. Generally, reduced metabolic rate and expression of cold shock response genes, caused by low temperatures lead to reduction in the growth rate²⁰. It has also been reported that cell cycle arrest has occurred when cells were exposed to high temperature (42°C)^{21,22}. Our results are consistent with these previous reports. Although the mature chondrocyte is seldom proliferating in the living body, in a process of cartilage regeneration, there is a possibility that 37°C is suitable when cell growth is required.

In research on the cell death, acute reactions caused by heat stimuli have been well-studied^{23,24}. Wheatley *et al.*¹⁸ reported that mammalian cells seemed to tolerate

temperatures of 42°C, whereas 43°C appeared to be a critical threshold, which rapidly led to cell death. Hojo *et al.*²⁵ have reported that heat stimulus at 41°C for 15 or 30 min increased the viability of chondrocytes, and the borderline temperature that determines the increase or decrease of their metabolism would be between 41°C and 43°C. Moreover, they have indicated that these effects vary, not only according to temperature, but also according to the duration of exposure. However, the influence of chronic exposure is still unclear²⁴. In this study, we demonstrated that the cell viability of the cartilage chondrocytes did not change in the temperature of the range of 32–41°C until Day 8 of culture (Fig. 1B). The proportion of TUNEL-positive cells at Day 3 was also low at < 1% of the total cells (Fig. 1C). Therefore, at temperatures from 32–41°C, chondrocytes may not experience cell death, even if exposed for a long period of time.

The expression of the genes related to cartilage ECM were up-regulated in a temperature-dependent manner (Fig. 2). This suggested that the production of cartilage ECM was enhanced by heat stimulus. Interestingly, MMP1 and MMP13 showed an opposite expression trend. MMP1 tended to be down-regulated while MMP13 tended to be up-regulated in a temperature-dependent manner. Although




Fig. 3. Heat shock protein 70 synthesis. (A) Relative mRNA expression of heat shock protein 70 (HSP70) analyzed by real-time PCR and (B) relative protein synthesis level analyzed by western blotting when cultured at 32°C, 37°C, and 41°C for 2 days and 3 days, respectively, are shown ($n = 3$). Values represent the means and SD. * $P < 0.05$. ** $P < 0.01$.

this up-regulation of MMP13, which is a specific degradation factor of type II collagen protein, indicates the induction of cartilage degeneration, both TIMP1 and TIMP2, which are cartilage-protective factors, were also up-regulated in a temperature-dependent manner. It has been reported that TIMP proteins combine with MMP proteins and inactivate the function of the MMP proteins²⁶. Therefore, it is necessary to consider the relative amounts of both MMP and TIMP proteins. However, these amounts cannot be assessed at the protein level, but only at the mRNA level in this study.

The MMP1 mRNA expression, which is of a different MMP family, was higher at 32°C in our study conditions, however, it has been reported that the mRNA expression of MMP1, induced by heat stimulus, was up-regulated in a temperature-dependent manner in human skin fibroblasts and epidermal keratinocytes^{27,28}. Although the detailed mechanism was unknown, these results imply that the influence of the thermal environment on MMP1 mRNA ex-




Fig. 4. Heat stress tolerance. The chondrocytes cultured at 32°C or 41°C for 7 days were exposed to excessive heat stimulus at 48°C for 1 h on a hot plate. (A) The cell number and (B) the proportion of TUNEL-positive cells 1 day following heat stimulus were counted ($n = 20$). Values represent the means and SD. * $P < 0.05$.

pression may differ depending on the cell species.

The cells cultured at 41°C were more tolerant to heat stress than those cultured at 32°C (Fig. 4A, B). It was assumed that the acquisition of this heat tolerance was related to induction of the HSP family, mainly HSP70. HSPs are a vital set of chaperone proteins that respond to thermal stress by assisting denatured proteins in the cytosol to refold into their native, functional conformations, thereby restoring the homeostasis of the cell²⁹. It has been reported that the stress tolerance by HSPs shows tolerance to not only heat stress³⁰, but also to oxidant stress³¹, and mechanical stress³². Moreover, these effects were confirmed not only in the articular cartilage^{11,12} but also in various organs such as skeletal muscles^{33–35}, liver³⁶, and blood vessels³⁷. Therefore, the heat stimulus at 41°C might reduce the apoptosis of chondrocytes, which has been reported as a part of OA pathology, through the induction of HSPs.

We identified five limitations of this study. First, our results were obtained from *in vitro* experiments. Since the same result may not be obtained *in vivo*, it is necessary to verify these findings using animal experiments, and to also further confirm in a clinical study. Second, the heat stress tolerance acquired by the heat stimulus did not verify

whether the tolerance is actually shown to stress other than the excessive heat stress used in this study, even though it has generally been reported that tolerance is shown to various other stressors. Third, we used 6-month-old pigs in this study, which are skeletally immature and not osteoarthritic, and it is likely that juvenile chondrocytes will react differently to temperature than adult chondrocytes. Fourth, our results assessing the metabolic change were obtained from basically mRNA level, not protein level. Since mRNA expression level does not always correlate with protein synthesis level, we should have also confirmed the effects of the thermal environment on articular chondrocyte metabolism in the protein level. Finally, the effects on other cell types that exist in the constituted synovial joint were not investigated. It has been reported that the metabolic reaction and the thermotolerance will differ among cell types^{38–40}. To establish a safe and effective thermotherapeutic approach, it is essential to investigate its influence on chondrocytes and also all other cell types.

Conclusions

We investigated the effects of the thermal environment on articular chondrocyte metabolism *in vitro* to facilitate the establishment of an effective thermotherapy for OA. Several points were suggested by our results. First, articular chondrocyte cell death is not promoted even when cells are cultured at 32–41°C for a long period of time in a monolayer culture. Second, cell proliferation is promoted at 37°C, which is near internal body temperature, and is inhibited at temperatures higher or lower than 37°C. Third, in an environment with temperature higher than 32°C, which is the normal intra-articular temperature, the expression of the genes related to the ECM is up-regulated, and the ECM production may be enhanced. Fourth, although the expression of MMP13, the main destructive enzyme of type II collagen, is up-regulated in a temperature-dependent manner, the expression of TIMP1 and TIMP2, which are MMP inhibitors, are also up-regulated in a similar manner. Therefore, it cannot be simply concluded that a high-temperature environment promotes cartilage destruction. Finally, the chondrocytes cultured at 41°C may acquire heat stress tolerance, in part, due to the up-regulation of HSP70, and thus may inhibit the apoptosis of the chondrocytes induced by various stresses, which is observed in OA. Taken together, thermal environment could affect articular chondrocyte metabolism, and heat stimulus at approximately 41°C may enhance chondrocyte anabolism and induce heat stress tolerance.

Acknowledgements

This study was supported in part by Grant-in-Aid for JSPS Research Fellows (number 820130600018), JSPS

KAKENHI Grant-in-Aid for Scientific Research (A) (number 25242055), and JSPS KAKENHI Grant-in-Aid for Challenging Exploratory Research (number 25560258).

Declaration of interest

The authors declare no competing interests associated with manuscript.

References

- 1) Muraki S, Oka H, Akune T, Mabuchi A, En-Yo Y, Yoshida M, Saika A, Suzuki T, Yoshida H, Ishibashi H, Yamamoto S, Nakamura K, Kawaguchi H, and Yoshimura N: Prevalence of radiographic knee osteoarthritis and its association with knee pain in the elderly of Japanese population-based cohorts: The road study. *Osteoarthritis Cartilage*. 2009, 17: 1137–1143.
- 2) Kim HK, Moran ME, and Salter RB: The potential for regeneration of articular cartilage in defects created by chondral shaving and subchondral abrasion. An experimental investigation in rabbits. *J Bone Joint Surg Am*. 1991, 73: 1301–1315.
- 3) Hayes DW Jr., Brower RL, and John KJ: Articular cartilage. Anatomy, injury, and repair. *Clin Podiatr Med Surg*. 2001, 18: 35–53.
- 4) Hunziker EB: Articular cartilage repair: Basic science and clinical progress. A review of the current status and prospects. *Osteoarthritis Cartilage*. 2002, 10: 432–463.
- 5) Fransen M, and McConnell S: Exercise for osteoarthritis of the knee. *Cochrane Database Syst Rev*. 2008: CD004376.
- 6) Uthman OA, Van Der Windt DA, Jordan JL, Dziedzic KS, Healey EL, Peat GM, and Foster NE: Exercise for lower limb osteoarthritis: systematic review incorporating trial sequential analysis and network meta-analysis. *BMJ*. 2013, 347: f5555.
- 7) Zhang W, Nuki G, Moskowitz RW, Abramson S, Altman RD, Arden NK, Bierma-Zeinstra S, Brandt KD, Croft P, Doherty M, Dougados M, Hochberg M, Hunter DJ, Kwoh K, Lohmander LS, and Tugwell P: OARSI recommendations for the management of hip and knee osteoarthritis: Part III: changes in evidence following systematic cumulative update of research published through January 2009. *Osteoarthritis Cartilage*. 2010, 18: 476–499.
- 8) Jevsevar DS: Treatment of osteoarthritis of the knee: evidence-based guideline, 2nd ed. *J Am Acad Orthop Surg*. 2013, 21: 571–576.
- 9) Brosseau L, Yonge KA, Robinson V, Marchand S, Judd M, Wells G, and Tugwell P: Thermotherapy for treatment of osteoarthritis. *Cochrane Database Syst Rev*. 2003: CD004522.
- 10) Moriyama H, Iijima H, Kanemura N, Murata K, Hata Y, Nishihara K, Ozawa J, Takayanagi K, Gomi T, and Tobimatsu Y: Efficacy of thermotherapy on musculoskeletal disorders. *J Phys med*. 2009, 20: 260–268. (Japanese)
- 11) Tonomura H, Takahashi KA, Mazda O, Arai Y, Shin-Ya M, Inoue A, Honjo K, Hojo T, Imanishi J, and Kubo T: Effects of heat stimulation via microwave applicator on cartilage matrix gene and hsp70 expression in the rabbit knee joint. *J Orthop Res*. 2008, 26: 34–41.
- 12) Fujita S, Arai Y, Nakagawa S, Takahashi KA, Terauchi R, Inoue A, Tonomura H, Hiraoka N, Inoue H, Tsuchida S,

- Mazda O, and Kubo T: Combined microwave irradiation and intraarticular glutamine administration-induced hsp70 expression therapy prevents cartilage degradation in a rat osteoarthritis model. *J Orthop Res.* 2012, 30: 401–407.
- 13) Kim H, Suzuki T, Saito K, Kim M, Kojima N, Ishizaki T, Yamashiro Y, Hosoi E, and Yoshida H: Effectiveness of exercise with or without thermal therapy for community-dwelling elderly Japanese women with non-specific knee pain: a randomized controlled trial. *Arch Gerontol Geriatr.* 2013, 57: 352–359.
 - 14) Ito A, Aoyama T, Yamaguchi S, Zhang X, Akiyama H, and Kuroki H: Low-intensity pulsed ultrasound inhibits messenger RNA expression of matrix metalloproteinase-13 induced by interleukin-1 β in chondrocytes in an intensity-dependent manner. *Ultrasound Med Biol.* 2012, 38: 1726–1733.
 - 15) Oosterveld FG, and Rasker JJ: Treating arthritis with locally applied heat or cold. *Semin Arthritis Rheum.* 1994, 24: 82–90.
 - 16) Sanchez-Inchausti G, Vaquero-Martin J, and Vidal-Fernandez C: Effect of arthroscopy and continuous cryotherapy on the intra-articular temperature of the knee. *Arthroscopy.* 2005, 21: 552–556.
 - 17) Dewey WC, Hopwood LE, Sapareto SA, and Gerweck LE: Cellular responses to combinations of hyperthermia and radiation. *Radiology.* 1977, 123: 463–474.
 - 18) Wheatley DN, Kerr C, and Gregory DW: Heat-induced damage to HeLa-S3 cells: correlation of viability, permeability, osmosensitivity, phase-contrast light-, scanning electron- and transmission electron-microscopical findings. *Int J Hyperthermia.* 1989, 5: 145–162.
 - 19) Imada S, Yamamoto M, Tanaka K, Seiwa C, Watanabe K, Kamei Y, Kozuma S, Taketani Y, and Asou H: Hypothermia-induced increase of oligodendrocyte precursor cells: possible involvement of plasmalemmal voltage-dependent anion channel 1. *J Neurosci Res.* 2010, 88: 3457–3466.
 - 20) Al-Fageeh MB, and Smales CM: Control and regulation of the cellular responses to cold shock: the responses in yeast and mammalian systems. *Biochem J.* 2006, 397: 247–259.
 - 21) Read RA, Fox MH, and Bedford JS: The cell cycle dependence of thermotolerance. I. CHO cells heated at 42 degrees C. *Radiat Res.* 1983, 93: 93–106.
 - 22) Mackey MA, and Dewey WC: Cell cycle progression during chronic hyperthermia in S phase CHO cells. *Int J Hyperthermia.* 1989, 5: 405–415.
 - 23) Dewhirst MW, Viglianti BL, Lora-Michiels M, Hanson M, and Hoopes PJ: Basic principles of thermal dosimetry and thermal thresholds for tissue damage from hyperthermia. *Int J Hyperthermia.* 2003, 19: 267–294.
 - 24) Yarmolenko PS, Moon EJ, Landon C, Manzoor A, Hochman DW, Viglianti BL, and Dewhirst MW: Thresholds for thermal damage to normal tissues: an update. *Int J Hyperthermia.* 2011, 27: 320–343.
 - 25) Hojo T, Fujioka M, Otsuka G, Inoue S, Kim U, and Kubo T: Effect of heat stimulation on viability and proteoglycan metabolism of cultured chondrocytes: Preliminary report. *J Orthop Sci.* 2003, 8: 396–399.
 - 26) Kafienah W, Al-Fayez F, Hollander AP, and Barker MD: Inhibition of cartilage degradation: a combined tissue engineering and gene therapy approach. *Arthritis Rheum.* 2003, 48: 709–718.
 - 27) Park CH, Lee MJ, Ahn J, Kim S, Kim HH, Kim KH, Eun HC, and Chung JH: Heat shock-induced matrix metalloproteinase (MMP)-1 and MMP-3 are mediated through ERK and JNK activation and via an autocrine interleukin-6 loop. *J Invest Dermatol.* 2004, 123: 1012–1019.
 - 28) Li WH, Lee YM, Kim JY, Kang S, Kim S, Kim KH, Park CH, and Chung JH: Transient receptor potential vanilloid-1 mediates heat-shock-induced matrix metalloproteinase-1 expression in human epidermal keratinocytes. *J Invest Dermatol.* 2007, 127: 2328–2335.
 - 29) Morimoto RI, Kroeger PE, and Cotto JJ: The transcriptional regulation of heat shock genes: a plethora of heat shock factors and regulatory conditions. *EXS.* 1996, 77: 139–163.
 - 30) Beckham JT, Wilbanks GJ, Mackanos MA, Takahashi K, Contag CH, Takahashi T, and Jansen ED: Role of hsp70 in cellular thermotolerance. *Lasers Surg Med.* 2008, 40: 704–715.
 - 31) Terauchi R, Takahashi KA, Arai Y, Ikeda T, Ohashi S, Imanishi J, Mazda O, and Kubo T: Hsp70 prevents nitric oxide-induced apoptosis in articular chondrocytes. *Arthritis Rheum.* 2003, 48: 1562–1568.
 - 32) Mcardle A, Dillmann WH, Mestrl R, Faulkner JA, and Jackson MJ: Overexpression of hsp70 in mouse skeletal muscle protects against muscle damage and age-related muscle dysfunction. *FASEB J.* 2004, 18: 355–357.
 - 33) Okita M, Nakai K, Kataoka H, Toyoda N, Nakano J, Origuchi T, and Yoshimura T: Effects of heat stress on prevention of disuse muscle atrophy in rat soleus muscle. *Rigakuryohogaku.* 2004, 31: 63–69. (Japanese)
 - 34) Fujino H, Kohzaki H, Takeda I, Tasaki H, Kondo H, Ishida T, and Kajiya F: Protective effects of a pre-conditioning exercise on soleus muscle atrophy by tail suspension. *Rigakuryohogaku.* 2005, 32: 400–405. (Japanese)
 - 35) Watanabe Y, Yoshikawa S, Kataoka H, Kataoka N, Sakamoto J, Nakano J, and Okita M: Effects of differences in heating methods on inhibiting the progression of disuse muscle atrophy in rat soleus muscle. –comparison of a warm water bath and electrical heating plate. *Rigakuryohogaku.* 2006, 33: 355–362. (Japanese)
 - 36) Li L, Zhang T, Zhou L, Xing G, Chen Y, and Xin Y: Schisan-drin B attenuates acetaminophen-induced hepatic injury through overexpression of heat shock protein 27 and 70 in mice. *J Gastroenterol Hepatol.* 2013, doi: 10.1111/jgh.12425. [Epub ahead of print]
 - 37) Chen Y, Ross BM, and Currie RW: Heat shock treatment protects against angiotensin II-induced hypertension and inflammation in aorta. *Cell Stress Chaperones.* 2004, 9: 99–107.
 - 38) Flour MP, Ronot X, Vincent F, Benoit B, and Adolphe M: Differential temperature sensitivity of cultured cells from cartilaginous or bone origin. *Biol Cell.* 1992, 75: 83–87.
 - 39) Shui CX, and Scutt A: Mild heat shock induces proliferation, alkaline phosphatase activity, and mineralization in human bone marrow stromal cells and MG-63 cells in vitro. *J Bone Miner Res.* 2001, 16: 731–741.
 - 40) Morrissey JJ, Higashikubo R, Goswami PC, and Dixon P: Mild hyperthermia as a potential mechanism to locally enhance cell growth kinetics. *J Drug Target.* 2009, 17: 719–723.