文 献 紹 介

文献委員会

21220

The Radiologic Study of the Normal Spine: Cervical, Thoracic, Lumbar, and Sacral.

Christenson, P.C. The Radiologic Clinics of North America. 15(2), 133, (1977).

21262

Number of Projections in Mammography: Influence on Detection of Breast Disease

Ingvar Andersson et al, Amer. J. Roentgenol. 130 (2), 349~351, (1978).

Summary

A total of 491 cases of breast cancer were reviewed to evaluate the diagnostic importance of the three standard projections; craniocaudal, mediolateral, and oblique. With only one projection, 90% of the malignancies would have been detected; with two and three projections the percentages rose to 94% and 95%, respectively. Nonvisualization was due to either juxtathoracic position of the tumor or superimposition of dense tissues. A relation was found between the position of a juxtathoracic tumor and the projection in which it was visualized. The mean diameter of tumors not visible due to superimposition of dense tissues was smaller than that of the whole material. In breast screening programs, use of two projections, the oblique and craniocaudal, is recommended.

21322, 24420

A Simple Catheter Curve for Cerebral Angiography

Stephen B. Fierstien, Radiology. 126 (2), 524~525, (1978)

Summary

A simple catheter curve for use in cerebral angiography is discussed. A shorter length of guide wire is needed before the catheter tip will enter the vessel.

21324, 21319

Prostaglandin E_1 (PGE_1) in Angiography of Tumors of the Extremities.

Kjell Jonsson et al, Amer. J. Roentgenol. 130(1) 7~11, (1978).

Summary

Prostaglandin E_1 (PGE_1) is a potent vasodilator.

The value of this agent as an adjuvant in angiography of tumors of the extremities was evaluated in 10 patients. Except in one case involving a hemangioma, the diagnostic information concerning tumor vascularity and delineation of the extent of disease decreased after the use of PGE_1 . The visualization of small normal muscular arteries was improved. Therefore, the effect of PGE_1 may be to preferentially increase the blood flow in normal tissues relative to the tumor. PGE_1 seems to be of little value in enhancing the angiographic demonstration of neoplasms of the extremity.

21324

Simplified Outpatient Lower Extremity Venography.

Helene Pavlov et al, Radiology. 126(2), 525~527, (1978).

Summary

Improved lower extremity venography has been achieved by a distally directed superficial foot venipuncture. Supine position and short procedure time afford excellent patient acceptance. Dilute contrast is believed responsible for eliminating venospasm and postvenography phlebitis; the low sodium concentration permits safe evaluation of cardiac patients. Sequential imaging (and the use of anteroposterior and lateral projections) eliminates false positive and negative interpretations. Skip areas are avoided by the use of 90-cm (36-inch) cassettes.

21390, 21243

An Easy Method to Demonstrate the Cruciate Ligaments by Double Contrast Arthrography.

Helene Pavlov et al, Radiology, 126(3), 817~818, (1978).

Summary

A reproducible technique which does not require special equipment for demonstrating the cruciate ligaments by double contrast arthrography is described. The positive contrast medium coats the synovial surfaces of the ligaments, *i.e.*, the anterior aspect of the anterior cruciate ligament and the posterior aspect of the posterior cruciate ligaments. The ligaments appear