EXPANDED ABSTRACT

Monitoring of IP₃ dynamics during the mechanical stimulation-induced intra- and intercellular Ca²⁺ waves in HSY human parotid cell line

Akihiro Nezu, Akihiko Tanimura, Takao Morita, and Yosuke Tojyo

Department of Pharmacology, School of Dentistry, Health Sciences University of Hokkaido, Hokkaido, Japan

Keywords : mechanical stimulation, Ca²⁺ dynamics, inositol 1,4,5-trisphosphate (IP₃), ATP

J. Med. Invest. 56 Suppl. : 388-390, December, 2009

INTRODUCTION

It is well known that mechanical stimulation elicits Ca²⁺ responses in various cell types. Touch stimulation on a single cell induces an increase in the cytosolic Ca²⁺ concentration ([Ca²⁺]_i), which subsequently propagates to neighboring cells as an intercellular Ca²⁺ wave. In addition to Ca²⁺ entry through mechanosensitive channels, this Ca²⁺ response is thought to be induced, at least in part, by inositol 1,4,5-trisphosphate (IP₃)-dependent Ca²⁺ release from intracellular Ca²⁺ stores (1-3). However, this hypothesis has not been tested directly due to a lack of the quantitative method to measure the cytosolic concentration of IP₃ ([IP₃]_i) in a single living cell.

Recently, we developed a FRET-based IP₃ biosensor, LIBRAvIII, and a method to measure [IP₃]_i quantitatively in the single living cell (4). In order to examine the involvement of IP₃ in mechanical stimulation-induced Ca²⁺ responses, we monitored IP₃ and Ca²⁺ responses simultaneously in HSY-EA1 cells, a human parotid cell line, using LIBRAvIII and the Ca²⁺ indicator fura-2.

METHODS

HSY-EA1 cells were cultured for 1 week in a

recording chamber. LIBRAvIII and LIBRAvN, an IP₃ insensitive variant of LIBRAvIII (4), were transiently expressed using Lipofectamine 2000. These HSY-EA1 cells were incubated with 2 μM fura-2/AM.

HSY-EA1 cells were stimulated by poking (0.1 sec) the cell membrane surface using a glass micropipette equipped with a micromanipulator (Fig. 1a). Simultaneous monitoring of LIBRAvIII, LIBRAvN and fura-2 was performed with sequential excitation at 380 nm (for fura-2) and 430 nm (for LIBRAvIII and LIBRAvN). Dual emission fluorescence was acquired on an AQUACOSMOS/ASHURA imaging system, and fluorescence signals were recorded by a cooled 3CCD color camera [CFP signal was detected using the C channel (420-500 nm), and fura-2 or Venus signals using the Y channel (500-565 nm)]. The [IP₃]_i in individual cells was estimated as described (4).

Fig. 1 Measurement of mechanical stimulation-induced IP₃ and Ca²⁺ responses in HSY-EA1 cells.

Illustration of the system used for mechanical stimulation and measurement of LIBRAvIII and fura-2 fluorescence. Cells were poked by a glass micropipette as shown in (a).

Received for publication October 19, 2009 ; accepted October 26, 2009.

Address correspondence and reprint requests to Akihiro Nezu, Department of Pharmacology, School of Dentistry, Health Sciences University of Hokkaido, 061-0293, Japan and Fax : +81-133-23-1399.

RESULTS AND DISCUSSION

When a HSY-EA1 cell was poked using a glass micropipette, the Ca^{2+} response initiated at the stimulated region and then spread throughout the cell. Subsequently, the Ca^{2+} response propagated to neighboring cells as an intercellular Ca^{2+} wave. In order to examine the mechanisms by which mechanical stimulation induced Ca^{2+} responses, the cells were stimulated in various experimental conditions. Mechanical stimulation-induced Ca^{2+} responses were observed in both stimulated and neighboring cells in the absence of extracellular Ca^{2+} . Pretreatment of cells with 10 μM U-73122, a phospholipase C (PLC) inhibitor, completely attenuated the Ca^{2+} response in neighboring cells. These results indicate that Ca^{2+} responses in neighboring cells are primarily mediated by the Ca^{2+} release from intracellular stores. In contrast, U-73122 treatment reduced Ca^{2+} responses in mechanically stimulated cells by $\sim 50\%$ in the presence of extracellular Ca^{2+} , but the responses were attenuated completely in its absence. These results indicate that the Ca^{2+} response in mechanically stimulated cells results from Ca^{2+} entry and PLC-mediated Ca^{2+} release.

It has been reported that ATP is involved in the propagation of intercellular Ca^{2+} waves after mechanical stimulation (1, 5, 6). Pretreatment of cells with 100 μM suramin, a purinergic receptor blocker, inhibited Ca^{2+} responses in neighboring cells almost completely. Although gap junctions have been shown to contribute to intercellular Ca^{2+} waves in some cell types (7, 8), the gap junction inhibitors 1-octanol and 16-DSA had no effect on Ca^{2+} responses in HSY-EA1 cells. Taken together, these data indicate that mechanical stimulation-induced intercellular Ca^{2+} waves in HSY-EA1 cells primarily result from ATP released by stimulated cells (Fig. 2).

In contrast to neighboring cells, suramin reduced Ca^{2+} responses in mechanically stimulated cells by only $\sim 50\%$. These suramin resistant Ca^{2+} responses were observed even in the absence of extracellular Ca^{2+} . One might assume that this Ca^{2+} release may be attributable to the disruption of intracellular Ca^{2+} stores by mechanical stimulation. However, Ca^{2+} release in mechanically stimulated cells was completely inhibited by U-73122. These results suggest that Ca^{2+} release in mechanically stimulated cells is induced, at least in part, by an ATP-independent mechanism (Fig. 2).

Our studies indicate that mechanical stimulation-induced Ca^{2+} release is mediated by ATP-dependent

Fig. 2 Mechanical stimulation induces IP_3 generation via ATP-dependent and ATP-independent PLC activation in stimulated and neighboring cells.

ER: endoplasmic reticulum. IP_3R : IP_3 receptor. MS cell: mechanically stimulated cell. NB cell: neighboring cell.

and/or ATP-independent activation of PLC. Therefore, IP_3 generation would be expected to contribute to this process. In order to examine this idea directly, we monitored changes in $[\text{IP}_3]_i$ during mechanical stimulation-induced Ca^{2+} responses in HSY-EA1 cells using LIBRAvIII and fura-2. Mechanical stimulation induced a transient increase in the emission ratio of LIBRAvIII in both stimulated and neighboring cells, and these responses correlated with the Ca^{2+} response. In the presence of 10 μM U-73122, the mechanical stimulation-induced increases in the emission ratio of LIBRAvIII in both stimulated and neighboring cells were completely blocked. In addition, mechanical stimulation did not change the emission ratio of LIBRAvN, an IP_3 insensitive variant of LIBRAvIII, even though large Ca^{2+} responses were observed. These experiments indicate that mechanical stimulation-induced changes in the emission ratio of LIBRAvIII indeed reflect changes in $[\text{IP}_3]_i$.

We then examined the effect of suramin on mechanical stimulation-induced increases in $[\text{IP}_3]_i$ in HSY-EA1 cells. In the presence of 100 μM suramin, mechanical stimulation-induced increases in $[\text{IP}_3]_i$ in neighboring cells were completely attenuated. In contrast, suramin had only a small effect on the increase in $[\text{IP}_3]_i$ in mechanically stimulated cells. These results indicate that mechanical stimulation-induced generation of IP_3 in stimulated cells involves an ATP-independent PLC activation pathway. In neighboring cells, however, IP_3 generation and Ca^{2+} responses were mediated by ATP released from stimulated cells (Fig. 2). Given that $[\text{IP}_3]_i$ was modulated by released ATP, we postulated that ATP may have autocrine effects on mechanical stimulation-induced IP_3 generation. Nevertheless, our data

indicate that the contribution of ATP-mediated pathways in mechanically stimulated cells is very small.

CONCLUSION

In this study, we show that IP₃-induced Ca²⁺ release contributes to mechanical stimulation-induced Ca²⁺ responses in both stimulated and neighboring cells. Our results suggest that Ca²⁺ release in mechanically stimulated cells is primarily induced by an ATP-independent pathway, whereas Ca²⁺ responses in neighboring cells are mediated by ATP released from stimulated cells. Furthermore, we demonstrate directly that mechanical stimulation induces an increase in [IP₃]_i in both stimulated and neighboring cells during Ca²⁺ responses. The mechanism of IP₃ generation in mechanically stimulated cells remains unknown, and further experiments will be required to elucidate these pathways.

REFERENCES

1. Sanderson MJ, Charles AC, Dirksen ER : Mechanical stimulation and intercellular communication increases intracellular Ca²⁺ in epithelial cells. *Cell Regul* 1 : 585-596, 1990
2. Boitano S, Dirksen ER, Sanderson MJ : Intercellular propagation of calcium waves mediated by inositol trisphosphate. *Science* 258 : 292-295, 1992
3. Moerenhout M, Himpens B, Vereecke J : Intercellular communication upon mechanical stimulation of CPAE- endothelial cells is mediated by nucleotides. *Cell Calcium* 29 : 125-136, 2001
4. Tanimura A, Morita T, Nezu A, Shitara A, Hashimoto N, Tojyo Y : Use of fluorescence resonance energy transfer-based biosensors for the quantitative analysis of inositol 1,4,5-trisphosphate dynamics in calcium oscillations. *J Biol Chem* 284 : 8910-8917, 2009
5. Osipchuk Y, Cahalan M : Cell-to-cell spread of calcium signals mediated by ATP receptors in mast cells. *Nature* 359 : 241-244, 1992
6. Enomoto K, Furuya K, Yamagishi S, Oka T, Maeno T : The increase in the intracellular Ca²⁺ concentration induced by mechanical stimulation is propagated via release of pyrophosphorylated nucleotides in mammary epithelial cells. *Pflügers Arch* 427 : 533-542, 1994
7. Charles AC, Naus CC, Zhu D, Kidder GM, Dirksen ER, Sanderson MJ : Intercellular calcium signaling via gap junctions in glioma cells. *J Cell Biol* 118 : 195-201, 1992
8. Hansen M, Boitano S, Dirksen ER, Sanderson MJ : A role for phospholipase C activity but not ryanodine receptors in the initiation and propagation of intercellular calcium waves. *J Cell Sci* 108 : 2583-2590, 1995