CASE REPORT

Usefulness of central venous oxygen saturation monitoring during bidirectional Glenn shunt

Nami Kakuta¹, Shinji Kawahito¹, Naoji Mita¹, Noriko Kambe¹, Asuka Kasai¹,
Narutomo Wakamatsu¹, Toshiko Katayama¹, Tomohiro Soga¹, Fumihiko Tada²,
Takashi Kitaichi³, Tetsuya Kitagawa³, and Hiroshi Kitahata⁴

¹Department of Anesthesiology, Tokushima University Hospital, Tokushima, Japan, ²Department of Anesthesia, Shikoku Medical Center for Children and Adults, Kagawa, Japan, ³Department of Cardiovascular Surgery, Tokushima University Hospital, Tokushima, Japan, ⁴Department of Dental Anesthesiology, Tokushima University Hospital, Tokushima, Japan

Abstract : A PediaSat™ oximetry catheter (PediaSat : Edwards Lifesciences Co., Ltd., Irvine, CA, U.S.A.), which facilitates continuous measurement of central venous oxygen saturation (ScvO₂), may be useful for surgery for pediatric congenital heart disease. We used PediaSat during a bidirectional Glenn shunt. The patient was a 13-month-old boy. Under a diagnosis of left single ventricle (pulmonary atresia, right ventricular hypoplasia, atrial septal defect) and residual left aortic arch/left superior vena cava, a modified right Blalock-Taussig shunt was performed. Cyanosis deteriorated, so a bidirectional Glenn shunt was scheduled. After anesthesia induction, a 4.5 Fr double-lumen (8 cm) PediaSat was inserted through the right internal jugular vein for continuous ScvO₂ monitoring. Furthermore, the probe of a near-infrared, mixed blood oxygen saturation-measuring monitor was attached to the forehead for continuous monitoring of the regional brain tissue mixed blood oxygen saturation (rSO₂) (INVOS™ 5100C, Covidien ; Boulder, CO, U.S.A.). Blockage of the right pulmonary artery and right superior vena cava decreased the oxygen saturation, ScvO₂, and rSO₂, but increased the central venous pressure. Although changes in ScvO₂ were parallel to those in rSO₂, the former showed more marked changes. A combination of ScvO₂ and rSO₂ for monitoring during Glenn shunt may be safer. *J. Med. Invest.* 60 : 272-275, August, 2013

Keywords : bidirectional Glenn shunt, circulatory monitoring, central venous oxygen saturation, PediaSat oximetry catheter, near-infrared spectroscopy

INTRODUCTION

A PediaSat™ oximetry catheter (PediaSat : Edwards Lifesciences Co., Ltd., Irvine, CA, U.S.A.),

Received for publication May 13, 2013 ; accepted July 20, 2013.

Address correspondence and reprint requests to Nami Kakuta, M.D., Department of Anesthesiology, Tokushima University Hospital, 3-18-15, Kuramoto, Tokushima 770-8503, Japan and Fax : +81-88-633-7182.

which facilitates continuous measurement of the central venous oxygen saturation (ScvO₂), may be useful for surgery for pediatric congenital heart disease, which induces marked changes in intraoperative circulatory kinetics (1, 2). We used PediaSat during a bidirectional Glenn shunt, and confirmed the usefulness of continuous ScvO₂ measurement during surgery.

CASE REPORT

The patient was a 13-month-old boy. His height and body weight were 74 cm and 10 kg, respectively. Under a diagnosis of left single ventricle (pulmonary atresia, right ventricular hypoplasia, atrial septal defect) and residual left aortic arch/left superior vena cava, a modified right Blalock-Taussig (BT) shunt was performed 11 days after birth. Cyanosis deteriorated, and balloon dilatation was conducted to treat shunt anastomotic stenosis. However, there was no response. A bidirectional Glenn (hemi-Fontan) shunt was scheduled. Anesthesia was induced with midazolam, and muscle relaxation was achieved with rocuronium. Endotracheal intubation was performed using a tube measuring 4.5 mm in inner diameter (ID). Anesthesia was maintained with sevoflurane and fentanyl. After anesthesia induction, a 4.5 Fr double-lumen (8 cm) PediaSat was inserted through the right internal jugular vein for continuous ScvO₂ monitoring. Furthermore, the probe of a near-infrared, mixed blood oxygen saturation-measuring monitor was attached to the forehead for continuous monitoring of the regional brain tissue mixed blood oxygen saturation (rSO₂) (INVOS™ 5100C, Covidien ; Boulder, CO, U.S.A.).

For transesophageal echocardiography, a multi-plane for children (Philips, Inc. ; Andover, MA, U.S.A.) was used.

Cardiopulmonary bypass was not performed. Blockage of the right pulmonary artery and right superior vena cava decreased the oxygen saturation, ScvO₂, and rSO₂, but increased the central venous pressure (CVP). Immediately before blockage, SpO₂, CVP, rSO₂, and ScvO₂ were 81%, 16 mmHg, 66%, and 83%, respectively. However, after blockage, the values were 71%, 25 mmHg, 50%, and 50%, respectively. Although the changes in ScvO₂ paralleled those in rSO₂, the former showed more marked changes. The response of ScvO₂ was prompt, and there were no artifacts ; the value was stable. As both rSO₂ and ScvO₂ were maintained at 50%, surgery was continued. After the blockage was relieved, all parameters promptly returned to the pre-blockage values. Changes upon blockage of the left pulmonary artery and left superior vena cava were slight. Subsequently, respiratory/circulatory kinetics stabilized, and surgery was completed (Figure). The postoperative course was favorable. The patient recovered well in the intensive care unit without any neurological issues.

Figure : Anesthesia record

HR : Heart rate, SpO₂ : Percutaneous oxygen saturation, CoreT : Core temperature, NBP : Noninvasive blood pressure, ABP : Arterial blood pressure, etCO₂ : End-tidal carbon dioxide, rSO₂ : Continuous monitoring of the regional brain tissue mixed blood oxygen saturation, ScvO₂ : Continuous measurement of the central venous oxygen saturation,
First arrow : Blockage of the right pulmonary artery and superior vena cava
Second arrow : Blockage of the left pulmonary artery and superior vena cava

DISCUSSION

The bidirectional Glenn shunt is a commonly performed procedure in the treatment of univentricular physiology. It can be performed with or without cardiopulmonary bypass as long as pulmonary perfusion is maintained (3, 4). During the procedure, the superior vena cava (SVC) is clamped and the proximal pressure can rise as high as 40 to 55 mmHg. This can hamper cerebral perfusion. The duration of the procedure is variable, ranging from five to 30 minutes. During this period, reliable cerebral monitoring is essential.

The goal of continuous ScvO₂ monitoring is to improve the conditions of severe-status patients by maintaining an appropriate balance between oxygen supply and consumption (5). The use of a PreSep™ CV Oximetry Catheter in adults and a PediaSat™ Oximetry Catheter in children facilitates continuous monitoring of the central venous oxygen saturation (ScvO₂). ScvO₂ changes in parallel to mixed venous oxygen saturation (SvO₂) (6). As ScvO₂ depends on the cardiac output/hemoglobin level/arterial blood oxygen saturation involved in oxygen transport, as well as metabolism involved in oxygen consumption, continuous monitoring of the oxygen demand/supply balance is possible (7). A study performed a protocol involving goal-directed therapy (GDT) (8) to maintain ScvO₂ at 70% or more in high-risk patients, and reported that a decrease in the mortality rate and reduction in the admission period were achieved (9). Systemic management using ScvO₂ as an index was introduced to guidelines for septic shock treatment. Many studies have suggested the usefulness of ScvO₂ (10, 11). In children, the monitoring of changes in ScvO₂ also facilitates prompt management, decreasing the incidence of complications and the mortality rate in those with congenital heart disease (1, 2).

As an index of cerebral perfusion, noninvasive monitoring of regional cerebral oxygen saturation by the use of near-infrared spectroscopy (NIRS) cerebral oximetry (Invivo™) has been used. The usefulness of rSO₂ continuous monitoring during pediatric cardiac surgery has also been reported. It was reported that a close correlation was observed between cerebral NIRS and SVC oxygen saturation in neonates with congenital heart disease (12).

We performed the simultaneous monitoring of ScvO₂ and rSO₂ during a bidirectional Glenn shunt, during which blockage of the pulmonary artery/SVC may cause cerebral edema/ischemia, and confirmed

the usefulness of the two parameters. In particular, the changes in ScvO₂ were more marked than those in rSO₂. The maximum frequency of ScvO₂ renewal was once every two seconds. It was lower than that of rSO₂ monitoring (6 times per second), but a sensitive response was achieved. Although rSO₂, which is measured by attaching a probe to the forehead, also reflects information on venous blood to some extent, it facilitates brain-tissue monitoring, and is strongly influenced by the autoregulation capacity. As described above, the changes in ScvO₂ were more marked than those in rSO₂. This was possibly because the influence of the cerebral autoregulation capacity on ScvO₂ was less marked than that on rSO₂, since ScvO₂ reflects the upper body venous blood oxygen saturation.

In this case, ScvO₂ showed higher values during surgery than we expected (especially before blockage of the pulmonary artery and superior vena cava; ScvO₂ 83%). During a hyperdynamic state of septic shock, so-called warm shock, ScvO₂ is said possibly to show an abnormally high value by oxygen utilization disorder in the tissue or increase of shunts (11). However, a definitive conclusion cannot be drawn on this issue because we did not measure the real value of oxygen saturation in the superior vena cava by blood gas analysis; however, the patients might enter a hyperdynamic state by the effects of a high concentration of oxygen, general anesthesia, catecholamine, and so on. It is thought to be important to observe not the absolute value but the change over time of ScvO₂. In addition, continuous measurement of ScvO₂, in combination with other surrogates of organ perfusion (rSO₂, vital signs, lactate, etc.), can be used as a reliable monitor during bidirectional Glenn shunt.

CONCLUSION

We reported a patient in whom the usefulness of central venous oxygen saturation monitoring during bidirectional Glenn shunt was confirmed. Continuous ScvO₂ monitoring with a PediaSat™ Oximetry Catheter may be particularly significant for performing cardiac surgery in children in whom there are marked changes in hemodynamics. This procedure may facilitate the prediction of a sign of hemodynamic disturbance that cannot be detected using other monitoring methods. As ScvO₂ shows more marked changes than rSO₂, a combination of these two parameters for monitoring during Glenn shunt

may be safer. In particular, it may be very useful for bidirectional Glenn shunt in which cardiopulmonary bypass is not performed.

CONFLICT OF INTEREST

No conflicts of interest declared.

REFERENCES

1. Liakopoulos OJ, Ho JK, Yezbick A, Sanchez E, Naddell C, Buckberg GD, Crowley R, Mahajan A : An experimental and clinical evaluation of a novel central venous catheter with integrated oximetry for pediatric patients undergoing cardiac surgery. *Anesth Analg* 105 : 1598-1604, 2007
2. Spenceley N, Skippen P, Krahn G, Kissoon N : Continuous central venous saturation monitoring in pediatrics : a case report. *Pediatr Crit Care Med* 9 : e13-e16, 2008
3. Murthy KS, Coelho R, Naik SK, Punnoose A, Thomas W, Cherian KM : Novel techniques of bidirectional Glenn shunt without cardiopulmonary bypass. *Ann Thorac Surg* 67 : 1771-1774, 1999
4. Hussain ST, Bhan A, Sapra S, Juneja R, Das S, Sharma S : The bidirectional cavopulmonary (Glenn) shunt without cardiopulmonary bypass : is it a safe option? *Interact Cardiovasc Thorac Surg* 6 : 77-82, 2007
5. Collaborative study group on perioperative ScvO₂ monitoring : Multicentre study on peri- and postoperative central venous oxygen saturation in high-risk surgical patients. *Crit Care* 10 : R158, 2006
6. Reinhart K, Kuhn HJ, Hartog C, Bredle DL : Continuous central venous and pulmonary artery oxygen saturation monitoring in the critically ill. *Intensive Care Med* 30 : 1572-1578, 2004
7. Reinhart K, Bloos F : The value of venous oximetry. *Curr Opin Crit Care* 11 : 259-263, 2005
8. Mayer J, Boldt J, Mengistu AM, Röhm KD, Suttner S : Goal-directed intraoperative therapy based on autocalibrated arterial pressure waveform analysis reduces hospital stay in high-risk surgical patients : a randomized, controlled trial. *Crit Care* 14 : R18, 2010
9. Rivers E, Nguyen B, Havstad S, Ressler J, Muzzin A, Knoblich B, Peterson E, Tomlanovich M : Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med* 345 : 1368-1377, 2001
10. Perner A, Haase N, Wiis J, White JO, Delaney A : Central venous oxygen saturation for the diagnosis of low cardiac output in septic shock patients. *Acta Anaesthesiol Scand* 54 : 98-102, 2010
11. Pope JV, Jones AE, Gaieski DF, Arnold RC, Trzeciak S, Shapiro NI, Emergency Medicine Shock Research Network (EMShockNet) Investigators : Multicenter study of central venous oxygen saturation (ScvO₂) as a predictor of mortality in patients with sepsis. *Ann Emerg Med* 55 : 40-46, 2010
12. Ricci Z, Garisto C, Favia I, Schioderer U, Giorni C, Fragasso T, Picardo S : Cerebral NIRS as a marker of superior vena cava oxygen saturation in neonates with congenital heart disease. *Pediatr Anaesth* 20 : 1040-1045, 2010