EARLY ONLINE RELEASE

This is a PDF of a manuscript that has been peer-reviewed and accepted for publication. As the article has not yet been formatted, copy edited or proofread, the final published version may be different from the early online release.

This pre-publication manuscript may be downloaded, distributed and used under the provisions of the Creative Commons Attribution 4.0 International (CC BY 4.0) license. It may be cited using the DOI below.

The DOI for this manuscript is

DOI:10.2151/jmsj.2021-066

J-STAGE Advance published date: July 2nd, 2021

The final manuscript after publication will replace the preliminary version at the above DOI once it is available.

1
2 **On the semidiurnal variation in surface rainfall rate over the tropics**
3 **in a global cloud-resolving model simulation and satellite observations**
4

5 Toshiro INOUE¹

6 Atmosphere and Ocean Research Institute, The University of Tokyo, Kashiwa, Chiba, Japan
7

8 Kavirajan RAJENDRAN

9 Multi-Scale Modelling Programme, CSIR Fourth Paradigm Institute, Bangalore, India
10

11 Masaki SATOH

12 Atmosphere and Ocean Research Institute, The University of Tokyo, Kashiwa, Chiba, Japan
13

14 Hiroaki MIURA

15 Department of Earth and Planetary Science, The University of Tokyo, Tokyo, Japan
16
17
18
19
20
21

22 (Submitted to JMSJ, 13 March 2020

23 Revised 31 May 2021)
24
25
26
27
28
29
30
31
32
33
34

¹ *Corresponding author address:* Toshiro Inoue, Atmosphere and Ocean Research Institute, The University of Tokyo, Kashiwa, Chiba, Japan.

E-mail: toshiroy@aoori.u-tokyo.ac.jp

Abstract

The dual peak semidiurnal variation in surface rainfall rate over the tropics, simulated by a 3.5-km mesh Non-hydrostatic Icosahedral Atmospheric Model (NICAM) for 26–31 December 2006, is analyzed and compared with data from the 17-year winter precipitation climatology of Tropical Rainfall Measuring Mission (TRMM) TMI (TRMM Microwave Imager), Precipitation Radar (PR), and the same 6-day data of Global Satellite Mapping of Precipitation, as well as infrared data from geostationary satellites.

We focus on land areas including southern Africa and the Amazon. Over these land areas, the NICAM simulation captures the primary peak in the afternoon and the secondary peak in the early morning, at similar times to those captured by TRMM data. In the PR observation, the primary peak of rainfall is mainly due to convective rain, whereas the secondary peak is due to stratiform rain. In the NICAM simulation, if a simple method is used for classification of convective/stratiform rain, convective rain is dominant all day long and the rainfall rate is generally higher than in the PR observation. However, an analysis of deep convection (DC) areas indicates consistency between the observation and NICAM; the primary peak of rainfall rate occurs at the mature stage of the number of DC areas, while the secondary peak occurs when the mean size of DC areas is almost at its highest point. In the NICAM simulation, however, the relative magnitudes of the two peaks are not represented well, and the contribution of the stratiform rain is underestimated.

54 The present study indicates that a high-resolution global nonhydrostatic model like NICAM
55 has the potential to overcome the limitations of coarse-resolution general circulation models by
56 reproducing semi-diurnal variation of deep convection, though there is room for improvement.

57

58 **Keywords** semidiurnal variation, surface rainfall, cloud resolving model, Tropical Rainfall
59 Measuring Mission, deep convection

60

61

62 **1. Introduction**

63 Rainfall over the tropics plays an important role in the transport of heat and moisture in the
64 atmosphere. Latent heat release by deep convection (DC) over the tropics is the major energy source
65 for the general circulation of the atmosphere. There has been a large number of studies on the diurnal
66 variation in rainfall rate (e.g., Kraus 1963; Andersson 1970; Wallace 1975; Gray and Jacobson 1977;
67 Mapes and Houze 1993; Janowiak et al. 1994; Sui et al. 1997; Dai 2001; Takayabu 2002; Hirose and
68 Nakamura 2005; Yamamoto et al. 2008; Schroder et al. 2009). These studies used a variety of
69 observations, ranging from regional rainfall data in a limited area and time to global satellite datasets.
70 Mechanisms of the diurnal variation in rainfall rate are summarized and reviewed in Yang and Smith
71 (2006). For the most part, however, they focused on the highest peak of rainfall rate in a day and a

72 few studies noticed on semidiurnal variation in rainfall rate.

73 Nitta and Sekine (1994) proposed that there is semidiurnal variation in convective activity over the
74 Intertropical Convergence Zone (ITCZ) and the South Pacific Convergence Zone (SPCZ) based on
75 the Fourier analysis of infrared data observed by the Geostationary Meteorological Satellite (GMS)
76 of Japan. Yang and Smith (2008) showed the importance of the study of convective/stratiform rain
77 and noted the secondary peak of rainfall with seasonal variation using the global observations of
78 rainfall by the Tropical Rainfall Measuring Mission (TRMM). Kikuchi and Wang (2008) studied the
79 daily variation in rainfall rate using 8 years of TRMM data. Their empirical orthogonal function
80 (EOF) analysis also indicated semidiurnal variation, although it was marginal compared to the diurnal
81 component of the EOFs. Furthermore, Yang et al. (2008) showed the persistent nature of the globally
82 prevalent semidiurnal variation in rainfall using 8 years of TRMM data.

83 Previously, Brier and Simpson (1969) showed the existence of semidiurnal variation in cloudiness
84 and rainfall over the tropical western Pacific and linked with the solar atmospheric tide. Lindzen and
85 Chapman (1969) explained that the semidiurnal mode is caused mainly by heating of stratospheric
86 ozone. Lindzen (1978) studied semidiurnal atmospheric tide using rainfall data at several small
87 tropical islands.

88 Diagnostic analysis of the simulation of diurnal variation in rainfall rate is important for the
89 verification and improvement of numerical models. General circulation models (GCMs) or global
90 climate models generally do not represent diurnal variations very well (Lin et al. 2000; Covey et al.

2016; Yuan et al. 2013). Improper representation of the diurnal cycle of convection in GCMs affects surface water balance and shortwave cloud forcing (Del Genio and Wu, 2010). Woolnough et al. (2004) studied the diurnal and semidiurnal variations of precipitation using an aqua-planet GCM, which is beneficial for isolating these variations from other fluctuations in rainfall. Dirmeyer et al. (2011) studied the diurnal variation in rainfall rate using GCMs. One reason that GCMs have difficulty reproducing diurnal variations is related to the representation of cumulus parameterization (e.g., Yang and Slingo, 2001; Takayabu and Kimoto, 2008). One way to overcome this cumulus parameterization problem is to use a finer mesh non-hydrostatic model with explicit cloud processes. Del Genio and Wu (2010) used a cloud resolving model to improve the simulation of the diurnal mode in terms of the entrainment rate by focusing on the representation of the transition of continental convection from shallow to deep.

It is expected that high-resolution non-hydrostatic models over the global domain capture the diurnal cycle of the rainfall rate well compared to GCMs. In fact, Sato et al. (2009) showed that the phase of the diurnal cycle of rainfall rate was well simulated by the Nonhydrostatic Icosahedral Atmospheric Model (NICAM; Tomita and Satoh 2004; Satoh et al. 2008a; Satoh et al. 2014), although NICAM overestimates mean rainfall and the magnitude of the diurnal variation (Dirmeyer et al. 2011). Various studies of the features of diurnal variation in rainfall rate using NICAM simulations have considered resolution dependency of horizontal grid size in the non-hydrostatic regime less than about 5 km (e.g., Sato et al. 2009; Noda et al. 2012; Yashiro et al. 2016). Such high-resolution global

110 atmospheric models are becoming more popular (Sato et al. 2019). The inter-comparison study of
111 the high-resolution global atmospheric model, called DYAMOND (The DYnamics of the
112 Atmospheric general circulation Modeled On Non-hydrostatic Domains), used global models with
113 mesh size less than 5 km. Almost all these models were found to capture diurnal variation very well
114 (Stevens et al. 2019).

115 Better representation of the semidiurnal variation in rainfall rate is also important for GCMs.
116 Tomita et al. (2005) showed that the semidiurnal variation in rainfall rate is accurately represented in
117 the NICAM aqua-planet experiment. Yasunaga et al. (2013) proposed a squeezing mechanism to
118 account for the afternoon peak in the NICAM aqua-planet experiment; that is, condensation of water
119 vapor is enhanced and more water vapor is squeezed within a cloud due to reduced temperature.
120 However, the semidiurnal variation in more realistic representations with a land and sea contrast has
121 not yet been analyzed and compared with observations.

122 Miura et al. (2007) simulated the realistic structure of a Madden-Julian Oscillation (MJO) event
123 that occurred in December 2006 with 3.5-km and 7-km mesh versions of NICAM. Using this dataset,
124 Sato et al. (2009) analyzed the diurnal cycle of precipitation and found that the phase and amplitude
125 of the diurnal cycle of precipitation over land has dependency on the horizontal resolution. The 3.5-
126 km mesh simulation produced the peak time and amplitude that are closer to those in the TRMM
127 Precipitation Radar (PR) observations. In the present study, we analyze data from the NICAM 3.5-
128 km mesh hindcast experiment performed by Miura et al. (2007). Inoue et al. (2012) analyzed the

129 semidiurnal variation in surface rainfall rate in the NICAM 3.5-km mesh simulation by Miura et al.
130 (2007) and compared it with satellite observations. Here, as a follow-up study of Inoue et al. (2012),
131 we extend the diagnostic study to analyze the characteristics of the semidiurnal variation in rainfall
132 rate simulated by the NICAM in detail and compare them with TRMM and infrared data from a
133 geostationary satellite.

134 The structure of the paper is as follows. The model, experimental design and observation datasets
135 are presented in Section 2. The overall characteristics of the global rainfall distribution and the
136 semidiurnal variation in rainfall rate are analyzed in Section 3. Detailed analysis of the semidiurnal
137 variation in rainfall rate is provided separately for different areas: first we focus on the land areas
138 including southern Africa and the Amazon, and then compare them with the eastern Pacific Ocean
139 areas and the maritime continent areas in Supplement. In Section 4, we present our concludes.

140

141 **2. Model, Simulation and Datasets**

142 **2.1 NICAM**

143 NICAM is configured to run with an explicit cloud microphysics scheme using a mesh size of a
144 few kilometers without using a cumulus parameterization scheme. The cloud microphysics scheme
145 by Grabowski (1998) is employed in the simulation data used for this study (Miura et al. 2007). It is
146 a simple three-category scheme in which airborne and precipitating hydrometeors are prognostic
147 variables. Cloud ice and snow are diagnosed by the temperature-dependent ratio between liquid and

ice phases from airborne and precipitating hydrometeors. The vertical grid size is 40 levels from 80.8 m at the lowest level to 38.1 km above ground level. Miura et al. (2007) integrated a 3.5-km mesh configuration of NICAM globally for a week [00 UTC on 26 December 2006 to 24 UTC on 31 December 2006, for the period of an active MJO event] with an initial condition of National Centers for Environmental Prediction (NCEP) global analysis data at 00UTC on 25 December 2006. The multi-scale organization of convection associated with MJO was analyzed by Nasuno et al. (2009).

In this study, the output data of the 3.5-km mesh NICAM for the one-week simulation is analyzed to understand the representation of semidiurnal variation. Although this is a short integration, the high resolution simulation provides a lot of samples from which to obtain statistical behaviors of cloud systems, and they can be compared directly with those of the real world as observed by satellite. We have constructed two-dimensional fields of mean surface rainfall rate and outgoing longwave radiation (OLR) on a $0.05^\circ \times 0.05^\circ$ latitude-longitude grid at 1.5-hour temporal resolution.

160

161 2.2 TRMM data

The TRMM 3G68 version 7 data from 1998–2014 are used in this study. The 3G68 data were constructed from the precipitation data observed by PR and TRMM Microwave Imager (TMI). Although spatial and temporal coverage of observations is limited along the satellite orbitals (PR: 220km swath; TMI :760 km swath), many samples have been collected over more than a decade. The TRMM satellite was launched on 28 November 1997, and observations stopped on 8 April 2015; the

167 Global Precipitation Measurement Mission (GPM) succeeded the TRMM observation (Hou et al.
168 2014). Because the period of the present simulation overlaps the TRMM observation, we used almost
169 the whole life-time observational data from TRMM to construct composite diurnal variations in area
170 mean rainfall rate. The precipitation data from TRMM was considered as the most reliable rainfall
171 measurements over the global domain and was succeeded by GPM (Seto et al. 2021). The TRMM
172 3G68 provides us with orbital rainfall information at $0.05^\circ \times 0.05^\circ$ longitude-latitude grids based on
173 the standard products of 2A25 (radar algorithm; Iguchi et al. 2000), 2A12 (passive microwave
174 algorithm; Kummerow et al. 2001) and 2B31 (combined algorithm; Haddad et al. 1997). We construct
175 the hourly area mean rainfall rate over tropics observed by PR and TMI (Rajendran and Nakazawa
176 2005). In this study, we use PR and TMI data as standard datasets of TRMM. Here, we use the 17-
177 year data for the total period 1998–2014 for PR and TMI.

178

179 2.3 Infrared data from the geostationary satellite

180 Infrared data from Spinning Enhanced Visible and Infrared Imager (SEVIRI) onboard Meteosat-
181 8 (Schmetz et al. 2002) and TRMM ancillary data (Janowiak et al. 2017) constructed from infrared
182 data from all geostationary satellites in 26–31 December 2006 are also gridded onto a $0.05^\circ \times 0.05^\circ$
183 latitude-longitude grid. Split window data of the SEVIRI is used to classify optically thicker and
184 thinner clouds (Inoue, 1985; Inoue 1987a) over southern Africa. Infrared data from TRMM ancillary
185 data are used for the analysis of DC over Amazon.

186 We also constructed a cumulative frequency histogram for both 11 μm brightness temperature (TB),
187 and NICAM-simulated OLR to find the corresponding OLR value for specific TB. The TBs of 213,
188 235, and 253 K are conventionally used to define DC. OLR values corresponding to the above TBs
189 in NICAM are determined from the accumulated frequency of TB with that of NICAM simulated
190 OLR (Inoue et al. 2008), although the values are slightly different region by region.

191

192 2.4 GSMaP data

193 The Global Satellite Mapping of Precipitation (GSMaP) is a precipitation product that uses
194 combined data from passive microwave (PMW) sensors in low Earth orbit and infrared radiometers
195 in geostationary Earth orbit (Kubota et al. 2020). The GSMaP dataset is based on a Kalman filter
196 model that refines the precipitation rate by using the cloud moving vector derived from two successive
197 infrared images (Ushio et al. 2009).

198 The GSMaP was developed by the Japan Aerospace Exploration Agency (JAXA) for the GPM
199 mission as the Japanese GPM standard product. The horizontal resolution is a $0.1^\circ \times 0.1^\circ$ latitude-
200 longitude grid, and the temporal resolution is 1 hour. The GSMaP_Gauge is a product adjusted with
201 the National Oceanic and Atmospheric Administration/Climate Prediction Center (NOAA/CPC)
202 unified gauge-based analysis of global daily precipitation (Mega et al. 2019). The algorithm uses an
203 optical estimation scheme, in which the solution is calculated by maximizing the probability density

204 function defined in the system model. Here, we use the reanalysis version of the GSMaP_Gauge
205 (GSMaP_Gauge_reanal) hourly data in 26-31 December 2006.

206

207 **3. Background Characteristics**

208 **3.1 Mean rainfall rate over global tropics**

209 Figure 1 shows the spatial distribution of mean rainfall rates from PR (a) and TMI (b) observations
210 and mean rainfall rate estimated from the 3.5-km mesh NICAM simulation (c) for the period of 00
211 UTC on 26 December 2006 to 24 UTC on 31 December 2006. The rainfall rate is averaged over each
212 $0.5^\circ \times 0.5^\circ$ latitude-longitude grid. Both TMI and PR indicate slightly noisy spatial patterns due to
213 limited observation time and coverage area by the TRMM satellite (PR: 220 km swath; TMI: 760 km
214 swath). However, the major rainfall areas in the NICAM simulation over ITCZ, SPCZ, southern
215 Africa, the maritime continent, and the Amazon correspond well to the PR and TMI observations.
216 There are some differences over latitudes higher than 30°N (the TRMM observation is limited to
217 latitudes lower than 35°N/S), particularly over the eastern part of the USA and the Atlantic Ocean.
218 The rainfall rate simulated by NICAM is slightly higher than the rate observed over the northern parts
219 (Equator– 15°S) of South America (hereafter Eq. is used to denote Equator).

220

221 **3.2 Semidiurnal variation in surface rainfall rate from TRMM (TMI, PR) and NICAM**

222 Yang et al. (2008) first showed that the secondary peak is globally prevalent and widespread in
223 nature using 8 years of TRMM data. They studied the diurnal variation in rainfall rate at $20^\circ \times 30^\circ$,
224 $10^\circ \times 10^\circ$ and $5^\circ \times 5^\circ$ latitude-longitude grid arrays over the area covered by the TRMM observation.
225 They found that the semidiurnal mode appears in 78% of the boxes in the $20^\circ \times 30^\circ$, 69% in the 10°
226 $\times 10^\circ$, and 65% in the $5^\circ \times 5^\circ$ grid arrays.

227 We also construct the global view of daily variation in rainfall rate normalized with the maximum
228 rainfall rate at each grid, from 17 years of TMI (Fig.2a: green) and PR (Fig.2a: blue) data from TRMM
229 3G68 for the period covering December, January, and February (DJF). Each figure in Fig. 2 consists
230 of a $10^\circ \times 15^\circ$ latitude/longitude grid array over 20°N – 20°S (24×4 grid boxes). The abscissa of each
231 grid box indicates the local standard time (LST) from 0 to 23 hours. The ordinate of each grid box is
232 the daily mean rainfall rate normalized to the maximum rainfall rate with a range of 0–1 and we
233 applied a three-time-step running mean twice to smooth out noisy variations. Inspection of the grid
234 boxes shows that many have a semidiurnal variation (two local peaks a day) in rainfall rate, especially
235 over Africa, the maritime continent, and the Amazon. There are some grids where diurnal variation
236 differs between the PR and TMI datasets, probably due to the difference in observation coverage
237 between PR and TMI. There also are few grids that indicate more than two peaks. Nonetheless, we
238 have validated each box using harmonic analysis. Here, we assigned A1 as the amplitude of the
239 diurnal (one peak a day) mode and A2 as the amplitude of the semidiurnal (two peaks a day) mode.
240 Hereafter, we refer to these modes as “diurnal variation” and “semidiurnal variation”, respectively,

241 and refer to the variation in a day as “daily variation”. We computed the amplitude of A1 and A2
242 over all 96 boxes and found that the amplitude of A2 is comparable to that of A1; the amplitude of
243 A2 is larger than that of A1 for 26 (29) boxes and is larger than two-thirds of the amplitude of A1 for
244 42 (44) boxes for TMI (PR) observations in the tropics. Figure 2b shows the distribution of the ratio
245 of amplitudes ($A2/A1$) greater than 0.5 corresponding to the areas shown in Fig. 2a. Note that x -axis
246 (longitude) is 0° – 360° in Fig. 2b. We can see from Figs. 2a, b that it is likely for rainfall rate to peak
247 twice a day over the tropics in $10^{\circ} \times 15^{\circ}$ latitude/longitude boxes from 17 years of the TRMM data.

248 Figure 2c shows the global view of daily variation in rainfall rate normalized to the maximum value
249 in each grid from the NICAM simulation. Figure 2d shows the distribution of the amplitude ratio of
250 (semidiurnal variation)/(diurnal variation) derived from the harmonic analysis for NICAM. We can
251 see that NICAM simulates the semidiurnal variation in rainfall rate over Africa, the maritime
252 continent, and the Amazon. However, the semidiurnal variation is not reproduced by the NICAM
253 simulation over some basins such as the South Pacific and the Indian Ocean where TMI and PR
254 indicate a relatively smaller signature of semidiurnal variation. In the next section, we focus on the
255 semidiurnal variation in rainfall rate over southern Africa and the Amazon where the dual peaks are
256 clearly seen in the NICAM simulation as well as in the observations, and compare the areal
257 characteristics of the semidiurnal variations. The additional analyses on the eastern Pacific Ocean
258 area and the maritime continent area are also given in Supplement.

259

260 **4. Areal characteristics of semidiurnal variation in rainfall rate**

261 **4.1 Southern Africa**

262 As shown in Fig. 1, NICAM reproduces a realistic distribution of mean rainfall over southern
263 Africa and the adjacent coastal areas and the oceanic rain belt along the east coast of Africa by
264 comparing with the PR and TMI observations. To analyze in detail the features of rainfall over
265 southern Africa, we selected the area of (15°–28°E, Eq.–20°S) avoiding large lakes and higher
266 mountains to capture the overall characteristics of NICAM performance in simulating rainfall for
267 relatively smooth land area. The day-to-day variation in simulated mean surface rainfall rate
268 averaged over this region on 26–31 December 2006 in UTC is shown in Fig. 3 (NICAM in red and
269 GSMaP in purple). We can see the obvious differences between NICAM and GSMaP for
270 magnitudes on 26 and 27, mismatch of one peak on 29 and 31 December. However, as implied by
271 Fig. 2c, the two local peaks of rainfall rate per day in the NICAM simulation in a day are clearly
272 seen during this six-day period except on 28 December, which is consistent with the GSMaP data,
273 which also shows two local peaks on several days (26, 29, and 30 December).

274 Figure 4 shows the mean daily variation in rainfall rate by the NICAM simulation (red), and the
275 TMI (green), PR (blue), and GSMaP (purple) observations over the southern African area. Here the
276 NICAM data and the GSMaP data are mean values computed from one week during 26–31 December,
277 while the TMI and PR data are the mean evolution of the diurnal cycle averaged for the 17 years of
278 data. We can see two peaks at 4.5 LST and 15 LST in the NICAM simulation. Two local peaks of

279 rainfall rate from PR (TMI) are also seen at 3.5 (0.5) LST and 15.5 (15.5) LST, although the morning
280 peak is tiny. We note that rainfall rate in the NICAM simulation is generally higher than that in the
281 PR, TMI, and GSMaP observation, especially in the morning. The time of the afternoon peak of
282 rainfall rate simulated by NICAM is close to that of PR, TMI and GSMaP. The NICAM simulation
283 and the TRMM and GSMaP all indicate semidiurnal variations for this area.

284 The relative magnitude of the two peaks of the semidiurnal variation and their timings are similar
285 between NICAM and PR, although the rainfall rate simulated by NICAM is higher than the rate in
286 the PR observation at all times of the day. TMI rainfall rate is much higher than PR rainfall rate in the
287 afternoon, although their peak times coincide. The TMI rainfall rate is generally higher than the PR
288 rainfall rate during the afternoon over land and cold surfaces. Furuzawa and Nakamura (2005) and
289 Rajendran and Nakazawa (2005) suspected that the systematic difference in rainfall rate between PR
290 and TMI, especially over land, might be caused by the life stage of DC. They suggested that scattering
291 signals from anvil cloud associated with afternoon DC caused a higher rainfall rate for the estimations
292 of the TMI rainfall rate than the PR rainfall rate.

293 To further explore the contributions of DC to the semidiurnal cycle, we examine the detailed
294 features of DC which is closely related to rainfall over the tropics. Figure 5a shows the spatial
295 distribution of DC over southern Africa at 03, 06, 09, 12, 15, 18, 21 and 24 UTC on 26 December
296 2006, based on Meteosat-8 infrared data. Here, we use UTC instead of LST; the area is close to the
297 prime meridian (the time difference between UTC and LST is about 1.5 hour (that is, 00 UTC is about

1.5 LST). The DC is defined as a cloud grid colder than 253 K in TB. Further split-window ($11\ \mu\text{m}$ and $12\ \mu\text{m}$) data from SEVIRI is used to classify optically thicker or thinner parts within the DC (Inoue, 1987a). In Fig 5a, green (blue) represents an optically thinner (thicker) part within the DC, while purple represents cloud top colder than 213 K. By this color coding, we can see the inner structure within the DC with the edge of 253 K. Figure 5a exemplifies the diurnal variation in the spatial distribution of DC. The size of DC is largest at about 03 UTC (~ 4.5 LST), decreases towards 12 UTC (~ 13.5 LST), especially in the southern part of the figure ($12^\circ\text{--}27^\circ\text{S}$), and then expands rapidly at 18 UTC (~ 19.5 LST). The size of DC decreases again towards 24 UTC (~ 1.5 LST). The thinner part of the DC area (shown in green in Fig. 5a) is smallest at 12 UTC (~ 13.5 LST) and larger towards 18 UTC (~ 19.5 LST). Then this area of DC shrinks toward 24 UTC (1.5 LST) on this day.

Figure 5b illustrates simulated DC defined by OLR less than 210, 165 and $102\ \text{W m}^{-2}$, which correspond to 253, 235, and 213 K of TB in the geostationary satellite, respectively, using the relationship between the cumulative frequencies of TB and OLR in NICAM (Inoue et al. 2008). Note that the OLR in NICAM is not directly related to the optical thickness of the cloud; we simply used green, blue, and purple to show the DC with OLR less than 210, 165, and $102\ \text{W m}^{-2}$, respectively in Fig. 5b. The large DC area at 03 UTC (~ 4.5 LST) shrinks toward 09 UTC (~ 10.5 LST). At 12 UTC (~ 13.5 LST), a large number of DCs of small size (blue and green) are represented, whereas at 18 UTC, a DC of large area (blue) is dominant. Then the areas of DC shrink toward 24 UTC (~ 1.5 LST). Such evolution of size and number of DC areas is similar to the characteristics seen in TB (Fig. 5a).

317 The 3.5-km mesh NICAM successfully captures the aforementioned characteristics of diurnal
318 variation in DC over the southern African region on this day. The realistic representation of the
319 diurnal variation in DC is one of the advantages of global cloud-resolving models (Sato et al. 2019;
320 Stevens et al. 2019).

321 The coverage of cold clouds over an analysis area has frequently been used to infer rainfall from
322 infrared data (e.g., Arkin and Meisner 1987). Figure 6 shows the observed diurnal variation for the
323 anomaly of number of DC grids ($0.05^\circ \times 0.05^\circ$ in latitude-longitude) defined by infrared data over
324 the analysis area (15° - 28° E, Eq. -20° S) for 26–31 December 2006 in UTC. The DC grid is defined by
325 TB colder than 213, 235, and 253K. In addition, we define the 253K-Cb and 253K-Ci cloud grids to
326 represent optically thick and thin clouds, respectively; these are defined as a cloud with TB that is
327 colder than 253 K with a relatively small or large TB difference for the split window, respectively
328 (Inoue, 1987a, b). Since the number of cloud grid over the analysis area is different depending on the
329 threshold, an anomaly from the daily mean is used in Fig.6. From this diurnal variation in cold cloud
330 grid, however, the semidiurnal mode of rainfall rate (cloud coverage) is not seen, although there is a
331 slight secondary peak at 24 UTC (~ 1.5 LST) in 253 K-Cb. Here, we note the difference in variation
332 between 253 K cloud (green) and 253 K-Cb cloud (red). The diurnal variation in 253 K-Cb appears
333 to be similar to that of 235 K (blue) and 213 K (purple), while the 253 K (green) peak is broad at 16–
334 22 UTC and similar to that of 253 K-Ci. This suggests that 253 K DC includes optically thinner anvil
335 that prevails for a longer period as seen in 253K-Ci. The 213 K peak appears earlier, around 14.5

336 UTC than the 235 K peak (at 15.5 UTC), and has timing similar to that of the 253 K-Cb peak; this
337 also suggests that the vertical enhancement of DC comes earlier than anvil expansion over southern
338 Africa.

339 Next, we investigated the size and number of DC areas, which are defined by the connected areas
340 of a specific TB. We utilize this method because the total area of DC identified by using a simple TB
341 threshold does not clearly show the semidiurnal variation. The analysis method is explained in
342 Appendix. We define a DC area as comprising all connected adjacent grids that have a TB below 213
343 K. The size of each DC area is defined by the number of grids in that area. The number of DC areas
344 includes only those DC areas whose centers are located within the analysis area. The mean size of
345 DC area is the average size of all DC areas whose centers are located within the analysis area. Figure
346 7a shows the daily variation in the number and mean size of DC areas. The analysis target is the area
347 (15° – 28° E, Eq.– 20° S) during the period of 26–31 December 2006, and the Meteosat-8 observation
348 data is used. The mean size of the DC areas (green) clearly has two local peaks in the early morning
349 and late afternoon, while the number of DC areas (blue) has a peak in the afternoon and a small
350 secondary peak early morning. The primary peak of the number of DC areas appears at 15.5 LST,
351 which is almost the same time as the PR rainfall peak (13.5–15.5 LST) in Fig. 4. The morning peak
352 of the mean size of the DC area at 4.5 LST almost coincides with the secondary peak of the PR rainfall
353 at 3.5 LST in Fig. 4. The primary peak of the rainfall rate observed by PR over this area coincides
354 with the primary peak of the number of DC areas which are rapidly expanding in size. This is

355 consistent with the finding by Inoue et al. (2009) that the rainfall rate is high at the earlier stage of
356 DC, based on a comparison between the life stage defined by the split-window and PR rainfall data.
357 Using SEVIRI data, Schroder et al. (2009) also found that deep convective systems start with high
358 rates of change in infrared TB in the formative stage, then continue with the minimum infrared TB
359 around 15.5 LST. Thus, it is important to note that the areal peak of rainfall is caused by a large
360 number of developing stages of DC. Figure 7a also suggests that large cloud size at 18 LST does not
361 necessarily coincide with a high rainfall rate in the late afternoon. The mean size of DC area is
362 expected to be large in the early morning due to radiation cooling at the cloud top (Gray and Jacobson
363 1977). The early morning peak of rainfall rate in PR seems to occur almost at the same time as the
364 secondary peak of the mean size of DC area.

365 Figure 7b shows the daily variation in mean size and number of DC areas defined by an OLR less
366 than 102 W m^{-2} (correspond to 213 K from cumulative analysis) in the NICAM simulation. We can
367 see the semidiurnal mode for both mean size and number of DC area; similar feature appear in the
368 infrared data for observations, although the peak times are slightly different. The primary peak of the
369 size of DC area appears later than the primary peak of the number of DC areas. The primary peak of
370 the number of DC areas coincides with the time at which the DC area increases in size. The morning
371 peak of the rainfall rate in the NICAM simulation (red curve in Fig.4) occurs slightly later than the
372 peak of size and number of DC area defined by OLR.

373 To examine how much DC contributes to the semidiurnal variations, we analyzed the diurnal

374 variations in convective/stratiform rain in PR observations and the NICAM simulation. Takayabu
375 (2002) showed that the convective rain in PR, defined by a rain type classification (Awaka et al. 1997;
376 2007), has a distinct maximum in the afternoon (15–18 LST), while stratiform rain has a maximum
377 over land around midnight (24–03LST). Figure 7c shows the daily variation in convective (blue) and
378 stratiform (green) rainfall rates defined by the PR observation (solid line) and the OLR of the NICAM
379 simulation (dotted line) over southern Africa (15°–28°E, Eq.–20°S). Convective/stratiform rainfall
380 rates in PR observations are defined by an algorithm based on Awaka et al. (1997, 2007). The
381 convective rainfall rate in PR data has a primary peak at 15.5 LST and a weak secondary peak at 3.5
382 LST. By contrast, stratiform rainfall rate is higher than the convective rainfall rate during the morning.
383 In PR observations, the primary peak of stratiform rainfall rate in the early morning (3.5 LST)
384 coincides with the secondary peak of convective rainfall rate. The secondary peak of total rainfall rate
385 in PR observations corresponds to both morning peaks of stratiform and convective rainfall rates.

386 For numerical simulation data, there are several methods for classifying convective/stratiform rain
387 types (Lang et al. 2003). Satoh et al. (2008b) used the rainfall rate of 10 mm hr⁻¹ as a threshold to
388 define convective rain following Grabowski (2003), and compared the rainfall statistics between the
389 NICAM 3.5-km mesh aqua-planet simulation and PR observations. From an analysis of the
390 probability distribution function of rainfall rates, they showed that the probability of rainfall rates for
391 heavy rain in NICAM simulations is similar to that of PR data. For the classification of stratiform
392 rain types, we use a threshold of 5 mm hr⁻¹ following evidence from Yang and Nesbitt (2014), based

393 on 13 years of PR observations, that 93% (92%) of stratiform rain events over the land (ocean) area
394 has a rainfall rate less than 5 mm hr⁻¹.

395 We use the threshold of 10 mm hr⁻¹ (5mm hr⁻¹) as an index for classifying of the convective
396 (stratiform) rain. The diurnal variation in convective and stratiform rainfall rates for the NICAM
397 simulation over the southern Africa analysis area is shown in Fig.7c by blue and green dotted lines,
398 respectively. For the convective rain, both the NICAM simulation and PR observations show the two
399 peaks, with similar local time maxima, but different magnitudes in the morning peaks. For the
400 stratiform rain, the NICAM simulation shows no clear peaks but with a flat peak between 3 and 15
401 LST. On the contrary, the PR observations show the morning peak at 3 LST, with no afternoon peak
402 in the stratiform rain rate. Further study is required for this difference between NICAM and PR
403 observation in terms of convective/stratiform rain.

404 To further understand the rainfall characteristics of the semidiurnal variation, we next studied the
405 size of the rainfall area in the NICAM simulation, as we did for the DC area (Figs. 7a, b). The rainfall
406 area is defined as the connected grids with a rainfall rate greater than 1 mm hr⁻¹. As before, a
407 convective rain grid is defined as a grid in which the rainfall rate is greater than 10 mm hr⁻¹. Figure
408 7d shows the daily variation in the size of the rainfall area (green, right ordinate: number of grid) and
409 the mean percentage of convective rain area within all the rain area (blue, left ordinate). This shows
410 that the size of the rain area is high during the morning and that the mean percentage of convective
411 rain areas within each rain area is large during the afternoon. The percentage of rain area without a

convective rain grid is large during the morning (not shown). These results suggest that the rainfall in the morning is characterized by stratiform rain to some extent, whereas the rainfall in the afternoon is characterized by convective rain. Therefore, in some measures, it can be said that the NICAM simulation reproduces the daily variation in convective/stratiform rain.

In summary, the semidiurnal variation in rainfall is seen both in the observation and the NICAM simulation over southern Africa. From the PR observation, we found that the afternoon peak is due to convective rain and that the morning peak is due to stratiform rain. For the NICAM simulation, as seen in Fig.7c, the contribution of convective/stratiform rain over the analysis area is different from the PR observations if a simple rain classification method with a threshold of 10 mm hr^{-1} is used. However, when we classify based on rain area, the NICAM simulation captured the characterization of stratiform rain type in the morning and convective rain type in the afternoon. For the rainfall rate intensity, the NICAM simulation overestimates the morning peak. This difference from the PR observation could be related to the known bias of the NICAM simulation of reproducing stronger convective rainfall rate than observed (Masunaga et al.; 2006, Satoh et al. 2010; Nasuno and Satoh, 2011). The difference might also be related to the time difference between the NICAM one-week simulation and the 17-year dataset of the TRMM observations.

428

4.2 Amazon Region

430 As shown in Fig. 1, the spatial distribution of mean rainfall rate over the Amazon is simulated well
431 by NICAM, although the rainfall rate of NICAM is higher (warmer colors) than in the TRMM PR
432 observations. Here, we analyze the characteristics of daily variation in rainfall rate, focusing on the
433 area of (75°–65°W, Eq.–10°S), to avoid the Andes Mountain area. Figure 8 shows the day-to-day
434 variation in mean rainfall rate over the area of the NICAM simulation (red) during 26–31 December
435 2006 compares it to GSMaP data. The NICAM simulation indicates clear semidiurnal variation (i.e.,
436 two-peaks a day) except for 30 December 2006. The GSMaP (purple) shows two local peaks per day
437 on 26, 27, 29, and 31 December.

438 Figure 9 shows the mean daily variation in rainfall rate in NICAM (red), PR (blue), TMI (green),
439 and GSMaP (purple) over the Amazon region. NICAM and PR clearly show two local peaks, but the
440 morning peaks in the TMI and GSMaP are not clear. The NICAM simulation shows significantly
441 higher rainfall rate than the TRMM and GSMaP daily observations. *In situ* observation over the
442 Amazon is provided by a Peruvian project of the Hydro-geochemistry of the Amazonian Basin
443 (HYBAM) (<https://hybam.obs-mip.fr/data/daily-rainfall-1980-2009-over-the-entire-amazon-basin/>).
444 The mean rainfall rate over the analysis area is estimated as 0.32 mm hr⁻¹ for December 26–31 2006.
445 Figure 9 indicates that this value of the observed mean rainfall rate is closer to that of the TRMM PR,
446 TMI, and GSMaP data.

447 Both TRMM PR and NICAM show the semidiurnal variation in rainfall rate, although the NICAM
448 simulation shows a higher rainfall rate than PR. The semidiurnal variation is not clear in GSMaP; in

449 particular, the morning peak is unclear. The NICAM simulation shows the primary peak at 15 LST
450 and a secondary peak at around 3–6 LST. TRMM PR indicates the primary peak at around 13.5–15.5
451 LST and a secondary peak at 2.5 LST. In TMI, the primary peak appears at 15.5 LST. The amplitude
452 of rainfall rate is similar between PR and TMI.

453 Figure 10a shows the daily variation in size (green) and number (blue) of DCs defined by 235K
454 TB over the Amazon using TRMM ancillary data. Note that we used 235 K to define DC over the
455 Amazon because cold clouds (< 213 K), which was used for southern Africa (Fig. 7), are rare over
456 the Amazon during this period. The size of the DC indicates a peak at 21.5 LST. There is also one
457 tiny peak at 4.5 LST. The number of DC areas shows one afternoon peak at 15.5 LST. The afternoon
458 peak of rainfall observed by TRMM at around 11.5–14.5 LST (Fig.9) corresponds to the timing at
459 which a large number of DC areas increase in size. The morning peak at around 3.5–6.5 LST in PR
460 is after the size of DC areas becomes largest at 21.5–24 LST. The daily variation in size and number
461 of DC areas defined by 253 K of TB infrared data are consistent with those found over southern
462 Africa (Fig.7a) for the afternoon rainfall peak.

463 Figure 10b shows a diurnal variation in size (green) and number (blue) of DC areas defined by
464 NICAM OLR less than 150 W m^{-2} . There is one broad peak of the size of DC area during the night,
465 although there is a tiny minimum at 22.5 LST. The peak of the rainfall rate in the afternoon at 15 LST
466 (Fig. 9) corresponds to the larger number of DC areas. This feature is similar to the observed evolution
467 of the size and number of DC areas.

468 Figure 10c shows the diurnal variation in convective (blue) and stratiform (green) rain in the PR
469 observation (solid line) and the NICAM simulation (dotted line). The convective rain in PR indicates
470 a semidiurnal mode with a primary peak at 13.5 LST and a secondary peak at 2.5 LST, although the
471 morning peak is small. In the PR observation, stratiform rainfall rate is higher during the late evening
472 to morning compared with convective rainfall rate.

473 Daily variation in convective ($> 10 \text{ mm hr}^{-1}$) and stratiform ($< 5 \text{ mm hr}^{-1}$) rainfall rate simulated
474 by NICAM is shown in blue and green dotted lines, respectively, in Fig 10c. The primary peak of
475 convective rainfall rate in the NICAM simulation is slightly later than that in the PR observation. The
476 morning peak of convective rainfall rate is higher in the NICAM simulation than in the PR
477 observation, similar to that seen over southern Africa (Fig. 7d). The stratiform rainfall rate in the
478 NICAM simulation is comparable to that in the PR observation, although it is much lower than the
479 convective rainfall rate in the NICAM simulation.

480 Figure 10d shows the daily variation in the rain area size (right ordinate) and the mean percentage
481 of convective rain area (left ordinate) within the rain area. The rain area size is large during the
482 morning over the Amazon. The mean percentage of convective rain area within the rain area is high
483 during the afternoon, similar to that seen over southern Africa (Fig. 7d). Although the classification
484 of convective/stratiform rain types is not unique, if we use the percentage of convective rain within
485 each rain area to define each rain type, then the NICAM simulation indicates that the afternoon peak

486 is characterized by convective rain, whereas the morning peak is characterized by stratiform rain,
487 consistent with the PR observation.

488

489 **5. Conclusions**

490 We analyzed the semidiurnal (two peaks per day) variation in surface rainfall rate simulated by the
491 NICAM 3.5-km mesh global cloud resolving model for 26-31 December 2006 and compared it with
492 observations from TRMM PR/TMI, GSMaP, and infrared data from geostationary satellites. The
493 hourly data on a 0.05° latitude/longitude grid from the NICAM one-week simulation and infrared
494 data from a geostationary satellite are constructed for 26-31 December 2006. We also construct the
495 climatological hourly data on a $0.5^\circ \times 0.5^\circ$ latitude-longitude grid from 17 years of 3G68 data of TMI
496 and PR observations during December, January, and February. The global hourly data with $0.1^\circ \times$
497 0.1° latitude-longitude grids of GSMaP for the same period of the NICAM simulation are used as
498 well. The timing of the two peaks may be influenced by many factors such as cloud–radiation
499 interactions, air–sea and air–land interactions and large-scale disturbances. The difference in the
500 analysis period between the NICAM one-week simulation and the 17-year TRMM climatology may
501 also affect the nature of the semidiurnal variation in surface precipitation. Thus, we also used the
502 hourly data of GSMaP for the same one-week period as the NICAM simulation. The rainfall rate of
503 the TRMM PR/TMI is more reliable, while the quality of the GSMaP rainfall may be affected by the
504 availability of satellite passive microwave measurements, which are used to calibrate rain retrievals

505 from Infrared measurements.

506 We first confirm the persistent nature of the semidiurnal variation in mean rainfall rate in the
507 daily time series on a $10^\circ \times 15^\circ$ latitude-longitude grid array over the tropics (20°N – 20°S) using
508 TRMM data. Using harmonic analysis, we show that about 45% of grids show an amplitude of the
509 semidiurnal mode that is more than two-thirds of the amplitude of the diurnal mode in daily variation.
510 The TRMM data implies that there are generally two peaks of rainfall rate a day over the tropics.

511 The semidiurnal variation in rainfall is analyzed over land areas including southern Africa
512 and the Amazon. Over southern Africa, the mean rainfall data in the NICAM simulation have two
513 peaks a day, with a primary peak in the afternoon at 15 LST and a secondary peak in the morning at
514 4.5 LST; in PR (TMI) observations, there is a primary peak in the afternoon at 15.5 (15.5) LST and a
515 secondary peak at 3.5 (0.5) LST. Similarly in the other areas, the primary and secondary peaks
516 simulated by NICAM are very close to those observed by PR/TMI, although the timing of the two
517 peaks are different. Generally, the rainfall rate is higher in the NICAM simulation than in the PR/TMI
518 observations, but the rainfall rate of NICAM simulation is closer to the GSMaP rainfall rate over
519 some areas particularly over the oceans.

520 We then analyzed the characteristics of daily variation by evolution of DC over southern
521 Africa and the Amazon, since DC is closely related to rainfall over the tropics. The DC area is defined
522 as grids with the specific thresholds of the observed TBs and OLR of the NICAM simulation. The
523 temporal variation and spatial distribution of DCs defined by NICAM is similar for DC defined by

524 TB. The size and number of DC areas within the analysis area of southern Africa and the Amazon are
525 analyzed as well. The two peaks in the morning and afternoon are seen in the mean size and number
526 of DC area, both in the observation and the simulation over southern Africa, although they are not
527 clear over the Amazon. The primary peak of total rainfall rate in the afternoon in PR and TMI
528 coincides with the time at which there are a large number of DC areas whose mean size are rapidly
529 increasing over both southern Africa and the Amazon (Figs. 7a, b and Figs. 10a, b).

530 Diurnal variation in convective and stratiform rainfall rates is also examined over southern Africa
531 and the Amazon using PR data. The present analysis shows that convective rainfall rate is significant
532 in the afternoon, while stratiform rainfall rate is larger than convective rainfall rate in the morning
533 over both southern Africa and the Amazon in the PR observation. Although the classification of
534 convective and stratiform rains in the simulation is not unique, if a simple rainfall threshold is used
535 for this classification, then the afternoon peak of convective rainfall rate in NICAM is close to that in
536 the PR observation. However, the daily variation in stratiform rainfall rate in NICAM is generally
537 smaller than the daily variation in convective rainfall rate in NICAM.

538 As another method of classification of rain types, we analyzed the rain area size and the percentage
539 of convective rain within the rain area simulated by NICAM over southern Africa and the Amazon.
540 The mean size of the rain area simulated by NICAM is large in the morning. The mean percentage of
541 convective rain area simulated by NICAM within each rain area is large in the afternoon (Figs. 7d,

542 10d). These results indicate that NICAM could be able to reproduce stratiform rain occurrence in the
543 morning and convective rain occurrence in the afternoon.

544 Although the classification of convective and stratiform rain types in the NICAM simulation is
545 somewhat arbitrary, stratiform rainfall rates tend to be lower and convective rainfall rates are
546 generally higher. This characteristic of rainfall in the NICAM simulation is similar to that reported in
547 the previous studies of NICAM (Masunaga et al. 2006; Satoh et al. 2010; Nasuno and Satoh 2011).
548 The improvement of precipitation in the numerical model needs more evaluation using observation
549 data and detailed process studies, and the cloud microphysics schemes in NICAM are being improved
550 (Satoh et al. 2014; Roh and Satoh 2014; Roh et al. 2017; Kodama et al. 2021).

551 In this study, as a representative of global cloud-resolving models, we showed for the first time
552 that NICAM simulates the major features of the semidiurnal variation in rainfall and DCs in realistic
553 configuration. Our analysis shows that spatial and temporal variation in DCs simulated by NICAM
554 corresponds well to the DCs defined by TBs using infrared data from geostationary satellites. The
555 benefit of NICAM is that evolution of DCs is reproduced well, where a DC is associated with
556 mesoscale circulation which consists of convective and stratiform rains. We believe that accurate
557 simulations of DCs are required to reproduce the semidiurnal variation in surface rainfall rate,
558 although DCs and rainfall are different. For the morning peak, however, NICAM overestimates
559 rainfall rate over land areas. Continued efforts to improve cloud and precipitation processes are

560 required to better reproduce the semidiurnal variation in rainfall rate, particularly for the
561 representation of stratiform rainfall in NICAM.

562

563 **Supplement**

564 Analyses on the eastern Pacific Ocean areas (S.1) and the maritime continent area (S.2) are given
565 with Figures S1-S10.

566

567 **Appendix: Calculating the number and mean size of DC areas**

568 A DC area is defined as an area comprising all connected adjacent grids that have a TB
569 below 213 K. The suitable value of the threshold of TB may depend on the analysis target. The size
570 of each DC area is defined by the number of grids in that area. The center of each DC area is
571 located by averaging the longitude and latitude of all grids in that area. The number and mean size
572 of DC areas are counted only for those DC areas whose centers are located within the analysis area.
573 Figure A shows a sample case for the definitions of the size and number of DC areas.

574

575 **Acknowledgments**

576 The NICAM simulation was carried out on the Earth Simulator. We acknowledge anonymous
577 reviewers for their comments and support which improved the clarity of our paper. KR acknowledges

578 the support from the National Monsoon Mission Phase-II Project (GAP-1013) of Ministry of Earth
579 Sciences (MoES), India.

580

581

References

582 Andersson, T., 1970: The diurnal variation of precipitation frequency over Weather Ship. *J. Appl.*
583 *Meteor.*, **9**, 17–19.

584 Arkin, P. A., and B. N. Meisner, 1987: The relationship between large-scale convective rainfall and
585 cold cloud over the western Hemisphere during 1982-84. *Mon. Wea. Rev.*, **115**, 51–74.

586 Awaka, J., T. Iguchi, H. Kumagai, and K. Okamoto, 1997: Rain type classification algorithm for
587 TRMM Precipitation Radar. *Proc. IGARSS'97*, Singapore, IEEE, 1633–1635.

588 Awaka, J., T. Iguchi, and K. Okamoto, 2007: Rain type classification algorithm. *Measuring*
589 *Precipitation from Space: EURAINSAT and the Future*. V. Levizzani, P. Bauer, and F. J. Turk
590 (Eds.), Springer, 213–224.

591 Brier, G. W., and J. Simpson, 1969: Tropical cloudiness and rainfall related to pressure and tidal
592 variations. *Quart. J. Roy. Meteor. Soc.*, **95**, 120-147.

593 Covey, C., P. J. Gleckler, C. Doutriaux, D. N. Williams, A. Dai, J. Fasullo, K. Trenberth, and A. Berg,
594 2016: Metrics for the diurnal cycle of precipitation: Toward routine benchmarks for climate
595 models. *J. Climate*, **29**, 4461–4471.

596 Dai, A., 2001: Global precipitation and thunderstorm frequencies, Part II: Diurnal Variations. *J.*

597 Climate, **14**, 1112–1128.

598 Del Genio, A.D., and J. Wu, 2010: The role of entrainment in the diurnal cycle of continental
599 convection. *J. Climate*, **23**, 2722–2738.

600 Dirmeyer, P. A., B. A. Cash, J. L. Kinter III, T. Jung, L. Marx, M. Satoh, C. Stan, H. Tomita, P. Towers,
601 N. Wedi, D. Achuthavarier, J. M. Adams, E. L. Altshuler, B. Huang, E. K. Jin, and J.
602 Manganello, 2011: Simulating the diurnal cycle of rainfall in global climate models: Resolution
603 versus parameterization. *Clim. Dyn.*, **39**, 399–418.

604 Furuzawa, F. A., and K. Nakamura, 2005: Differences of rainfall estimates over land by Tropical
605 Rainfall Measuring Mission (TRMM) Precipitation Radar (PR) and TRMM Microwave
606 Imager (TMI) -Dependence on storm height. *J. Appl. Meteor.*, **44**, 367–382.

607 Grabowski, W. W., 1998: Toward cloud resolving modeling of large-scale tropical circulations: A
608 simple cloud microphysics parameterization. *J. Atmos. Sci.*, **55**, 3283–3298.

609 Grabowski, W.W., 2003: Impact of ice microphysics on multiscale organization of tropical convection
610 in two-dimensional cloud-resolving simulations. *Q. J. R. Meteorol. Soc.*, 129, 67–81.

611 Gray, W. M., and R. W. Jacobson, 1977: Diurnal variation of deep cumulus convection. *Mon. Wea.*
612 *Rev.*, **105**, 1171–1188.

613 Haddad, Z. S., E. A. Smith, C. D. Kummerow, T. Iguchi, M. R. Farrar, S. L. Durden, M. Alves, and
614 W. S. Olson, 1997: The TRMM “Day-1” radar/radiometer combined rain-profile algorithm. *J.*
615 *Meteor. Soc. Japan*, **75**, 799–809.

616 Hamilton, K., 1981: Latent heat release as a possible forcing mechanism for atmospheric tides. Mon.
617 Weather Rev., **109**, 3—17.

618 Hamilton, K., 1981: A Note on the Observed Diurnal and Semidiurnal Rainfall Variations. J. Geophys.
619 Res., **86**, 12,122-12,1226.

620 Hirose, M., and K. Nakamura, 2005: Spatial and diurnal variation of precipitation systems over Asia
621 observed by the TRMM Precipitation Radar. J. Geophys. Res., **110**, D05106, doi:
622 10.1029/2004JD004815.

623 Hou, A.Y., R. K. Kakar, S. Neeck, A. A. Azarbarzin, C. D. Kummerow, M. Kojima, R. Oki, K.
624 Nakamura, and T. Iguchi, 2014: The Global Precipitation Measurement Mission. Bulletin of
625 the American Meteorological Society, **95**, 701–722.

626 Iguchi, T., T. Kozu, R. Meneghini, J. Awaka, and K. Okamoto, 2000: Rain-profiling algorithm for
627 the TRMM precipitation radar. J. Appl. Meteor., **39**, 2038–2052.

628 Inoue, T., 1985: On the temperature and effective emissivity determination of semi-transparent cirrus
629 clouds by bi-spectral measurements in the 10 μ m window region. J. Meteor. Soc. Japan, **63**,
630 88–99.

631 Inoue, T., 1987a: A cloud type classification with NOAA-7 split window measurements. J. Geophys.
632 Res., **92**, 3991–4000.

633 Inoue, T., 1987b: An instantantaneous delineation of convective rainfall area using split window data
634 of NOAA-7 AVHRR. J. Meteor. Soc. Japan, **65**, 469-481.

635 Inoue, T., M. Satoh, H. Miura, and B. Mapes, 2008: Characteristics of cloud size of deep convection
636 simulated by a global cloud resolving model. *J. Meteor. Soc., Japan*, **86A**, 1-15.

637 Inoue, T., D. Vila, K. Rajendran, A. Hamada, X. Wu, and L. A. Machado, 2009: Life cycle of deep
638 convective systems over the eastern tropical Pacific observed by TRMM and GOES-W. *J.*
639 *Meteor. Soc. Japan*, **87A**, 381–391.

640 Inoue, T., K. Rajendran, M. Satoh, H. Miura, and J. Schmetz, 2012: Semi-diurnal variation of surface
641 rainfall studied from global cloud-system resolving model and satellite observations, *Proc.*
642 *SPIE 8529, Remote Sensing and Modeling of the Atmosphere, Oceans, and Interactions IV*,
643 85290K (8 November 2012), doi: 10.1117/12.977609.

644 Janowiak, J., P. A. Arkin, and M. Morrissey, 1994: An examination of the diurnal cycle in oceanic
645 tropical rainfall using satellite and in situ data. *Mon. Wea. Rev.*, **122**, 2296–2311.

646 Janowiak, J., B. Joyce, and P. Xie, 2017: NCEP/CPC L3 Half hourly 4km global (60S-60N) merged
647 IR V1, Edited by Andrey Savtchenko, Greenbelt, MD, Goddard Earth Sciences Data and
648 Information Services Center.

649 Kikuchi, K., and B. Wang, 2008: Diurnal precipitation regimes in the global tropics. *J. Climate*, **21**,
650 2680–2696.

651 Kodama, C., T. Ohno, T. Seiki, H. Yashiro, A. T. Noda, M. Nakano, Y. Yamada, W. Roh, M. Satoh,
652 T. Nitta, D. Goto, H. Miura, T. Nasuno, T. Miyakawa, Y.-W. Chen, and M. Sugi, 2021: The
653 Nonhydrostatic ICosahedral Atmospheric Model for CMIP6 HighResMIP simulations

654 (NICAM16-S): Experimental design, model description, and impacts of model updates. *Geosci.*
655 *Model Dev.*, **14**, 795–820.

656 Kraus, E. B., 1963: The diurnal precipitation change over the sea. *J. Atmos. Sci.*, **20**, 546–551.

657 Kubota, T., K. Aonashi, T. Ushio, S. Shige, Y. N. Takayabu, M. Kachi, Y. Arai, T. Tashima, T. Masaki,
658 N. Kawamoto, T. Mega, M. K. Yamamoto, A. Hamada, M. Yamaji, G. Liu, and R. Oki, 2020:
659 Global Satellite Mapping of Precipitation (GSMaP) products in the GPM era. *Satellite*
660 *Precipitation Measurement*. Levizzani, V., C. Kidd, D. B. Kirschbaum, C. D. Kummerow, K.
661 Nakamura, and F. J. Turk (eds.), Springer, Cham, 355–373.

662 Kummerow, C., Y. Hong, W. S. Olson, S. Yang, R. F. Adler, J. McCollum, R. Ferraro, G. Petty, D. B.
663 Shin, and T. T. Wilheit, 2001: The evolution of the Goddard profiling algorithm (GPROF) for
664 rainfall estimation from passive microwave sensors. *J. Appl. Meteor.*, **40**, 1801–1840.

665 Lang, S., W.-K. Tao, J. Simpson, and B. Ferrier, 2003: Modeling of convective-stratiform
666 precipitation processes: Sensitivity to partitioning methods. *J. Applied. Meteor.*, **42**, 505-527.

667 Lin, X, D. A. Randall, and L. D. Fowler, 2000: Diurnal variability of the hydrologic cycle and
668 radiative fluxes: comparisons between observations and a GCM. *J. Clim.* **13**, 4159–4179.

669 Lindzen, R. S., 1978: Effect of daily variations in cumulonimbus activity on the atmospheric
670 semidiurnal tide. *Mon. Wea. Rev.*, **106**, 526— 533.

671 Lindzen, R. S., and S. Chapman, 1969: Atmospheric Tides. *Space Sci. Rev.*, **10**, 3-188.

672 Mapes, B. E., and R. A. House, Jr., 1993: Cloud clusters and superclusters over the oceanic warm

673 pool. Mon. Wea. Rev., **121**, 1398–1416.

674 Masunaga, H., M. Satoh, and H. Miura, 2008: A joint satellite and global CRM analysis of an MJO
675 event: Model diagnosis. J. Geophys. Res., **113**, D17210, doi:10.1029/2008JD009986.

676 Mega, T., T. Ushio, T. Matsuda, T. Kubota, M. Kachi, and R. Oki, 2019: Gauge-adjusted Global
677 Satellite Mapping of Precipitation (GSMaP_Gauge). IEEE Transactions on Geoscience and
678 Remote Sensing, **57**, 1928-1935.

679 Miura, H., M. Satoh, T. Nasuno, A. T. Noda, and K. Oouchi, 2007: A Madden-Julian Oscillation event
680 simulated using a global cloud-resolving model. Science, **318**, 1763–1765.

681 Mori, S., J. Hamada, Y. I. Tauhid, M. D. Yamanaka, N. Okamoto, F. Murata, N. Sakurai, H.
682 Hashiguchi, and T. Sribimawati, 2004: Diurnal land–sea rainfall peak migration over Sumatera
683 Island, Indonesian Maritime Continent, observed by TRMM satellite and intensive rawinsonde
684 soundings. Mon. Wea. Rev., **132**, 2021–2039.

685 Nasuno, T., H. Miura, M. Satoh, A. T. Noda, and K. Oouchi, 2009: Multi-scale organization of
686 convection in a global numerical simulation of the December 2006 MJO event using explicit
687 moist processes. J. Meteor. Soc. Japan, **87**, 335-345.

688 Nasuno, T., and M. Satoh, 2011: Statistical relation between maximum vertical velocity and surface
689 precipitation of tropical convective clouds in a global nonhydrostatic aquaplanet experiment. J.
690 Meteor. Soc. Japan, **89**, 553-561.

691 Nitta, T., and S. Sekine, 1994: Diurnal variation of convective activity over the tropical western

692 Pacific. J. Meteorol. Soc. Japan, **72**, 627– 641.

693 Noda, A. T., K. Oouchi, M. Satoh, and H. Tomita, 2012: Quantitative assessment of diurnal variation
694 of tropical convection simulated by a global nonhydrostatic model without cumulus
695 parameterization. *J. Climate*, **25**, 5119-5134.

696 Rajendran, K., and T. Nakazawa, 2005: Systematic difference between TRMM 3G68 PR and TMI
697 rainfall estimates and the possible association with life cycle of convection. *SOLA*, **1**, 165–168.

698 Roh, W., and Satoh, M., 2014: Evaluation of precipitating hydrometeor parameterizations in a single-
699 moment bulk microphysics scheme for deep convective systems over the tropical open ocean.
700 *J. Atmos. Sci.*, **71**, 2654-2673.

701 Roh, W., M. Satoh, and T. Nasuno, 2017: Improvement of a cloud microphysics scheme for a global
702 nonhydrostatic model using TRMM and a satellite simulator. *J. Atmos. Sci.*, **74**, 167-184.

703 Sato, T., H. Miura, M. Satoh, Y. N. Takayabu, and Y. Wang, 2009: Diurnal cycle of precipitation in
704 the tropics simulated in a global cloud-resolving model. *J. Climate*, **22**, 4809–4826.

705 Satoh, M., T. Matsuno, H. Tomita, H. Miura, T. Nasuno, and S. Iga, 2008b: Nonhydrostatic
706 Icosahedral Atmospheric Model (NICAM) for global cloud resolving simulations. *J. Comp.*
707 *Phys.*, **227**, 3486–3514.

708 Satoh M., T. Nasuno, H. Miura, H. Tomita, S. Iga, and Y. Takayabu, 2008b: Precipitation Statistics
709 Comparison Between Global Cloud Resolving Simulation with NICAM and TRMM PR Data.
710 *Precipitation statistics comparison between global cloud resolving simulation with NICAM and*

711 *TRMM PR data*. Hamilton K., and Ohfuchi W. (eds), Springer, 99-112.

712 Satoh, M., T. Inoue, and H. Miura, 2010: Evaluations of cloud properties of global and local cloud
 713 system resolving models using CALIPSO/CloudSat simulators. *J. Geophys. Res.*, **115**, D00H14,
 714 doi:10.1029/2009JD012247.

715 Satoh, M., H. Tomita, H. Yashiro, H. Miura, C. Kodama, T. Seiki, A. T. Noda, Y. Yamada, D. Goto,
 716 M. Sawada, T. Miyoshi, Y. Niwa, M. Hara, Y. Ohno, S. Iga, T. Arakawa, T. Inoue, and H.
 717 Kubokawa, 2014: The Non-hydrostatic Icosahedral Atmospheric Model: Description and
 718 development. *Progress in Earth and Planetary Science*. 1, 18, doi: 10.1186/s40645-014-0018-1.

719 Satoh, M., B. Stevens, F. Judd, M. Khairoutdinov, S. Lin, W. M. Putman, and P. Dben, 2019: Global
 720 cloud-resolving models. *Current Climate Change Reports*, **5**, 172-184.

721 Schmetz, J., P. Pili, S. Tjemkes, D. Just, J. Kerkmann, S. Rota, and A. Eaiter, 2002: An introduction
 722 to METEOSAT Second Generation (MSG). *Bull. Am. Meteorol. Soc.*, **83**, 977–992.

723 Schroder, M., M. Konig, and J. Schmetz, 2009: Deep convection observed by the spinning enhanced
 724 visible and infrared imager on board METEOSAT-8: Spatial distribution and temporal evolution
 725 over Africa in summer and winter 2006. *J. Geophys. Res.*, **114**, D05109, doi:
 726 10.1029/2008JD010653.

727 Seto, S., T. Iguchi, R. Meneghini, J. Awaka, T. Kubota, T. Masaki, and N. Takahashi, 2021: The
 728 Precipitation rate retrieval algorithms for the GPM Dual-frequency Precipitation Radar. *J.*
 729 *Meteor. Soc. Japan*, **99**, <https://doi.org/10.2151/jmsj.2021-011>.

730 Stevens, B., M. Satoh, L. Auger, J. Biercamp, C. Bretherton, X. Chen, P. Duben, F. Judt, M.
731 Khairoutdinov, D. Klocke, C. Kodama, L. Kornbluh, S.-L. Lin, W. Putman, R. Shibuya, P.
732 Neumann, N. Rober, B. Vannier, P.-L. Vidale, N. Wedi, L. Zhou, 2019: DYAMOND: The
733 DYnamics of the Atmospheric general circulation Modeled On Non-hydrostatic Domains.
734 Progress in Earth and Planetary Science, 6, 61, <https://doi.org/10.1186/s40645-019-0304-z>.

735 Sui, C.-H., K.-M. Lau, Y. N. Takayabu, and D. A. Short, 1997: Diurnal variations in tropical oceanic
736 cumulus convection during TOGA COARE. J. Atmos. Sci., **54**, 639–655.

737 Takayabu, Y. N., 2002: Spectral representation of rain profiles and diurnal variations observed with
738 TRMM PR over the equatorial area. Geophys. Res. Lett., **29**, 1584, doi:10.1029/2001GL014113.

739 Takayabu, Y. N., and K. Kimoto, 2008: Diurnal march of rainfall simulated in a T106 AGCM and
740 dependence on cumulus schemes. J. Meteor. Soc. Japan, **86A**, 163–173.

741 Tomita, H., and M. Satoh, (2004) A new dynamical framework of nonhydrostatic global model using
742 the icosahedral grid. Fluid Dyn. Res., **34**, 357-400.

743 Tomita, H., H. Miura, S. Iga, T. Nasuno, and M. Satoh, 2005: A global cloud-resolving simulation:
744 Preliminary results from an aqua planet experiment. Geophys. Res. Lett., **32**, L08805,
745 doi:10.1029/2005GL022459.

746 Ushio, T., K. Sasashige, T. Kubota, S. Shige, K. Okamoto, K. Aonashi, T. Inoue, N., Takahashi, T.
747 Iguchi, M. Kachi, R. Oki, T. Morimoto, and Z.-I. Kawasaki, 2009: A Kalman filter approach to
748 the Global Satellite Mapping of Precipitation (GSMaP) from combined passive microwave and

749 infrared radiometric data. *J. Meteor. Soc. Japan*, 87A, 137–151.

750 Wallace, J. M., 1975: Diurnal variations in precipitation and thunderstorm frequency over the
751 conterminous United States. *Mon. Wea. Rev.*, 103, 406–419.

752 Woolnough, S., J., J. M. Slingo, and B. J. Hoskins, 2004: The diurnal cycle of convection and
753 atmospheric tides in an aquaplanet GCM. *J. Atmos. Sci.*, **61**, 2559–2573.

754 Yamamoto, M. K., F. A. Furuzawa, A. Higuchi, and K. Nakamura, 2008: Comparison of diurnal
755 variation in precipitation system observed by TRMM PR, TMI, and VIRS. *J. Climate*, **21**, 4011–
756 4028.

757 Yang, G-Y, and J. Slingo, 2001: The diurnal cycle in the tropics. *Mon. Wea. Rev.*, 129, 784–801

758 Yang, S., and E. A. Smith, 2006: Mechanisms for diurnal variability of global tropical rainfall
759 observed from TRMM. *J. Climate*, **19**, 5190–5226.

760 Yang, S., and E. A. Smith, 2008, Convective-stratiform precipitation variability at seasonal scale from
761 eight years of TRMM observations: Implications for multiple modes of diurnal variability. *J.*
762 *Clim.*, **21**, 4087–4114.

763 Yang, S., K-S Kuo, and E. A. Smith, 2008: Persistent Nature of Secondary Diurnal Modes of
764 Precipitation over Oceanic and Continental Regimes. *J. Clim.*, **21**, 4115–4131.

765 Yang, S., and S. W. Nesbitt, 2014, Statistical properties of precipitation as observed by the TRMM
766 precipitation radar. *Geophys. Res. Lett.*, **41**, 5636–5643.

767 Yashiro, H., Y. Kajikawa, Y. Miyamoto, T. Yamaura, R. Yoshida, and H. Tomita, 2016: Resolution
768 dependence of the diurnal cycle of precipitation simulated by a global cloud-system resolving
769 model. SOLA, **12**, 272-276.

770 Yasunaga, K., T. Nasuno, H. Miura, Y. N. Takayabu, and M. Yoshizaki, 2013: Afternoon
771 precipitation peak in an Aqua-planet global cloud-resolving model (Aqua-planet NICAM). J.
772 Meteor. Soc. Japan, **91A**, 217-229.

773 Yuan, W. H., R. C. Yu, M. H. Zhang, W. Y. Lin, J. Li, and Y. F. Fu, 2013: Diurnal cycle of summer
774 precipitation over subtropical East Asia in CAM5. J. Clim., **26**, 3159–3172.

775

776 Figure captions

777

778 Figure 1:

779 (a) Spatial distribution of mean surface rainfall rate averaged for the period 00UTC on 26
780 December 2006 to 24 UTC on 31 December 2006 by TRMM PR. (b) Same as (a) but for TMI.
781 (c) Same as (a) but for the NICAM simulation.

782

783 Figure 2:

784 (a) Map of the time series of mean rainfall over the $10^{\circ} \times 15^{\circ}$ latitude-longitude grid arrays (boxes)
785 over the tropics (20°N-20°S) constructed from the TMI (green) and PR (blue) observations. In each
786 grid box, x-axis spans 0-23 LST, and the y-axis is rainfall rate normalized to the maximum rainfall

787 rate that box. (b) Distribution of the amplitude ratio of (semidiurnal variation)/(diurnal variation)
788 derived from the harmonic analysis for TRMM PR (top) and TRMM TMI (bottom). (c) Same as (a)
789 but for the NICAM simulation. (d) Same as (b) but for the NICAM simulation.

790

791

792

793

794

795

796

797 Figure 3:

798 Day-to-day variation of mean rainfall rate averaged over southern Africa (15° - 28° E, Eq.- 20° S) during
799 26 – 31 December 2006 represented by the NICAM simulation (red) and GSMaP (purple) in UTC.
800 The x-axis starts at 0.5 LST with vertical lines indicating 3-hour periods.

801

802 Figure 4:

803 Composite of daily variation in mean rainfall rate over the southern Africa (15° - 28° E, Eq.- 20° S) by
804 PR (blue), TMI (green), GSMaP (purple), and the NICAM simulation (red).

805

806 Figure 5:

807 (a) Spatial distribution of DC observed over southern Africa by Meteosat-8 infrared data at 03, 06,
808 09, 12, 15, 18, 21, and 24 UTC on 26 December 2006. (Note that LST over this area is close to UTC.)

809 DC is defined by TB colder than 253K. Optically thick (cumulonimbus type) cloud shown in blue
810 and optically thinner (cirrus type) cloud (green) are classified by the split-window method. Cloud
811 colder than 213K within DC is shown in purple. (b) Same as (a) but for the NICAM simulation. DC
812 is defined by OLR less than 210 W m^{-2} .

813

814 Figure 6:

815 Daily variation in cold cloud (DC) grid number anomaly constructed from the 6-day period of 26 -
816 31 December 2006 over Southern Africa (15° - 28° E, Eq.- 20° S). Cold cloud is defined by various TB
817 from Meteosat-8 data. Five classes of DC are defined: three classes colder than the threshold TB of
818 213 K (purple), 235 K (blue), 253 K (green) and two other classes 253 K-Cb (red) and 253 K-Ci
819 (green dashed), for optically thick and thin cloud colder than 253K, respectively.

820

821 Figure 7:

822 Daily variation in characteristics of DC over southern Africa (15° - 28° E, Eq.- 20° S). (a) Daily
823 variation in mean size (green) and number (blue) of DC defined by TB of infrared data from Meteosat-
824 8 constructed for 26–31 December 2006. The DC area is defined by the connected grids colder than
825 213 K. (b) Same as (a) except for OLR less than 102 W m^{-2} (dotted line) in the NICAM simulation.(c)
826 Daily variation in rainfall rate of convective rain (blue) and stratiform rain (green) observed by
827 TRMM PR (solid line) or simulated by NICAM (dotted line). Convective rain in the NICAM
828 simulation is defined as rainfall rate higher than 10 mm hr^{-1} and stratiform rain is defined as rainfall
829 rate lower than 5 mm hr^{-1} . (d) Daily variation in the size of the rain area in grid number (green: right

830 ordinate) and the mean percentage of convective rain area within each rain area (blue: left ordinate)
831 in the NICAM simulation. Rain area is defined as the connected rain grid.

832

833 Figure 8:

834 Same as Fig. 3 but for over the Amazon (75°-65°W, Eq.-10°S). The x-axis starts at 19.5 LST.

835

836 Figure 9:

837 Same as Fig. 4 but for over the Amazon (75°-65°W, Eq.-10°S).

838

839 Figure 10:

840 Same as Fig. 7 but for the Amazon (75°-65°W, Eq.-10° S) with DCs defined by the threshold values
841 of 235 K and 150 W m⁻² for TB of TRMM ancillary data and OLR of NICAM, respectively.

842

843 Figure A:

844 A sample case for calculating the number and mean size of DC areas. In this case, four DC areas are
845 defined: DC-1, DC-2, DC-3, and DC-4. However, as the center of DC-1 is outside the analysis area
846 (indicated by the rectangle), the number of DC areas is three. The mean size of DC areas is the average
847 of the sizes of DC-2, DC-3, and DC-4.

848

Figure1:

- (a) Spatial distribution of mean surface rainfall rate averaged for the period 00UTC on 26 December 2006 to 24 UTC on 31 December 2006 by TRMM PR. (b) Same as (a) but for TMI. (c) Same as (a) but for the NICAM simulation.

(a)

DJF TMI/PR Normalised Rain (LST)

(b)

Amplitude Ratio of Semidiurnal/Diurnal (PR)

Amplitude Ratio of Semidiurnal/Diurnal (TMI)

(c)

(d)

Figure 2:

(a) Map of the time series of mean rainfall over the $10^\circ \times 15^\circ$ latitude/longitude grid arrays

(boxes) over the tropics (20°N-20°S) constructed from the TMI (green) and PR (blue) observations constructed from 17 years TRMM data. In each grid box, x-axis spans 0-23 LST, and y-axis is rainfall rate normalized by the maximum rainfall rate at each grid box (0-1).

- (b) Distribution of the amplitude ratio of (semidiurnal variation)/(diurnal variation) derived from the harmonic analysis for TRMM PR (top) and TMI (bottom).
- (c) Same as (a) but for the NICAM simulation.
- (d) Same as (b) but for the NICAM simulation.

Figure 3:

Day-to-day variation of mean rainfall rate averaged over southern Africa (15°E - 28°E and Eq. - 20°S) during 26 – 31 December 2006 represented by the NICAM simulation (red) and GSMaP (purple) data in UTC. The x-axis starts at 0.5 LST with vertical lines indicating 3-hour periods.

Figure 4:

Daily variation of mean rainfall rate over the southern Africa (15°E-28°E and Eq.-20°S)

by TRMM PR (blue), TMI (green), GSMaP (purple), and the NICAM simulation (red).

(a)

(b)

Figure 5:

(a) Spatial distribution of DC observed over southern Africa by Meteosat-8 infrared data at 03, 06, 09, 12, 15, 18, 21 and 24UTC on 26 December 2006. (Note that LST over this area is close to UTC.) DC is defined by TB colder than 253K. Optically thick (cumulonimbus type) cloud shown in blue and optically thinner (cirrus type) cloud shown in green classified by the split window method. Cloud colder than 213K within DC is shown in purple. (b) Same as (a) but for the NICAM simulation. DC is defined by OLR smaller than 210 W m^{-2} .

Figure 6:

Daily variation of cold cloud (DC) grid number anomaly constructed from the 6-day period 26-31 December 2006 over Southern Africa (15E-28E; Eq.-20S). Cold cloud is defined by various brightness temperatures of Meteosat-8. Five classes of DCs are defined: the three class are those colder than the threshold brightness temperatures of 213, 235, 253K and the two classes are 253K-Cb and 253K-Ci (optically thick and thin cloud colder than 253K).

Figure 7:

Daily variation of characteristics of DC over southern Africa (15°-28°E, Eq.-20°S). (a)

Daily variation of mean size of DC (green) and number of DC (blue) defined by TB of

infrared data of Meteosat constructed for 26 -31 December 2006. DC is defined as

connected adjacent grids colder than 213K. (b) Same as (a) except for OLR less than 102

W m⁻² (dotted line) in the NICAM simulation. (c) Daily variation of rainfall rate of

convective rain (blue) and stratiform rain (green) observed by TRMM PR (solid line) and

the NICAM simulation (dotted line). Convective rain in NICAM is defined as rainfall rate

greater than 10mm hr⁻¹ and stratiform rain is defined as rainfall rate less than 5 mm hr⁻¹.

(d) Daily variation of the size of the rain area in grid number (green: right ordinate) and

the mean percentage of convective rain area within each rain area (blue: left ordinate) in

the NICAM one-week simulation. Rain area is defined as the adjacent connected rain grid.

Figure 8:

Same as Fig3 but for over Amazon (75° - 65° W and Eq.- 10° S) and x-axis starts at 19.5

LST.

Figure 9:

Same as Fig.4 but for over Amazon (75°-65°W and Eq.-10°S).

Figure 10:

Same as Fig. 7 but for Amazon (75°-65°W, Eq.-10°S) with DCs defined by the threshold

values of 235 K and 150W m^{-2} for TB of TRMM ancillary data and (b) OLR of NICAM, respectively.

Figure A:

A sample case for calculating the number and mean size of DC areas. In this case, four DC areas are defined: DC-1, DC-2, DC-3, and DC-4. However, as the center of DC-1 is outside the analysis area (indicated by the rectangle), the number of DC areas is three. The mean size of DC areas is the average of the sizes of DC-2, DC-3, and DC-4.