EARLY ONLINE RELEASE

This is a PDF of a manuscript that has been peer-reviewed and accepted for publication. As the article has not yet been formatted, copy edited or proofread, the final published version may be different from the early online release.

This pre-publication manuscript may be downloaded, distributed and used under the provisions of the Creative Commons Attribution 4.0 International (CC BY 4.0) license. It may be cited using the DOI below.

The DOI for this manuscript is

DOI:10.2151/jmsj.2022-003

J-STAGE Advance published date: September 30th, 2021

The final manuscript after publication will replace the preliminary version at the above DOI once it is available.

Characteristics of raindrop size distributions during Meiyu season in Mount Lushan, eastern China

Yi CHANG¹

*State Key Laboratory of Severe Weather (LASW) & Key Laboratory for Cloud Physics of
China Meteorological Administration, Chinese Academy of Meteorological Sciences,
Beijing, China*

Qianrong MA

School of Physical Science and Technology, Yangzhou University, Yangzhou, China

Lijun GUO¹, Jing DUAN, Jun LI

*State Key Laboratory of Severe Weather (LASW) & Key Laboratory for Cloud Physics of
China Meteorological Administration, Chinese Academy of Meteorological Sciences,
Beijing, China*

Xiaopeng ZHANG

Lushan Meteorological Bureau, Jiujiang, China

Xueliang GUO

Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, 100029, China

Xiaofeng LOU, and Baojun CHEN

*State Key Laboratory of Severe Weather (LASW) & Key Laboratory for Cloud Physics of
China Meteorological Administration, Chinese Academy of Meteorological Sciences,
Beijing, China*

1) Corresponding author: Yi Chang and Lijun Guo, State Key Laboratory of Severe
Weather (LASW) & Key Laboratory for Cloud Physics of China Meteorological
Administration, Chinese Academy of Meteorological Sciences, Zhong-guan-cun South
Avenue No.46, Haidian District, Beijing 100081
Email: changyi1017@qq.com & ljqiao@cma.gov.cn

Abstract

Meiyu front precipitation makes the region prone to frequent floods, mudslides, landslides, and other disasters, and has been the focus of ongoing and challenging meteorological research. Investigation of the Raindrop size distribution (RSD) is essential for exploring the characteristics and underlying physical precipitation processes. This study investigated the precipitation characteristics in Lushan mountainous areas during the Meiyu season using laser disdrometer observed RSD data from 2016 to 2019. For the average spectra of five rain rate classes, the concentrations of large raindrops (>0.5 mm) increased with rain rate (R), while the concentrations of small raindrops (<0.5 mm) increased only under rain rates higher than 10 mm/h. The gamma distribution parameters of N_0 (intercept parameter) and Λ (slope parameter) increased/decreased with rain rate, and the shape parameter μ exhibited negative values in different rain rate classes. The distribution pattern features were $N(D) = 721D^{-1.79}e^{-1.20D}$. Distributions of the frequency for mass-weighted mean diameter (D_m) and the logarithm of the generalized intercept parameter ($\log_{10}N_w$) both showed a unique bimodal type, and an exceptionally high N_w ($\log_{10}N_w > 4.5$) subset with small D_m was determined. The stratiform and convective rain of RSD were also investigated. $D_m - R$ and $N_w - R$ showed similar variations in two types of precipitation. The lower μ values resulted in higher primary and constant coefficients in the quadratic polynomial fitting for the $\mu - \Lambda$ relationship ($\Lambda = 0.0347\mu^2 + 1.180\mu + 2.495$). The $Z - R$ relationship (Z for radar reflectivity factor) in stratiform precipitation characteristics was $Z = 203R^{1.59}$. Further investigations showed that high N_w values usually occurred in persistent precipitation. The RSD can be characterized as high concentrations of the first two diameter classes with narrow spectrum width (<1 mm), which were captured during in-cloud rain with

60 a low but continuous rain rate (<5 mm/h). The mountainous topography plays an important
61 role in reshaping the characteristics of RSD and the physical processes of precipitation.

62

63 **Key Words** Meiyu Season; Mount Lushan; Raindrop size distribution; Stratiform and
64 convective rain

65

66 1. Introduction

67 More accurate precipitation prediction and related physical mechanisms researches are
68 vital for meteorology, climatology, and hydrology. Investigation of the Raindrop size
69 distribution (RSD) is essential for exploring the characteristics and underlying physical
70 precipitation processes (Rosenfeld and Ulbrich, 2003). The characteristics of RSD from
71 ground-based observations are synthetic, and address the microphysics and thermal
72 dynamics within, or outside, precipitation clouds (Rosenfeld and Ulbrich, 2003; Thomas et
73 al., 2021). The parameterization of RSD is fundamental to improving the microphysics of
74 clouds and precipitation in numerical model simulations (Haddad et al., 1996; Milbrandt and
75 Yau, 2005; Rotsteyn, 1997; Sun, 2005; Tapiador et al., 2014; Tapiador et al., 2019). RSD
76 researches have contributed to improving quantitative precipitation estimations for weather
77 radar and satellite observations (Atlas et al., 1999; Chandrasekar and Bringi, 1987;
78 Chandrasekar et al., 2003; Rosenfeld and Ulbrich, 2003; Seto et al., 2021; Tokay and Short,
79 1996; Ulbrich and Atlas, 1998; Zhang et al., 2001; Zhao et al., 2011).

80 The characteristics of RSD vary according to geographical locations (Chang et al., 2009;
81 Chen et al., 2017a; Dolan et al., 2018; Luo et al., 2020; Niu et al., 2010; Radhakrishna et al.,
82 2020; Sauvageot and Lacaux, 1995; Zhao et al., 2011), climate regimes (Bringi et al., 2003;
83 Yamaji et al., 2020), seasons (Chakravarty and Raj, 2013; Seela et al., 2018; Wen et al.,
84 2017), and rain types (Atlas and Ulbrich, 2006; Chen et al., 2016; Chen et al., 2013;
85 Thompson et al., 2015; Tokay and Short, 1996). RSD characteristics and their variations
86 may differ considerably between different disdrometer instruments (Angulo-Martínez et al.,
87 2018), or even between the same instrument but within a short distance (Tapiador et al.,
88 2010). Systematic RSD studies have been performed over the past two decades to better

understand the characteristics of RSD in the East Asian monsoon region. Statistical studies of RSD, ranging from northern China to the Tibetan Plateau have been conducted, and the RSD characteristics significantly vary in different regions. Physical processes between stratiform and convective rains also exhibit obvious differences in different climate regimes (Chen et al., 2017a; Chen et al., 2013; Ji et al., 2019; Luo et al., 2020; Niu et al., 2010; Wen et al., 2016; Zhang et al., 2019).

Meiyu is a climatological phenomenon that affects the weather and climate in eastern Asia and usually induces abundant precipitation (Ding, 1994). The Meiyu/Baiu front is characterized as a quasi-stationary front along the north edge of a subtropical high stretching from the Yangtze River basin to south Japan and presents unique weather systems during the Meiyu season. Previous research has shown that heavy precipitation events and related flood disasters are likely caused by the Meiyu/Baiu front (Tao and Ding, 1981; Xu et al., 2008; Yasunari and Miwa, 2006; Zhao et al., 2019). In addition, it can influence extreme precipitation in Mount Lushan by adjusting the low-level jet (Li et al., 2012). Hashimoto and Harimaya (2003) analyzed the Baiu RSD data obtained in south Japan using principal component analysis and found two principal components of variation of RSD, which could be attributed to the difference between convective and stratiform precipitation. Different characteristics of RSD variation were obtained in their subsequent study, the major precipitation mechanism in the mature and dissipation stages of the Meiyu/Baiu convective clouds (Hashimoto and Harimaya, 2005). By employing a C-band polarimetric radar and a disdrometer simultaneously, Oue et al. (2010) and Oue et al. (2011) found and characterized different RSD with low echo-top and large Z_h (radar reflectivity) in stratiform and convective rain zones within the Meiyu/Baiu frontal rainband in the Okinawa region. Chen et al. (2013)

112 found that the mass weighted diameter (D_m) and the generalized intercept parameter N_w
113 in eastern China were lower than those reported in Japan (Bringi et al., 2003). Wen et al.
114 (2016) studied the RSD characteristics and differences in convective, stratiform, and shallow
115 rains, and higher N_w , and lower D_m values than those of Chen et al. (2013) were obtained.
116 More recently, Fu et al. (2020) studied RSD characteristics in central China, and
117 demonstrated different N_w and D_m values and different $\mu - \Lambda$ and $Z - R$ relationships.
118 Although such studies have furthered the understanding of RSD in precipitation processes,
119 few have focused on mountainous regions, and even less attention has been given to the
120 impacts of mountainous topography on RSD characteristics and the related physical
121 processes.

122 More RSD observations and studies are needed to understand the physical processes
123 within the Meiyu precipitation concerning the high variety of RSD characteristics among
124 different geographical locations. The Lushan Mountains, with most peaks being more than
125 1000 m, are known as typical horst block mountains. They are in the northern Jiangxi
126 Province, which faces Poyang Lake to the east and borders the Yangtze River in the north
127 (Fig. 1a). The high mountainous topography of Mount Lushan can act as an elevated
128 platform to investigate the interactions between aerosols, clouds, and precipitation because
129 it is close to the cloud base, or within the cloud directly (Duan et al., 2021; Guo et al., 2019).
130 Continuous field observations of cloud and fog processes in Mount Lushan have been
131 conducted since 2015 at the Lushan Cloud and Fog Experiment Station, and in this study,
132 a laser precipitation monitor disdrometer was employed to observe the RSD characteristics.
133 The long-term observed RSD dataset in Mount Lushan provides opportunities for
134 investigating more comprehensive RSD characteristics and related physical processes

135 under mountainous topographical conditions.

136 The aim of this study is to explore the characteristics of RSDs during the Meiyu season
137 in Mount Lushan from 2016 to 2019. This study is structured as follows: The data and
138 methods are described in Section 2. The characteristics of RSDs for different rain rates and
139 rain types, and the relationships between RSD parameters and gamma fitting parameters
140 are presented in Section 3. Section 4 discusses the dataset with specific properties, and the
141 conclusions are summarized in Section 5.

142 2. Data and Methods

143 2.1 Observations

144 A Laser Precipitation Monitor (LPM) from Thies CLIMA was set at the observational
145 platform of the Lushan Cloud and Fog Experiment Station in Mount Lushan (22.57°N,
146 115.97°E, 1080 m ASL) (Fig. 1). The LPM is a ground-based optical disdrometer with a
147 sampling area of 45.6 cm² and a temporal resolution of 1 min. The RSD data were collected
148 every 1 min in a 22 × 20 matrix that recorded the raindrop numbers of each diameter and
149 velocity bin. The diameter ranged from 0.125 mm to larger than 8 mm, and the velocity
150 ranged from 0 m/s to 20 m/s.

151 2.2 Methods

152 a. LPM data quality control

153 The effective sampling area for laser optical sensors is affected by the border effects
154 (Battaglia et al., 2010). In the i th size class, the effective sampling area is
155 $228 \times (20 - D_i) \text{ mm}^2$ and D_i (mm), which is the mean diameter for the i th size diameter
156 class (Jaffrain and Berne, 2011). The incorrect records of small particles with high falling
157 speed are named as the “margin fallers” and detected when they partially cross the sampling

158 area (Yuter et al., 2006). Strong winds and splashing usually cause spurious particle
 159 recording, and according to previous studies, the empirical terminal velocity-diameter (V-D)
 160 relationship is constantly used to assess and eliminate spurious records (Chen et al., 2017a;
 161 Chen et al., 2013; Ji et al., 2019). In this study, we chose the empirical V-D relationship with
 162 the air density correction factor to assess the raindrop data (Atlas et al., 1973):

$$V(D) = (9.65 - 10.3e^{-0.6D}) \left(\frac{\rho_0}{\rho_a} \right)^{0.4}, \quad (1)$$

164 where ρ_0 and ρ_a (1.20 kg/m^3) are the air density at the observation level and sea level,
 165 respectively. The temperature and pressure data of the automatic meteorological station
 166 (AWS) of the Lushan Meteorological Bureau (29.57°N , 115.97°E , 1165 m) were used to
 167 retrieve the correction factor. Using the AWS data in June and the first half of July from 2016
 168 to 2020, the average pressure and temperature during the Meiyu season were first
 169 calculated, and ρ_0/ρ_a was then derived through the ideal gas state equation ($P = \rho RT$).
 170 Finally, a correction factor of 1.0655 was obtained. The terminal velocity $V(D_i)$ of each
 171 diameter bin D_i was achieved, and the particles out of the velocity range $V(D_i) \pm 0.6V(D_i)$
 172 were removed (Friedrich et al., 2013; Jaffrain and Berne, 2011).

173 *b. Raindrop size distribution parameters*

174 The raindrop size distribution is depicted as the number concentration of raindrops
 175 $N(D_i)$; $\text{m}^{-3} \text{ mm}^{-1}$) per unit volume from D_i to $D_i + \Delta D_i$ in size class i , and can be calculated
 176 from Equation (2):

$$N(D_i) = \sum_j^{20} \frac{n_{ij}}{A_i \cdot \Delta t \cdot V_j \cdot \Delta D_i}, \quad (2)$$

178 where n_{ij} is the number of raindrops for the i th diameter bin and j th velocity bin, A_i (m^2) is
 179 the effective sampling area, Δt is the sampling time interval (60 s for LPM), V_j (m/s) is the

180 average velocity of the j th velocity bin, and ΔD_i (mm) is the width of each diameter bin.

181 The bulk microphysical parameters total number concentration N_T (m^{-3}), rainwater
182 content W (g/m^3), rain rate R (mm/h), and radar reflectivity factor Z (mm^6m^{-3}) were
183 calculated as follows:

$$184 \quad N_T = \sum_{i=1}^{22} \sum_{j=1}^{20} \frac{n_{ij}}{A_i \cdot \Delta t \cdot V_j}, \quad (3)$$

$$185 \quad W = \frac{\pi}{6} \times 10^{-3} \cdot \rho_w \cdot \sum_{i=1}^{22} \sum_{j=1}^{20} D_i^3 \cdot \frac{n_{ij}}{A_i \cdot \Delta t \cdot V_j}, \quad (4)$$

$$186 \quad R = 6\pi \times 10^{-4} \cdot \sum_{i=1}^{22} \sum_{j=1}^{20} D_i^3 \cdot \frac{n_{ij}}{A_i \cdot \Delta t}, \quad (5)$$

$$187 \quad Z = \sum_{i=1}^{22} \sum_{j=1}^{20} D_i^6 \cdot \frac{n_{ij}}{A_i \cdot \Delta t \cdot V_j}, \quad (6)$$

188 where ρ_w is the density of water ($1 \text{ g}/\text{cm}^3$). The mass-weighted mean diameter D_m (mm)
189 and generalized intercept parameter N_w ($\text{m}^{-3} \text{ mm}^{-1}$) were introduced to describe the bulk
190 characteristics of RSD, which could be calculated according to Equations (7) and (8).

$$191 \quad D_m = \frac{\sum_{i=1}^{22} N(D_i) \cdot D_i^4 \cdot \Delta D_i}{\sum_{i=1}^{22} N(D_i) \cdot D_i^3 \cdot \Delta D_i}, \quad (7)$$

$$192 \quad N_w = \frac{4^4}{\pi \rho_w} \cdot \left(\frac{10^3 \cdot W}{D_m^4} \right). \quad (8)$$

193 The gamma distribution proposed by Ulbrich (1983) was used to describe the distribution of
194 the raindrop size, as shown in Equation (9).

$$195 \quad N(D) = N_0 D^\mu e^{-\Lambda D}, \quad (9)$$

196 where D (mm) is the diameter, N_0 ($\text{m}^{-3}\text{mm}^{-1-\mu}$) is the intercept parameter, μ is the shape
197 parameter, and Λ (mm^{-1}) is the slope parameter.

198 In this study, a rain event was defined as when there were continuous spectra with no
199 less than 30 min, and two events with a less than 1 h rain-free period would be regarded as

one (Chen et al., 2017a; Tokay and Bashor, 2010). The spectra with a rain rate of less than 0.1 mm/h or a total number of less than 10 would be treated as noise (Chen et al., 2013). During the Meiyu period from 2016 to 2019, a dataset of 186 rain events was derived from the quality-controlled observational data.

3. Results

3.1 RSD properties of different rain rates

The RSD of different rain rates have different characteristics (Chen et al., 2017a; Chen et al., 2013; Ji et al., 2019; Porcù et al., 2014). However, because of different geographic locations (e.g., Chen et al., 2017; Ji et al., 2019) and rain types (e.g., Porcù et al., 2014; Chen et al., 2013), the RSD properties of different rain rates might vary. In this study, the rain rates were divided into five classes: 1) $0.1 \leq R < 1$ mm/h, 2) $1 \leq R < 5$ mm/h, 3) $5 \leq R < 10$ mm/h, 4) $10 \leq R < 100$ mm/h, and 5) $R \geq 100$ mm/h. Samples with $R < 5$ mm/h contributed 79% to the total sample numbers and 22.74% to the precipitation amount, and the samples with $5 \leq R < 10$ mm/h and $R \geq 10$ mm/h were 10.25% and 14.72%, 10.74%, and 62.54%, respectively (Fig. 3). The percentages of sample counts and precipitation amounts were similar to the results for Nanjing, a city with a lower altitude in east China (Chen et al., 2013), however, there were more samples (0.32%, 135 samples) with higher rain rates in Mount Lushan.

For large raindrops (>0.5 mm), the number concentration increases with the rain rate, and the spectra rise in parallel. For small raindrops (<0.5 mm), when the rain rates are lower than 10 mm/h, the concentrations for smaller/larger than 0.0375 mm raindrops will decrease/increase, which concurs with the 'rotational movement' reported by Hashimoto and Harimaya (2003). When the rain rates exceeded 10 mm/h, the spectra tended to expand

223 in parallel, and the concentrations of all diameter bins will rise with the rain rate. The average
224 spectrum is like that of $5 \leq R < 10$ mm/h, but with a higher concentration at the smaller and
225 larger end of the diameter range, and a lower concentration at the middle diameter range
226 (Fig. 4).

227 To compare the spectral characteristics for different rain rate classes, the standard
228 deviation of the mass spectrum $\sigma_M = D_m / (4 + \mu)^{1/2}$ and the normalized standard deviation
229 of the mass spectrum σ_M / D_m , which were defined by Ulbrich and Atlas (1998), were used.
230 To obtain the μ parameter from the RSD, the fitting methods were first compared. The
231 maximum likelihood (ML) (Brawn and Upton, 2007; Brawn and Upton, 2008) and truncated
232 moments (MM) (Ulbrich and Atlas, 1998; Zhang et al., 2003) methods were used to estimate
233 the parameters of the gamma distribution. The MM method fits better than the ML method
234 in terms of RSD shape. For rain rates of < 10 mm/h, the ML method
235 underestimates/overestimates the concentrations for large/small particles, and the ML
236 method also slightly overestimates the larger diameter classes (diameter > 0.5 mm) for rain
237 rates ≥ 10 mm/h. For the entire dataset average spectrum, the ML method overestimates the
238 concentrations in the diameter range of $0.5\text{--}3$ mm and the smallest diameter class (0.125--
239 0.25 mm), and underestimates the concentrations in the $0.25\text{--}0.5$ mm and > 3.5 mm ranges
240 (Fig. 5).

241 Table 1 presents the correlation coefficient, absolute bias, relative bias, N_T , and W for
242 the observation and fitting results for the two fitting methods. The absolute bias and relative
243 bias were calculated following Tokay and Bashor (2010). The ML method shows significant
244 advantages over the MM method according to the correlation coefficients and absolute
245 biases, and is similar to observed results in the Tibetan Plateau (Chen et al., 2017b).

246 However, the MM method shows superior results in terms of relative biases. For N_T , ML
 247 obtains better results for smaller rain rate classes (<10 mm/h), and MM obtains better results
 248 for larger rain rate classes (≥ 10 mm/h). For the entire dataset, the two methods exhibit
 249 similar results. For W , the MM method shows superior results to the ML method. The results
 250 will be based on the MM method to further discuss the characteristics of Meiyu precipitation,
 251 because it has advantages in terms of describing the RSD shape and estimating the
 252 precipitation.

253 Table 2 presents parameters and fitting results of the average spectra for different rain
 254 rates. According to Table 1, the N_T and W both increase with rain rate, and in Table 2,
 255 the Z , D_m , and σ_M exhibit similar variations, meaning that the rising parameters have close
 256 relationships with diameter, spectra width, and rain rate. In contrast, $\log_{10}N_w$ and σ_M/D_m ,
 257 which address concentration, diameter and spectra shape, scarcely vary with rain rate. For
 258 the fitting results, N_0 and Λ increase/decrease with rain rate, and μ varies little with rain rate
 259 and are negative for all rain rate classes. This differs from other observations (Chen et al.,
 260 2017a; Chen et al., 2016; Chen et al., 2013). The concentrations for small raindrops are
 261 slightly high, and as a result, lead to a left lower bending trend shape of the spectra, and
 262 negative μ values are derived (Figs. 4 and 5).

263 3.2 Differences between stratiform and convective rains in Meiyu

264 a. Classification of stratiform and convective rains

265 It has been widely demonstrated that stratiform rains have quite different physical
 266 processes compared to convective rains (Bringi et al., 2003; Chang et al., 2009; Chen et al.,
 267 2017a; Chen et al., 2013; Ji et al., 2019; Zhang et al., 2019), as well as microphysical
 268 schemes and retrieval methods in remote sensing (Atlas and Ulbrich, 2006; Chen et al.,

2017a; Dolan et al., 2018; Rosenfeld and Ulbrich, 2003; Ulbrich and Atlas, 1998). To determine the differences between stratiform and convective rains, two classification schemes were considered, namely, the T01 scheme followed by Testud et al. (2001), and B03 scheme (Bringi et al., 2003). For a rain rate time series $\{R_i\}$, in T01 scheme, if R_k and the adjacent 10 spectra ($R_{k-5} \sim R_{k+5}$) are all less than 10 mm/h, then the spectrum k will be classified as stratiform, otherwise the spectrum k will be classified as convective. As to the B03 scheme, if the standard deviation (σ_R) of 10 consecutive spectra is no larger than 1.5 mm/h, then the 10 consecutive spectra will be classified as stratiform, while when the σ_R is larger than 1.5 mm/h and the rain rate is no smaller than 5 mm/h, the spectrum will be classified as convective.

For the entire dataset (42,474 RSD samples), 33,518 (78.9%) samples were classified as stratiform and 8956 (21.1%) samples were classified as convective rain according to the T01 scheme. Using the B03 scheme, 30127 (70.9%) samples were classified as stratiform and 7866 (18.5%) samples were convective. T01 is a proper way to classify all samples to stratiform or convective rain, but the B03 scheme allows a higher rain rate to be classified as stratiform rain. The convective sample ratios for the stratiform samples were 26.1% and 26.7% for the T01 and B03 schemes, respectively, which were almost the same.

Figure 6 displays the distributions of D_m and $\log_{10}N_w$ frequency of stratiform and convective RSD samples. For the entire dataset, the frequency of D_m peaks at 0.5–0.6 mm and 1.0–1.1 mm and the mean value, standard deviation, and skewness for D_m are 1.20 mm, 0.57 mm, and 1.03, respectively. Compared with the results of Chen et al. (2013) and Bringi et al. (2006), the mean D_m is smaller, while the standard deviation and skewness are larger in Mount Lushan. For $\log_{10}N_w$, the peaks are 3.5–3.6 and 5.0–5.1, and the mean

value, standard deviation, and skewness are 3.88, 0.80, and 0.74, respectively. The mean $\log_{10}N_w$, which is close to the Marshall-Palmer value ($8000 \text{ m}^{-3}\text{m}^{-1}$ for N_w), is similar to Chen et al. (2013), but has a higher standard deviation and skewness. The bimodal distribution of D_m frequency was reported by Wen et al. (2016), and is caused by shallow rain. Furthermore, the frequency of $\log_{10}N_w$ exhibits a second peak. This was also reported and attributed to shallow rain by Wen et al. (2016) and Fu et al. (2020). In the present study, the elevation of Mount Lushan will result in insufficient coalescence and thus reduce the concentration of large raindrop. Also, it is easier for precipitating clouds to reach the ground because of the elevation, lack of evaporation in saturated environment will increase the concentration of small raindrop. So the second peak of $\log_{10}N_w$ in Mount Lushan can be affected jointly by shallow rain as well as the elevation of mountain topography. Moreover, the first peak in the present study differs from some Meiyu studies (Fu et al., 2020; Wen et al., 2016). The higher standard deviation and skewness also indicate significantly higher RSD variability s in Mount Lushan during the Meiyu season.

b. Distributions of D_m and $\log_{10}N_w$ in stratiform and convective rains

The stratiform-convective rain type classification methods of T01 and B03 were compared (Fig. 6), and the frequencies of D_m and $\log_{10}N_w$ for stratiform and convective rains of both methods exhibit similar results. The average value, standard deviation, and skewness of D_m of T01/B03 methods are 1.04/1.02 mm, 0.44/0.49 mm, 1.04/1.02, and 1.77/1.87 mm, 0.61/0.58 mm, 0.96/1.23 for stratiform and convective rains, respectively. For $\log_{10}N_w$, the average value, standard deviation, and skewness of T01/B03 methods are 3.91/3.93, 0.86/0.88, 0.70/0.70, and 3.76/3.77, 0.52/0.49, -0.24/-0.24 for stratiform and convective rains, respectively. There is no essential difference between the two methods,

315 however, the B03 scheme displays more distinct stratiform/convective characteristics.
316 Therefore, here, the B03 scheme is considered as the classification method.

317 The difference between the stratiform and convective rains mainly manifests in the
318 different peaks of the frequency distributions for D_m and $\log_{10}N_w$. However, different peaks
319 and skewness of frequency distributions for D_m and $\log_{10}N_w$ have been found among
320 different geographical locations (Bringi et al., 2006; Chen et al., 2013; Fu et al., 2020; Wen
321 et al., 2016) in both stratiform and convective rains. Larger standard deviations for D_m and
322 $\log_{10}N_w$ were obtained, indicating a more notable variety of RSD in Mount Lushan during
323 the Meiyu season because of the mountainous topography.

324 The distribution between D_m and $\log_{10}N_w$ exhibits characteristics similar to those of
325 other Meiyu RSD studies along the Yangtze River Valley (Fig. 7) (Chen et al., 2013; Fu et
326 al., 2020; Wen et al., 2016), which is generally more stratiform-like. Compared to other Meiyu
327 research, there are more samples with larger $\log_{10}N_w$ and lower D_m values in the Mount
328 Lushan dataset (Bringi et al., 2003; Chen et al., 2013; Fu et al., 2020; Wen et al., 2016). A
329 larger standard deviation of $\log_{10}N_w$ for stratiform rain and a smaller standard deviation of
330 D_m for convective rain were found in the present study, but the mean $\log_{10}N_w$ values were
331 quite close. Compared to other related Meiyu precipitation studies (Chen et al., 2013; Fu et
332 al., 2020; Wen et al., 2016), the present study obtained a smaller mean D_m for stratiform
333 rain but a larger mean D_m for convective rain, indicating that the variety of D_m is larger in
334 Mount Lushan. For the mean D_m - $\log_{10}N_w$ pairs of all data in each study, all studies are
335 similar and fall along the stratiform rain line proposed by Bringi et al. (2003) in the D_m -
336 $\log_{10}N_w$ coordinate. No significant tendency for maritime or continental characteristics can
337 be found in the Meiyu precipitation (Bringi et al., 2003; Bringi et al., 2009; Thompson et al.,

338 2015).

339 The relationship between $D_m - R$ and $N_w - R$ was investigated (Fig. 8). D_m
340 increases with rain rate for both stratiform and convective rains, and the power fitting results
341 show little difference between the two rain types, which is similar to other Meiyu observations
342 (Chen et al., 2013; Fu et al., 2020). The exponential fitting coefficients are higher than those
343 of Chen et al. (2013) and Wen et al. (2016), indicating that D_m increases faster with R at
344 higher rain rates. The N_w of stratiform rains shows little variation with rain rate, while for
345 convective rains, N_w shows a rising trend at lower rain rates. Compared to other studies,
346 the exponential fitting coefficients in the present study are quite low (Chen et al., 2013; Fu
347 et al., 2020; Wen et al., 2016), particularly for stratiform rains.

348 D_m and N_w both show bimodal type distributions (Fig. 8a, b), which correspond to the
349 frequency distributions of stratiform rains (Fig. 6). The high N_w peak (Fig. 8a) corresponds
350 to the low D_m peak (Fig. 8b). To investigate the unique high N_w samples, a subset,
351 consisting of samples with $\log_{10}N_w > 4.5$, was evaluated from stratiform rains for further
352 analysis (Fig. 8a, b). The exponential fitting coefficients of the subset between D_m and R
353 are the same as those of stratiform rain, but with a lower primary coefficient, and N_w shows
354 little variation with R .

355 The average spectra of the subset of $\log_{10}N_w > 4.5$, $\log_{10}N_w < 4.5$, and the entire
356 stratiform rain are presented in Fig. 9. The concentrations of smaller/larger (divided by 0.875
357 mm) for the subset with $\log_{10}N_w > 4.5$ are higher/lower than those of $\log_{10}N_w < 4.5$, and the
358 subset with $\log_{10}N_w > 4.5$, has a narrower spectrum width. Rosenfeld and Ulbrich (2003)
359 summarized the processes that can modify the shape of RSD, and stated that typically,
360 coalescence and evaporation cause a decrease in small raindrops. For the samples with

361 $N_w > 4.5$, the narrow spectrum width and high small raindrop concentrations are the result of
362 a lack of coalescence. When precipitation occurs within a cloud, the RSD spectra will always
363 be narrow and with high concentrations for small raindrops because of insufficient
364 coalescence and the absence of evaporation. This type of precipitation is frequently seen in
365 Mount Lushan, and therefore, it can be assumed that the high- N_w subset is the result of in-
366 cloud precipitation.

367 *c. RSD spectra*

368 The average spectrum of convective rain shows higher concentrations than stratiform
369 rain, while the differential concentrations rise with diameter (Fig. 10) (Table 3). Compared to
370 other studies conducted at lower altitudes along the Yangtze River Valley (Chen et al., 2013;
371 Fu et al., 2020), the concentrations of raindrops in Mount Lushan are higher at the small
372 diameter end than particle size velocity (Parsivel) measurements, and this can be attributed
373 to raw particle size and fall velocity distribution differences estimated by two sensors
374 (Angulo-Martínez et al., 2018, see in Table 4). Furthermore, compared with Chen et al. (2013)
375 and Wen et al. (2016), higher raindrop concentrations at the large-diameter end were also
376 obtained in Mount Lushan, and may be related to the mountainous topography.

377 The gamma distribution fits quite well for both rain types, except for the underestimation
378 of the first diameter bin class (Fig. 10). The N_0 and μ for stratiform rains are higher than
379 convective rains, and the Λ for stratiform rains is smaller than convective rains. Compared
380 with Chen et al. (2013), N_0 and Λ are significantly lower in Mount Lushan, and μ is
381 negative for both rain types, indicating that the spectra are bent toward the left bottom. The
382 differences between the present study and those of Chen et al. (2013) can be attributed to
383 the high concentration of small raindrops resulting from the mountainous topography of

384 Mount Lushan.

385 3.3 Relationships of gamma distribution parameters

386 Theoretical calculations and observational results both show that the parameters of
387 gamma distribution are not independent (Atlas et al., 1973; Chandrasekar and Bringi, 1987;
388 Chen et al., 2017a; Haddad et al., 1996; Seela et al., 2018; Ulbrich, 1983; Zhang et al.,
389 2003). To investigate the relationships between the gamma distribution parameters and the
390 distinctiveness of RSD in Mount Lushan, the relationships between $\mu - N_0$ and $\mu - \Lambda$ were
391 analyzed (Fig. 11).

392 It has been found theoretically and empirically that N_0 and μ have an exponential
393 relationship of $N_0 = C_N \exp(3.2\mu)$, where C_N is a constant (Ulbrich, 1983). The distributions
394 of the two parameters show an obvious exponential relationship, and the fitting equations
395 for stratiform rain, convective rain, and the entire dataset are $N_0 = 3720 \exp(1.70\mu)$, $N_0 =$
396 $7406 \exp(1.29\mu)$, $N_0 = 13043 \exp(1.45\mu)$, respectively (Fig. 11). Compared to Ulbrich (1983),
397 the present study shows a lower exponential coefficient, but the C_N here is of the same
398 order of magnitude. However, when compared with the dataset on the Tibetan Plateau
399 (Chen et al., 2017a), the exponential coefficient is similar, but a higher C_N was found in
400 Mount Lushan. The differences between the three studies indicate that although $\mu - N_0$
401 has an exponential relationship, the fitting results vary significantly among different regions
402 and rain types, and it is inconvenient to assess the varieties because N_0 will change with
403 μ (Chen et al., 2017a).

404 Zhang et al. (2001) and Zhang et al. (2003) proposed a quadratic polynomial $\mu - \Lambda$
405 relationship ($\Lambda = 0.0365\mu^2 + 0.735\mu + 1.935$), which is thought to be a fundamental property
406 of RSD, especially in convective rain (Seifert, 2005). The $\mu - \Lambda$ relationship varies

407 according to geographic location, climate regime, and synoptic system (Bringi et al., 2003;
 408 Chen et al., 2011; Chen et al., 2017a; Chen et al., 2016; Chen et al., 2013; Ji et al., 2019;
 409 Luo et al., 2020). The $\mu - \Lambda$ relationship for stratiform rain, convective rain, and the entire
 410 dataset of this study are presented in Fig. 11b. The fitting equations for three datasets are
 411 $\Lambda = 0.0211\mu^2 + 1.365\mu + 2.374$, $\Lambda = 0.0533\mu^2 + 0.974\mu + 2.281$, and $\Lambda = 0.0347\mu^2 +$
 412 $1.180\mu + 2.495$, respectively. The $\mu - \Lambda$ of stratiform rain is similar to the results in the
 413 Tibetan Plateau, which is $\Lambda = 0.0217\mu^2 + 1.090\mu + 1.706$, but with a higher constant
 414 coefficient (Chen et al., 2017a). The $\mu - \Lambda$ of convective rain in Mount Lushan is similar to
 415 the result of a squall line case in Eastern China, which is $\Lambda = 0.0585\mu^2 + 0.812\mu + 1.934$,
 416 also with a higher constant coefficient (Chen et al., 2016). Compared with the results of
 417 Zhang et al. (2003), the quadratic coefficient for the entire dataset here is similar, but the
 418 primary and constant coefficients are higher. However, in Chen et al. (2013), which is $\Lambda =$
 419 $0.0149\mu^2 + 0.491\mu + 2.015$, all coefficients differ from Mount Lushan. The present study
 420 shows that under conditions using different instruments, the similarities of the $\mu - \Lambda$
 421 relationship can still be assessed for different geophysical locations, climate regimes, and
 422 synoptic systems.

423 3.4 $Z - R$ relationship

424 The $Z - R$ relationship, in the form of $Z = AR^b$ is an important criterion in radar
 425 quantitative precipitation estimation (QPE) (Chen et al., 2013; Porcù et al., 2014; Tokay and
 426 Short, 1996), and here, the $Z - R$ relationship in Mount Lushan was investigated (Fig. 12).
 427 The $Z - R$ relationship for the entire dataset is $Z = 203R^{1.59}$ (Fig. 12a), which is similar to
 428 the continental stratiform rain $Z = 200R^{1.6}$ (Marshall and Palmer, 1948), and indicates that
 429 the Meiyu precipitation in Mount Lushan is typically stratiform-like. The $Z - R$ relationships

for stratiform and convective rains are $Z = 194R^{1.53}$ and $Z = 198R^{1.63}$, respectively, which are also similar to those of stratiform rain, although there are minute differences (Marshall and Palmer, 1948). When comparing the operational WSR-88D $Z = 300R^{1.4}$ (Fulton et al., 1998) and Meiyu results in eastern China $Z = 368R^{1.21}$ (Chen et al., 2013), the rain rate R derived from Z will be overestimated/underestimated at lower/higher rain rate ranges (dividing rain rate approximately 5–10 mm/h) in Mount Lushan. The Meiyu precipitation in Mount Lushan can be generally characterized as typical stratiform rain.

The samples with $\log_{10}N_w > 4.5$ show significantly different characteristics, and therefore, the differences between samples with $\log_{10}N_w > 4.5$ and $\log_{10}N_w < 4.5$ were investigated. The $Z - R$ scatterplots show a clear boundary between samples with $\log_{10}N_w > 4.5$ and $\log_{10}N_w < 4.5$, for the entire dataset (Fig. 12a) and the stratiform dataset (Fig. 12b). The $Z - R$ relationships of samples with $\log_{10}N_w > 4.5$ for the entire dataset and stratiform rain dataset are $Z = 54R^{1.47}$ and $Z = 51R^{1.41}$, respectively, showing little difference. The characteristics of the spectra and $Z - R$ relationship for the high N_w samples are similar to the shallow rains diagnosed by Wen et al. (2016), and was attributed to warm rain processes. For those with $\log_{10}N_w < 4.5$, the $Z - R$ relationship for the entire dataset is $Z = 306R^{1.51}$, which is similar to that of WSR-88D (Fulton et al., 1998), but with a higher b value. The $Z - R$ relationship for stratiform rain samples with $\log_{10}N_w < 4.5$ is $Z = 203R^{1.59}$, which is very similar to continental stratiform rain (Marshall and Palmer, 1948). The $Z - R$ relationship for the high- N_w subset in the present study shows a higher rain rate at the same Z value. Yuter and Houze (1997) reported a bimodally distributed $Z - R$ relationship in the Pacific Warm Pool, which corresponds to the populations of large- and small-drop spectra. The present study also revealed a similar bimodal distribution, but with

453 a significantly lower rain rate. The underlying reason should be further investigated.

454 **4. Discussion**

455 In Section 3, a typical dataset with high N_w ($\log_{10}N_w > 4.5$) was assessed and showed
456 quite different RSD characteristics. To investigate the characteristics and underlying reasons
457 for the high N_w samples, a primary statistical analysis of the high N_w subset was
458 conducted. About one-third (60) of the rain events had high N_w samples with counts of more
459 than 30. A further investigation of two long-lasting precipitation events of Meiyu found that
460 high N_w samples always existed within continuous long-lasting precipitation processes (Fig.
461 13) that both lasted for more than one day, with few interruptions. The high N_w samples
462 were normally obtained during the interval of two relatively heavy rain periods or in the
463 dissipating period of the long-lasting precipitation (e.g., Fig. 13a, b). The samples with high
464 N_w can be characterized as: 1) high concentrations in the first two diameter classes, 2)
465 narrow spectrum width (no larger than 1 mm), and 3) low but continuous rain rate (mostly
466 lower than 5 mm/h).

467 Mount Lushan is well known for its changeable clouds and fogs, and it is believed that
468 in-cloud precipitation is frequently observed on the observational platform (Fig. 14). When a
469 rain event occurs within the cloud in Mount Lushan, the absence of evaporation will cause
470 a high concentration of small raindrops, and the low clouds will also lack sufficient
471 coalescence. As such, a narrow spectrum with a high small raindrop concentration will be
472 expected from in-cloud rain. To confirm this, a rain case was observed by the LPM, fog
473 monitor (DMT FM-100), and visibility meter (Vaisala PWD22) (Fig. 15). Under similar rain
474 rates, the N_w/D_m outside the cloud (before 09:00 on May 23) was lower/higher than that
475 inside the cloud (15:00–21:00 on May 23). When the rain began to dissipate, the N_w/D_m

476 became higher/lower. Wen et al. (2016) reported the shallow rain type during the Meiyu
477 season, which is like the high N_w subset and is regarded as forming through warm rain
478 processes. However, in the present study, a more highlighted high N_w subset can be
479 derived from the entire dataset (e.g., Figs. 6 and 12), indicating that in-cloud rain, which is
480 highly related to mountainous topography, plays an important role in generating a high- N_w
481 RSD.

482 Although the topography of Mount Lushan is critical in generating in-cloud precipitation,
483 to explain why the high- N_w samples always occur in long-lasting precipitation, the
484 underlying dynamics and thermodynamics need to be investigated in depth. Furthermore,
485 the microphysical mechanisms of conversion from cloud/fog droplets to raindrops, and the
486 interaction between cloud/fog and precipitation, also require investigation.

487 **5. Conclusions**

488 Using the RSD data collected by LPM in Mount Lushan from 2016 to 2020, the RSD
489 characteristics of Meiyu precipitation in Mount Lushan were studied. The special RSD
490 samples with high N_w were also discussed, and the results are as follows:

491 A dataset of 186 rain events was determined during the Meiyu season from 2016 to
492 2019 in Mount Lushan. For the entire dataset, lower rain rates (<5 mm/h) contributed to the
493 highest number of samples, while rain rates of 10–100 mm/h contributed the highest
494 precipitation amount. Extraordinarily high rain rates (>100 mm/h) were found in Mount
495 Lushan during the Meiyu season, and contributed to quantitative precipitation (8.62%)
496 compared with its low frequency (0.32%). For large diameter ranges (>0.5 mm), the
497 concentrations for each diameter bin of the average spectra will increase with rain rate, while
498 for small diameter ranges (<0.5 mm), the concentrations for each diameter bin will rise only

499 when the rain rates are higher (>10 mm/h).

500 The performance of the two gamma fitting methods, namely, the maximum likelihood
501 (ML) and the truncated moments (MM), were evaluated using the average spectra of each
502 rain rate class for the Meiyu dataset. The ML method obtained better results for correlation
503 coefficients and absolute biases, and the MM method obtained better results for relative
504 biases and shapes of RSD. In simulating the RSD parameters, the MM method had good
505 results for N_T and W . Therefore, the MM method achieved good results for simulating the
506 Meiyu RSD in Mount Lushan.

507 The RSD parameters of N_T , W , Z , D_m , and σ_M increased with rain rate, while N_w
508 and σ_M/D_m varied little with rain rate. For the gamma distribution parameters of the
509 average spectra for different rain rates, N_0 and Λ rose/decreased with rain rate, and μ had
510 negative values for all rain rate classes. The gamma distribution for the average spectrum
511 of the entire dataset was $N(D) = 721D^{-1.79}e^{-1.20D}$. The frequency distributions of D_m and
512 $\log_{10}N_w$ both showed a special bimodal type. An exceptionally high- N_w ($\log_{10}N_w > 4.5$)
513 subset was diagnosed, which corresponds to the unique secondary peaks of D_m and
514 $\log_{10}N_w$, which has a narrower and steeper average spectrum with high concentrations for
515 small raindrops (<0.875 mm).

516 Two classification schemes (T01 and B03) for stratiform/convective rain were discussed,
517 and the frequency distributions of D_m and $\log_{10}N_w$ for stratiform and convective rains
518 showed little difference between the two schemes. The distribution of D_m - $\log_{10}N_w$ showed
519 differences between the two rain types and no significant maritime/continental
520 characteristics. The $D_m - R$ and $N_w - R$ relationships showed little difference between
521 stratiform and convective rains, and N_w showed little variation with rain rate. The average

spectra for stratiform and convective rains both had high concentrations of small raindrops, but the convective rains had larger raindrops. The gamma distribution parameters were also calculated (Table 3).

The relationships between the gamma distribution parameters of $\mu - N_0$ and $\mu - \Lambda$ for stratiform and convective rain were examined. The $\mu - N_0$ showed quite different characteristics for stratiform and convective rains, and the fitting equation for the entire dataset was $N_0 = 13043 \exp(1.45\mu)$. The Meiyu dataset of Mount Lushan showed a similar $\mu - \Lambda$ relationship ($\Lambda = 0.0347\mu^2 + 1.180\mu + 2.495$) to other studies, but in the present study, the lower μ values resulted in higher primary and constant coefficients in the quadratic polynomial fitting results.

The $Z - R$ relationship for the Meiyu season in Mount Lushan showed featured stratiform rain characteristics with a $Z = 203R^{1.59}$ fitting result. The $Z - R$ relationships for stratiform and convective rains showed little difference, but the dataset of high N_w samples obtained a unique fitting result of $Z = 54R^{1.47}$.

The high- N_w subset was normally detected in long-lasting precipitation during the Meiyu season. The RSD can be characterized as high concentrations in the first two diameter classes, with narrow spectrum width (<1 mm), and low but continuous rain rate (<5 mm/h). Further investigations demonstrated that these samples can be attributed to in-cloud precipitation, which is related to the mountainous topography of Mount Lushan.

541

Acknowledgments

This work was jointly supported by Chinese Academy of Meteorological Sciences Basic Research and Operation Fund (2019Z013, 2016Z004, 2020Y010), S&T Development Fund

545 of CAMS (2021KJ017), and the National Key R&D Program of China (2019YFC1510300).

546

547

References

548 Angulo-Martínez, M., Beguería, S., Latorre, B., Fernández-Raga, M., 2018. Comparison of precipitation
549 measurements by OTT Parsivel2 and Thies LPM optical disdrometers. *Hydrol. Earth Syst. Sci.*, 22(5):
550 2811-2837. DOI:10.5194/hess-22-2811-2018

551 Atlas, D., Srivastava, R.C., Sekhon, R.S., 1973. Doppler radar characteristics of precipitation at vertical
552 incidence. *Reviews of Geophysics*, 11(1): 1-35. DOI:10.1029/RG011i001p00001

553 Atlas, D., Ulbrich, C., 2006. Drop size spectra and integral remote sensing parameters in the transition
554 from convective to stratiform rain. *Geophysical Research Letters*, 33(16).
555 DOI:https://doi.org/10.1029/2006GL026824

556 Atlas, D., Ulbrich, C.W., Marks Jr., F.D., Amitai, E., Williams, C.R., 1999. Systematic variation of drop size
557 and radar-rainfall relations. *Journal of Geophysical Research: Atmospheres*, 104(D6): 6155-6169.
558 DOI:https://doi.org/10.1029/1998JD200098

559 Battaglia, A., Rustemeier, E., Tokay, A., Blahak, U., Simmer, C., 2010. PARSIVEL Snow Observations: A
560 Critical Assessment. *Journal of Atmospheric and Oceanic Technology*, 27(2): 333-344.
561 DOI:10.1175/2009JTECHA1332.1

562 Brawn, D., Upton, G., 2007. Closed-form parameter estimates for a truncated gamma distribution.
563 *Environmetrics*, 18(6): 633-645. DOI:https://doi.org/10.1002/env.841

564 Brawn, D., Upton, G., 2008. Estimation of an atmospheric gamma drop size distribution using disdrometer
565 data. *Atmospheric Research*, 87(1): 66-79. DOI:https://doi.org/10.1016/j.atmosres.2007.07.006

566 Bringi, V.N. et al., 2003. Raindrop Size Distribution in Different Climatic Regimes from Disdrometer and
567 Dual-Polarized Radar Analysis. *Journal of the Atmospheric Sciences*, 60(2): 354-365.
568 DOI:10.1175/1520-0469(2003)060<0354:Rsdidc>2.0.Co;2

569 Bringi, V.N. et al., 2006. Rainfall Estimation from C-Band Polarimetric Radar in Okinawa, Japan:
570 Comparisons with 2D-Video Disdrometer and 400 MHz Wind Profiler. *Journal of the Meteorological*
571 *Society of Japan. Ser. II*, 84(4): 705-724. DOI:10.2151/jmsj.84.705

572 Bringi, V.N., Williams, C.R., Thurai, M., May, P.T., 2009. Using Dual-Polarized Radar and Dual-Frequency
573 Profiler for DSD Characterization: A Case Study from Darwin, Australia. *Journal of Atmospheric and*

574 Oceanic Technology, 26(10): 2107-2122. DOI:10.1175/2009JTECHA1258.1

575 Chakravarty, K., Raj, P.E., 2013. Raindrop size distributions and their association with characteristics of
576 clouds and precipitation during monsoon and post-monsoon periods over a tropical Indian station.
577 Atmospheric Research, 124: 181-189. DOI:https://doi.org/10.1016/j.atmosres.2013.01.005

578 Chandrasekar, V., Bringi, V.N., 1987. Simulation of Radar Reflectivity and Surface Measurements of
579 Rainfall. Journal of Atmospheric and Oceanic Technology, 4(3): 464-478. DOI:10.1175/1520-
580 0426(1987)004<0464:SORRAS>2.0.CO;2

581 Chandrasekar, V., Meneghini, R., Zawadzki, I., 2003. Global and Local Precipitation Measurements by
582 Radar. In: Wakimoto, R.M., Srivastava, R. (Eds.), Radar and Atmospheric Science: A Collection of
583 Essays in Honor of David Atlas. American Meteorological Society, Boston, MA, pp. 215-236.
584 DOI:10.1007/978-1-878220-36-3_9

585 Chang, W.-Y., Wang, T.-C.C., Lin, P.-L., 2009. Characteristics of the Raindrop Size Distribution and Drop
586 Shape Relation in Typhoon Systems in the Western Pacific from the 2D Video Disdrometer and NCU
587 C-Band Polarimetric Radar. Journal of Atmospheric and Oceanic Technology, 26(10): 1973-1993.
588 DOI:10.1175/2009jtech1236.1

589 Chen, B., Hu, W., Pu, J., 2011. Characteristics of the raindrop size distribution for freezing precipitation
590 observed in southern China. Journal of Geophysical Research, 116: D06201.
591 DOI:doi:10.1029/2010JD015305

592 Chen, B., Hu, Z., Liu, L., Zhang, G., 2017a. Raindrop Size Distribution Measurements at 4,500 m on the
593 Tibetan Plateau During TIPEX-III. Journal of Geophysical Research: Atmospheres, 122(20): 11,092-
594 11,106. DOI:10.1002/2017JD027233

595 Chen, B., Wang, J., Gong, D., 2016. Raindrop Size Distribution in a Midlatitude Continental Squall Line
596 Measured by Thies Optical Disdrometers over East China. Journal of Applied Meteorology and
597 Climatology, 55(3): 621-634. DOI:10.1175/jamc-d-15-0127.1

598 Chen, B., Yang, J., Pu, J., 2013. Statistical Characteristics of Raindrop Size Distribution in the Meiyu
599 Season Observed in Eastern China. Journal of the Meteorological Society of Japan. Ser. II, 91(2): 215-
600 227. DOI:10.2151/jmsj.2013-208

601 Chen, J., Wu, X., Yin, Y., Huang, Q., Xiao, H., 2017b. Characteristics of Cloud Systems over the Tibetan
602 Plateau and East China during Boreal Summer. Journal of Climate, 30(9): 3117-3137.
603 DOI:10.1175/JCLI-D-16-0169.1

604 Ding, Y., 1994. Monsoons over China. *Advances in Atmospheric Sciences*, 11(2): 252-252.
605 DOI:10.1007/BF02666553

606 Dolan, B., Fuchs, B., Rutledge, S.A., Barnes, E.A., Thompson, E.J., 2018. Primary Modes of Global Drop
607 Size Distributions. *Journal of the Atmospheric Sciences*, 75(5): 1453-1476. DOI:10.1175/JAS-D-17-
608 0242.1

609 Duan, J. et al., 2021. Cable-car measurements of vertical aerosol profiles impacted by mountain-valley
610 breezes in Lushan Mountain, East China. *Science of The Total Environment*, 768: 144198.
611 DOI:10.1016/j.scitotenv.2020.144198

612 Friedrich, K., Higgins, S., Masters, F.J., Lopez, C.R., 2013. Articulating and Stationary PARSIVEL
613 Disdrometer Measurements in Conditions with Strong Winds and Heavy Rainfall. *Journal of*
614 *Atmospheric and Oceanic Technology*, 30(9): 2063-2080. DOI:10.1175/jtech-d-12-00254.1

615 Fu, Z. et al., 2020. Statistical Characteristics of Raindrop Size Distributions and Parameters in Central
616 China During the Meiyu Seasons. *Journal of Geophysical Research: Atmospheres*, 125(19):
617 e2019JD031954. DOI:https://doi.org/10.1029/2019JD031954

618 Fulton, R.A., Breidenbach, J.P., Seo, D.-J., Miller, D.A., O'Bannon, T., 1998. The WSR-88D Rainfall
619 Algorithm. *Weather and Forecasting*, 13(2): 377-395. DOI:10.1175/1520-
620 0434(1998)013<0377:Twra>2.0.Co;2

621 Guo, L. et al., 2019. An observational study of diurnal and seasonal variations, and macroscopic and
622 microphysical properties of clouds and precipitation over Mount Lu, Jiangxi, China (in Chinese). *Acta*
623 *Meteorologica Sinica*, 77(5): 923-937. DOI:doi:10.11676/qxxb2019.056

624 Haddad, Z.S., Durden, S.L., Im, E., 1996. Parameterizing the Raindrop Size Distribution. *Journal of*
625 *Applied Meteorology and Climatology*, 35(1): 3-13. DOI:10.1175/1520-
626 0450(1996)035<0003:PTRSD>2.0.CO;2

627 Hashimoto, A., Harimaya, T., 2003. Characteristics of the Variation of Raindrop Size Distribution in Baiu
628 Season. *Journal of the Meteorological Society of Japan. Ser. II*, 81(4): 871-878.
629 DOI:10.2151/jmsj.81.871

630 Hashimoto, A., Harimaya, T., 2005. Characteristics of Raindrop Size Distribution Dependent on the Life
631 Stage of a Convective Precipitation Cloud in the Baiu Season. *Journal of the Meteorological Society of*
632 *Japan. Ser. II*, 83(4): 641-649. DOI:10.2151/jmsj.83.641

633 Jaffrain, J., Berne, A., 2011. Experimental Quantification of the Sampling Uncertainty Associated with

634 Measurements from PARSIVEL Disdrometers. *Journal of Hydrometeorology*, 12(3): 352-370.
 635 DOI:10.1175/2010JHM1244.1

636 Ji, L. et al., 2019. Raindrop Size Distributions and Rain Characteristics Observed by a PARSIVEL
 637 Disdrometer in Beijing, Northern China. *Remote Sensing*, 11: 1479.

638 Li, Y. et al., 2012. Characteristics of precipitation chemistry at Lushan Mountain, East China: 1992–2009.
 639 *Environmental Science and Pollution Research*, 19(6): 2329-2343. DOI:10.1007/s11356-012-0742-2

640 Luo, L. et al., 2020. Raindrop size distribution and microphysical characteristics of a great rainstorm in
 641 2016 in Beijing, China. *Atmospheric Research*, 239: 104895.
 642 DOI:<https://doi.org/10.1016/j.atmosres.2020.104895>

643 Marshall, J., Palmer, W., 1948. The Distribution of Raindrops with Size. *Journal of Atmospheric Sciences*,
 644 5: 165-166.

645 Milbrandt, J.A., Yau, M.K., 2005. A Multimoment Bulk Microphysics Parameterization. Part II: A Proposed
 646 Three-Moment Closure and Scheme Description. *Journal of the Atmospheric Sciences*, 62(9): 3065-
 647 3081. DOI:10.1175/JAS3535.1

648 Niu, S. et al., 2010. Distributions of Raindrop Sizes and Fall Velocities in a Semiarid Plateau Climate:
 649 Convective versus Stratiform Rains. *Journal of Applied Meteorology and Climatology*, 49(4): 632-645.
 650 DOI:10.1175/2009jamc2208.1

651 Oue, M., Uyeda, H., Lee, D.-I., 2011. Raindrop size distribution parameters estimated from polarimetric
 652 radar variables in convective cells around Okinawa Island during the Baiu period. *Asia-Pacific Journal*
 653 *of Atmospheric Sciences*, 47(1): 33-44. DOI:10.1007/s13143-011-1003-x

654 Oue, M., Uyeda, H., Shusse, Y., 2010. Two types of precipitation particle distribution in convective cells
 655 accompanying a Baiu frontal rainband around Okinawa Island, Japan. *Journal of Geophysical*
 656 *Research: Atmospheres*, 115(D2). DOI:<https://doi.org/10.1029/2009JD011957>

657 Porcù, F., D'Adderio, L.P., Prodi, F., Caracciolo, C., 2014. Rain drop size distribution over the Tibetan
 658 Plateau. *Atmospheric Research*, 150: 21-30. DOI:<https://doi.org/10.1016/j.atmosres.2014.07.005>

659 Rosenfeld, D., Ulbrich, C.W., 2003. Cloud Microphysical Properties, Processes, and Rainfall Estimation
 660 Opportunities. In: Wakimoto, R.M., Srivastava, R. (Eds.), *Radar and Atmospheric Science: A Collection*
 661 *of Essays in Honor of David Atlas*. American Meteorological Society, Boston, MA, pp. 237-258.
 662 DOI:10.1007/978-1-878220-36-3_10

663 Radhakrishna, B., K. Saikranthi, and T. N. Rao, 2020: Regional differences in raindrop size distribution

664 within Indian subcontinent and adjoining seas as inferred from global precipitation measurement dual-
665 frequency precipitation radar. J. Meteor. Soc. Japan, 98, 573-584. [https://doi.org/10.2151/jmsj.2020-](https://doi.org/10.2151/jmsj.2020-030)
666 030

667 Rotstayn, L.D., 1997. A physically based scheme for the treatment of stratiform clouds and precipitation
668 in large-scale models. I: Description and evaluation of the microphysical processes. Quarterly Journal
669 of the Royal Meteorological Society, 123(541): 1227-1282.
670 DOI:<https://doi.org/10.1002/qj.49712354106>

671 Sauvageot, H., Lacaux, J.-P., 1995. The Shape of Averaged Drop Size Distributions. Journal of
672 Atmospheric Sciences, 52(8): 1070-1083. DOI:10.1175/1520-
673 0469(1995)052<1070:TSOADS>2.0.CO;2

674 Seela, B.K., Janapati, J., Lin, P.-L., Wang, P.K., Lee, M.-T., 2018. Raindrop Size Distribution
675 Characteristics of Summer and Winter Season Rainfall Over North Taiwan. Journal of Geophysical
676 Research: Atmospheres, 123(20): 11,602-11,624. DOI:<https://doi.org/10.1029/2018JD028307>

677 Seifert, A., 2005. On the Shape–Slope Relation of Drop Size Distributions in Convective Rain. Journal of
678 Applied Meteorology, 44(7): 1146-1151. DOI:10.1175/JAM2254.1

679 Seto, S., T. Iguchi, R. Meneghini, J. Awaka, T. Kubota, T. Masaki, and N. Takahashi, 2021: The
680 precipitation rate retrieval algorithms for the GPM Dual-frequency Precipitation Radar. J. Meteor. Soc.
681 Japan, 99, 205-237. <https://doi.org/10.2151/jmsj.2021-011>

682 Sun, J., 2005. Initialization and Numerical Forecasting of a Supercell Storm Observed during STEPS.
683 Monthly Weather Review, 133(4): 793-813. DOI:10.1175/MWR2887.1

684 Tao, S.-y., Ding, Y.-h., 1981. Observational Evidence of the Influence of the Qinghai-Xizang (Tibet)
685 Plateau on the Occurrence of Heavy Rain and Severe Convective Storms in China. Bulletin of the
686 American Meteorological Society, 62(1): 23-30. DOI:10.1175/1520-
687 0477(1981)062<0023:OEOTIO>2.0.CO;2

688 Tapiador, F.J., Checa, R., de Castro, M., 2010. An experiment to measure the spatial variability of rain
689 drop size distribution using sixteen laser disdrometers. Geophysical Research Letters, 37(16): L044120.
690 DOI:<https://doi.org/10.1029/2010GL044120>

691 Tapiador, F.J., Haddad, Z.S., Turk, J., 2014. A Probabilistic View on Raindrop Size Distribution Modeling:
692 A Physical Interpretation of Rain Microphysics. Journal of Hydrometeorology, 15(1): 427-443.
693 DOI:10.1175/JHM-D-13-033.1

694 Tapiador, F.J., Sánchez, J.-L., García-Ortega, E., 2019. Empirical values and assumptions in the
695 microphysics of numerical models. *Atmospheric Research*, 215: 214-238.
696 DOI:<https://doi.org/10.1016/j.atmosres.2018.09.010>

697 Testud, J., Oury, S., Black, R.A., Amayenc, P., Dou, X., 2001. The Concept of “Normalized” Distribution
698 to Describe Raindrop Spectra: A Tool for Cloud Physics and Cloud Remote Sensing. *Journal of Applied*
699 *Meteorology*, 40(6): 1118-1140. DOI:10.1175/1520-0450(2001)040<1118:TCONDT>2.0.CO;2

700 Thomas, A., Kanawade, V., Chakravarty, K., Srivastava, A., 2021. Characterization of raindrop size
701 distributions and its response to cloud microphysical properties. *Atmospheric Research*, 249: 105292.
702 DOI:10.1016/j.atmosres.2020.105292

703 Thompson, E.J., Rutledge, S.A., Dolan, B., Thurai, M., 2015. Drop Size Distributions and Radar
704 Observations of Convective and Stratiform Rain over the Equatorial Indian and West Pacific Oceans.
705 *Journal of the Atmospheric Sciences*, 72(11): 4091-4125. DOI:10.1175/JAS-D-14-0206.1

706 Tokay, A., Bashor, P.G., 2010. An Experimental Study of Small-Scale Variability of Raindrop Size
707 Distribution. *Journal of Applied Meteorology and Climatology*, 49(11): 2348-2365.
708 DOI:10.1175/2010JAMC2269.1

709 Tokay, A., Short, D.A., 1996. Evidence from Tropical Raindrop Spectra of the Origin of Rain from
710 Stratiform versus Convective Clouds. *Journal of Applied Meteorology and Climatology*, 35(3): 355-371.
711 DOI:10.1175/1520-0450(1996)035<0355:EFTRSO>2.0.CO;2

712 Ulbrich, C.W., 1983. Natural Variations in the Analytical Form of the Raindrop Size Distribution. *Journal*
713 *of Climate and Applied Meteorology*, 22(10): 1764-1775. DOI:10.1175/1520-
714 0450(1983)022<1764:Nvitaf>2.0.Co;2

715 Ulbrich, C.W., Atlas, D., 1998. Rainfall Microphysics and Radar Properties: Analysis Methods for Drop
716 Size Spectra. *Journal of Applied Meteorology*, 37(9): 912-923. DOI:10.1175/1520-
717 0450(1998)037<0912:RMARPA>2.0.CO;2

718 Wen, G., Xiao, H., Yang, H., Bi, Y., Xu, W., 2017. Characteristics of summer and winter precipitation over
719 northern China. *Atmospheric Research*, 197. DOI:10.1016/j.atmosres.2017.07.023

720 Wen, L. et al., 2016. Statistical characteristics of raindrop size distributions observed in East China during
721 the Asian summer monsoon season using 2-D video disdrometer and Micro Rain Radar data. *Journal*
722 *of Geophysical Research: Atmospheres*, 121(5): 2265-2282.
723 DOI:<https://doi.org/10.1002/2015JD024160>

724 Xu, X., Lu, C., Shi, X., Gao, S., 2008. World water tower: An atmospheric perspective. *Geophysical*
725 *Research Letters*, 35(20): L035867. DOI:10.1029/2008GL035867

726 Yamaji, M., H. G. Takahashi, T. Kubota, R. Oki, A. Hamada, and Y. N. Takayabu, 2020: 4-year climatology
727 of global drop size distribution and its seasonal variability observed by spaceborne Dual-frequency
728 Precipitation Radar. *J. Meteor. Soc. Japan*, 98, 755-773. <https://doi.org/10.2151/jmsj.2020-038>

729 Yasunari, T., Miwa, T., 2006. Convective Cloud Systems over the Tibetan Plateau and Their Impact on
730 Meso-Scale Disturbances in the Meiyu/Baiu Frontal Zone —A Case Study in 1998—. *Journal of the*
731 *Meteorological Society of Japan. Ser. II*, 84(4): 783-803. DOI:10.2151/jmsj.84.783

732 Yuter, S.E., Houze, R.A., 1997. Measurements of Raindrop Size Distributions over the Pacific Warm Pool
733 and Implications for Z–R Relations. *Journal of Applied Meteorology*, 36(7): 847-867. DOI:10.1175/1520-
734 0450(1997)036<0847:MORSDO>2.0.CO;2

735 Yuter, S.E., Kingsmill, D.E., Nance, L., Löfflermang, M., 2006. Observations of Precipitation Size and Fall
736 Speed Characteristics within Coexisting Rain and Wet Snow. *Journal of Applied Meteorology and*
737 *Climatology*, 45(10): 1450-1464.

738 Zhang, A. et al., 2019. Statistical Characteristics of Raindrop Size Distribution in the Monsoon Season
739 Observed in Southern China. *Remote Sensing*, 11(4): 432.

740 Zhang, G., Vivekanandan, J., Brandes, E., 2001. A method for estimating rain rate and drop size
741 distribution from polarimetric radar measurements. *IEEE Transactions on Geoscience and Remote*
742 *Sensing*, 39(4): 830-841. DOI:10.1109/36.917906

743 Zhang, G., Vivekanandan, J., Brandes, E.A., Meneghini, R., Kozu, T., 2003. The Shape–Slope Relation
744 in Observed Gamma Raindrop Size Distributions: Statistical Error or Useful Information? *Journal of*
745 *Atmospheric and Oceanic Technology*, 20(8): 1106-1119. DOI:10.1175/1520-
746 0426(2003)020<1106:Tsriog>2.0.Co;2

747 Zhao, G. et al., 2011. Improving the rainfall rate estimation in the midstream of the Heihe River Basin
748 using raindrop size distribution. *Hydrology and Earth System Sciences - HYDROL EARTH SYST SCI*,
749 15: 943-951. DOI:10.5194/hess-15-943-2011

750 Zhao, Y. et al., 2019. Effects of convection over the Tibetan Plateau on rainstorms downstream of the
751 Yangtze River Basin. *Atmospheric Research*, 219: 24-35.
752 DOI:<https://doi.org/10.1016/j.atmosres.2018.12.019>

Fig. 1. Geographical location and surrounding topography of Mount Lushan (a), topographical features of Mount Lushan (b), overview of the observational platform (c), and the Thies CLIMA Laser Precipitation Monitor (LPM) (d). Locations of the observational platform (red pentagram) and Lushan Meteorological Bureau (blue pentagram) are marked in (b), and the position of the LPM (red arrow) is shown in (c).

Fig. 2. Schematic diagram of quality control. The color blocks indicate the numbers of the sampled particles and the solid line and dash lines show the corrected V-D relationship and the range of $V(D_i) \pm 0.6V(D_i)$. Particles falling outside the dash lines were removed.

766

767 **Fig. 3.** Accumulated sample numbers (blue) and precipitation amounts (orange) for each rain rate class

768 in Mount Lushan from the Meiyu dataset. Numbers show the percentages (%) for each rain rate class.

769

Fig. 4. Average RSD spectra for different rain rate classes of $0.1 \leq R < 1$ mm/h (blue), $1 \leq R < 5$ mm/h (orange), $5 \leq R < 10$ mm/h (yellow), $10 \leq R < 100$ mm/h (purple), $R \geq 100$ mm/h (green), and all data (black).

Fig. 5. Fitting results of the ML method (dash lines) and the MM method (dotted lines) for average spectra of each rain rate class (a)-(e) and the entire dataset. Blue lines indicate the observed spectra.

Fig. 6. Histograms of frequency for D_m (a) and $\log_{10} N_w$ (b). Gray shades, blue/red lines, and

cyan/magenta lines represent the entire dataset, stratiform rains classified by the T01 and B03 scheme, and convective rains classified by the T01 and B03 scheme, respectively.

Fig. 7. Scatter plots of D_m vs $\log_{10} N_w$ for stratiform (light blue dots) and convective (pink dots) rain types and the average values and standard deviations for stratiform (blue line) and convective (red line) RSD. The horizontal and oblique dash gray lines indicate the stratiform-convective rain type separation lines from Thompson et al. (2015) and Bringi et al. (2009), and the Maritime and Continental rain type rectangles (gray rectangles) are from Bringi et al. (2003). Average D_m - $\log_{10} N_w$ pairs from the results of Bringi et al. (2006, '×', 'Bringi06'), Wen et al. (2016, '▼', 'Wen16'), Chen et al. (2013, '●', 'Chen 13'), and Fu et al. (2020, '◆', 'Fu20') are also marked in the D_m - $\log_{10} N_w$ coordinate. Colors represent values of stratiform rain (blue), convective rain (red), and all data (black). S, C, and A represent stratiform, convective, and all, respectively.

Fig. 8. Distributions of D_m and N_w with R for stratiform (a, b) and convective (c, d) rains. Blue and magenta lines indicate the power fitting results for stratiform and convective rains, and the fitting equations are also presented in corresponding diagrams. The light blue scatterplots represent the samples with $\log_{10} N_w > 4.5$ and the cyan lines indicate their fitting results of corresponding samples.

799

800 **Fig. 9.** Average RSD spectra of $\log_{10} N_w > 4.5$ (cyan line), $\log_{10} N_w < 4.5$ (magenta line), and all samples
 801 (blue line) in stratiform rains.

802

Fig. 10. Average spectra (blue/orange dots) and fitting results (blue/red dash dotted lines) of stratiform/convective rains.

Fig. 11. Scatterplots of $\mu - N_0$ (a) and $\mu - \Lambda$ (b) for stratiform (blue dots) and convective (orange dots) rains. The blue, red, and black solid lines indicate the exponential (a) and quadratic polynomial (b) fitting results of $\mu - N_0$ and $\mu - \Lambda$ relationship for stratiform rains (S), convective rains (C), and the entire dataset (A). The fitting equations are also presented inside the diagrams. The black dashed line in (b) shows the $\mu - \Lambda$ fitting equation of Zhang et al. (2003), which is $\Lambda = 0.0365\mu^2 + 0.735\mu + 1.935$. All the data in Fig. 11 were filtered using the standard of $N_t > 1000 m^{-3}$ and $R > 5 mm/h$ (Zhang et al., 2003).

Fig. 12. $Z - R$ scatterplots and fitting results for the Meiyu season at Mount Lushan: (a) the blue and orange scatterplots represent the samples with $\log_{10}N_w > 4.5$ and $\log_{10}N_w < 4.5$ in the entire dataset and the power fitting results of samples with $\log_{10}N_w > 4.5$, $\log_{10}N_w < 4.5$, and the entire dataset are presented by red, blue, and magenta lines; (b) the blue and orange scatterplots indicate the stratiform and convective samples. Among the blue scatterplots, the light/deep blue color represents the samples with $\log_{10}N_w > 4.5/\log_{10}N_w < 4.5$. The red line, blue solid line, blue dash dotted line, and blue dotted line show the power fitting results of each dataset. The dash, dash dotted, and dotted black lines indicate the power laws of stratiform rain $Z = 200R^{1.6}$ (Marshall and Palmer, 1948), operational WSR-88D $Z = 300R^{1.4}$ (Fulton et al., 1998), and Meiyu result in Eastern China $Z = 368R^{1.21}$ (Chen et al., 2013).

Fig. 13. Two examples of precipitation processes with high N_w values. The first (a, b) is from 12:00 on June 28 to 07:40 on June 30, 2016, and the second is from 20:20 on July 15 to 06:00 July 17, 2019. The color maps and magenta lines in (a)/(c) represent the RSD and rain rate time series, and the blue and orange lines in (b)/(d) represent for the D_m and $\log N_w$ time series.

Fig. 14. An example of in-cloud precipitation (taken on August 12, 2020).

Fig. 15. (a) and (b) are the same as Fig. 13, but for the precipitation process from 00:00 May 23 to 16:00 on May 24, 2017, (c) shows the time series of drop size distributions from the fog monitor (color maps) and visibility (magenta line).

842 **Table 1.** Correlation coefficients, average absolute biases, average relative biases, N_T (m⁻³), and
 843 W (g/m³) of the fitting results using the ML and MM methods for each rain rate class and the entire
 844 dataset.

R class	Corr. Coef.		Abs. Bias		Rel. Bias (%)		N_T (m ⁻³)			W (g/m ³)		
	ML	MM	ML	MM	ML	MM	ML	MM	Obs.	ML	MM	Obs.
0.1≤R<1	0.86	0.50	418	684	126.5	31.6	1526	1503	1216	0.09	0.05	0.05
1≤R<5	0.93	0.56	195	558	109.4	25.1	1614	2265	1475	0.26	0.17	0.17
5≤R<10	0.89	0.69	212	357	77.9	29.7	1983	2457	1818	0.56	0.41	0.42
10≤R<100	0.65	0.58	1433	1494	56.5	33.9	7288	5668	5469	2.29	1.34	1.32
R≥100	0.53	0.49	6721	6234	75.6	49.6	31846	17165	20152	15.19	6.73	6.39
All	0.75	0.52	409	668	93.1	29.2	2371	2523	1899	0.56	0.28	0.29

845

846 **Table 2.** Integral Parameters derived from average RSDs of each rain rate class and the whole dataset.

R class	Z (dBZ)	D_m (mm)	$\log_{10}N_w$	σ_M (mm)	σ_M/D_m	N_0 ($m^{-3}mm^{-1-\mu}$)	μ	Λ (mm^{-1})
0.1≤R<1	20.0	0.73	4.15	0.44	0.60	1775	-1.25	3.52
1≤R<5	30.9	1.14	3.92	0.67	0.59	2326	-1.11	2.44
5≤R<10	38.1	1.55	3.77	0.86	0.56	3287	-0.76	2.07
10≤R<100	47.5	2.14	3.71	1.28	0.60	3366	-1.22	1.32
R≥100	58.3	2.97	3.82	1.82	0.61	7653	-1.33	0.96
All	39.6	1.78	3.37	1.20	0.67	721	-1.79	1.20

847

848

Table 3. Fitting results of gamma distribution for stratiform and convective rains.

Rain type	N_0 ($\text{m}^{-3}\text{mm}^{-1-\mu}$)	μ	λ (mm^{-1})
Stratiform	2368	-1.37	1.24
Convective	1220	-1.45	2.30
All	721	-1.79	1.20

849

850

Table 4. Size bins and velocity bins for LPM and Parsivel.

Size Bins (mm)		Velocity Bins (mm/h)	
LPM	Parsivel	LPM	Parsivel
	0.000-0.125		0.0-0.1
0.125-0.250	0.125-0.250	0.0-0.2	0.1-0.2
0.250-0.375	0.250-0.375	0.2-0.4	0.2-0.3
0.375-0.500	0.375-0.500	0.4-0.6	0.3-0.4
0.500-0.750	0.500-0.625	0.6-0.8	0.4-0.5
0.750-1.000	0.625-0.750	0.8-1.0	0.5-0.6
1.000-1.250	0.750-0.875	1.0-1.4	0.6-0.7
1.250-1.500	0.875-1.000	1.4-1.8	0.7-0.8
1.500-1.750	1.000-1.125	1.8-2.2	0.8-0.9
1.750-2.000	1.125-1.250	2.2-2.6	0.9-1.0
2.000-2.500	1.250-1.500	2.6-3.0	1.0-1.2
2.500-3.000	1.500-1.750	3.0-3.4	1.2-1.4
3.000-3.500	1.750-2.000	3.4-4.2	1.4-1.6
3.500-4.000	2.000-2.250	4.2-5.0	1.6-1.8
4.000-4.500	2.250-2.575	5.0-5.8	1.8-2.0
4.500-5.000	2.575-3.000	5.8-6.6	2.0-2.4
5.000-5.500	3.000-3.500	6.6-7.4	2.4-2.8
5.500-6.000	3.500-4.000	7.4-8.2	2.8-3.2
6.000-6.500	4.000-4.500	8.2-9.0	3.2-3.6
6.500-7.000	4.500-5.125	9.0-10.0	3.6-4.1
7.000-7.500	5.125-6.000	> 10.0	4.1-4.8
7.500-8.000	6.000-7.000		4.8-5.6
> 8.000	7.000-8.000		5.6-6.4
	8.000-9.000		6.4-7.2
	9.000-10.250		7.2-8.2
	10.250-12.000		8.2-9.6
	12.000-14.000		9.6-11.2
	14.000-16.000		11.2-12.8
	16.000-18.000		12.8-14.4
	18.000-20.000		14.4-16.4
	20.000-23.000		16.4-19.2
	23.000-26.000		19.2-21.4