551.509.313 : 551.511.32 : 551.510.528

Dynamical Effects of the Variable Tropopause

By **Takashi Nitta***

Electronic Computation Center, Japan Meteorological Agency, Tokyo

(Manuscript received 22 December 1964, in revised form 4 March 1965)

Abstract

A computational model for large scale motions which extends to the stratosphere is designed. The model has the tropopause and ground surface, and is favorable in treating the interaction between the tropospheric and stratospheric motions. A special σ -coordinate in the vertical is used, and the tropopause is assumed to be a material surface at which $d\sigma/dt$ vanishes.

Fundamental equations of the model are illustrated, and energetical considerations are given. Particular interest is put in the interaction of kinetic energy between the troposphere and the stratosphere in the short term evolution of the atmospheric motion. Calculation based on the actual data is made for the vertical transfer of kinetic energy through the tropopause.

Analytical solutions of perturbations superimposed on the basic zonal current are obtained for barotropic and baroclinic models and compared with those of p -coordinate system. The kinetic energy transport in vertical as well as control of the false westward retrogression of the ultra-long wave, due to the non-vanishing vertical mean divergence in the troposphere, are shown. Modification of the criterion of baroclinic instability due to the existence of the stratosphere is discussed.

1. Introduction

Recently the atmospheric model of numerical weather prediction has been refined by introduction of more detailed physical processes in lower troposphere, such as the non-adiabatic heating and cooling, and the effect of orography.

It seems, however, that there remain several important problems unsolved. One of them is the extension of model toward the stratosphere, which will enable us to treat the ultra-long waves more accurately. Gates (1957), Eliassen (1960) and Murakami (1964) have proposed atmospheric models for this purpose. Gates' model consists of two baroclinic layers (corresponding to the troposphere and the stratosphere) separated by a tropopause which is considered as a material surface. Eliassen has presented a quasi-Lagrangian model where the atmosphere is bounded by material surfaces, and the tropopause is chosen as one of

coordinate planes. To include the stratosphere in the model atmosphere, Murakami has recommended a vertical σ -coordinate as $\sigma = p/p_T$ for the stratosphere, and $\sigma = (p - p_T)/(p_S - p_T)$ for the troposphere, where p_T and p_S stand for pressures at the tropopause and the ground surface respectively.

The purposes of the present paper are to see the features of the kinetic energy transfer through the tropopause and to examine the behaviors of the tropospheric disturbance under the variable tropopause, with special interest on very long wave, with respect to an atmospheric model extending to the stratosphere. In the following, we will use the x, y, σ -coordinate system as the same as that of Murakami, and will also use the assumption that the tropopause is a material surface.

At first we derive the fundamental equations and the energetical relations on the σ -coordinate. Secondly, the interaction between the troposphere and the stratosphere are considered, and calculations are made to see the short term vertical propagation of kinetic energy through the tropopause. Finally, we obtain

* Present address: National Meteorological Center, U.S. Weather, Bureau, Suitland, Md., U.S.A.

wave solutions from the linearized equations, and discuss on their properties of the ultra-long waves and the effect of the variable tropopause on the tropospheric motion. Comparisons are made between the forecast results by this model and by the conventional model of p -system.

2. Coordinate and modelling assumption

Nothing can be said about the maintenance of the tropopause without taking into account the radiational effect in the atmosphere. In the case of short range forecasting, however, incorporation of the influence of the stratosphere through the tropopause in an atmospheric model could be accomplished by assuming the existence of the stratosphere at initial time and keeping physical properties of the stratosphere from distortion, by some means, in the process of numerical calculation.

In the following discussion, we use the σ -coordinate similar to that of Murakami, *i.e.*,

Fig. 1. Vertical resolution in σ -coordinate.

$$\left. \begin{aligned} \sigma &= \frac{p}{p_T}, \quad (0 \leq p \leq p_T): \\ &\quad \text{in the stratosphere} \\ \text{and} \\ \sigma &= (1-\varepsilon) \frac{p-p_T}{p_s-p_T} + \varepsilon, \quad (p_T \leq p \leq p_s): \\ &\quad \text{in the troposphere,} \end{aligned} \right\} \quad (2.1)$$

where $0 < \varepsilon < 1$, and p_T denotes the pressure at

the tropopause and p_s that at the ground. The value of ε is to be determined according to the degree of vertical resolution desired: for instance, $\varepsilon=1/2$ is suitable for representing the 2-layer model (the troposphere and the stratosphere), and $\varepsilon=1/3$ for 3-layer model (two layers in the troposphere and one in the stratosphere). Although there would be many versions of σ -coordinate other than mentioned above, we particularly choose (2.1) simply because σ is linear with p and then the difference of σ corresponds to the mass between the two levels. This fact enables us to solve the problem easily. But on the other hand it requires an additional boundary condition on $\dot{\sigma} = d\sigma/dt$ at the internal surface, *i.e.*, at the tropopause.

Concerning this point, we assume that the tropopause is a material surface; $\dot{\sigma}$ is zero at the level $\sigma=\varepsilon$. Moreover, it is also assumed that the atmosphere is isentropic in the stratosphere and accordingly $\dot{\sigma}$ vanishes on every coordinate plane.

3. Fundamental equations and energy equations

Using the definition of σ , we can derive fundamental equations of motion and the thermal equations on σ -coordinate for the stratosphere (St) and the troposphere (Tr) as follows.

(1) Equation of motion

$$\text{St: } \frac{dV}{dt} + f\mathbf{k} \times V + \nabla\phi - \frac{\sigma}{p_T} \left(\frac{\partial\phi}{\partial\sigma} \right) \nabla p_T = \mathbf{F}, \quad (3.1.1)$$

$$\text{Tr: } \frac{dV}{dt} + f\mathbf{k} \times V + \nabla\phi - \frac{1}{\pi} \left(\frac{\partial\phi}{\partial\sigma} \right) \times \nabla(\sigma\pi + p_T - \varepsilon p_s) = \mathbf{F}, \quad (3.1.2)$$

where $\pi = p_s - p_T$, $\frac{d}{dt} \equiv \frac{\partial}{\partial t} + \mathbf{V} \cdot \nabla + \dot{\sigma} \frac{\partial}{\partial \sigma} \equiv \frac{d_H}{dt} + \dot{\sigma} \frac{\partial}{\partial \sigma}$, $\mathbf{F}$ shows a dissipation and other notations are as usual. $\dot{\sigma}$ relates to $\omega \equiv dp/dt$ as

$$\text{St: } \omega = \frac{1}{\varepsilon} \left(\sigma \frac{d_H p_T}{dt} + p_T \dot{\sigma} \right), \quad (3.2.1)$$

$$\text{Tr: } \omega = \frac{1}{1-\varepsilon} \frac{d_H}{dt} (\sigma\pi + p_T - \varepsilon p_s) + \frac{\pi}{1-\varepsilon} \dot{\sigma}. \quad (3.2.2)$$

(2) Continuity equation

$$St: \frac{\partial}{\partial t} p_T + \nabla \cdot (p_T \mathbf{V}) + \frac{\partial}{\partial \sigma} (p_T \dot{\sigma}) = 0, \quad (3.3.1)$$

$$Tr: \frac{\partial}{\partial t} \pi + \nabla \cdot (\pi \mathbf{V}) + \frac{\partial}{\partial \sigma} (\pi \dot{\sigma}) = 0. \quad (3.3.2)$$

Integration of (3.3) with σ give the following formulae to determine $\dot{\sigma}$, $\partial p_T / \partial t$ and $\partial \pi / \partial t$.

$$St: p_T \dot{\sigma} = - \int_0^\sigma \nabla \cdot (p_T \mathbf{V}) d\sigma - \sigma \frac{\partial p_T}{\partial t}, \quad (3.4.1)$$

$$\frac{\partial p_T}{\partial t} = - \frac{1}{\varepsilon} \left\{ \int_0^\varepsilon \nabla \cdot (p_T \mathbf{V}) d\sigma + p_T \dot{\sigma}_T \right\}, \quad (3.5.1)$$

where $\dot{\sigma}_T = (\varepsilon / p_T) (\omega_T - d_H p_T / dt)$.

$$Tr: \pi \dot{\sigma} = \pi \dot{\sigma}_T - \int_\varepsilon^\sigma \nabla \cdot (\pi \mathbf{V}) d\sigma - (\sigma - \varepsilon) \frac{\partial \pi}{\partial t}, \quad (3.4.2)$$

$$\frac{\partial \pi}{\partial t} = \frac{1}{1 - \varepsilon} \left\{ \pi \dot{\sigma}_T - \int_\varepsilon^1 \nabla \cdot (\pi \mathbf{V}) d\sigma \right\}, \quad (3.5.2)$$

where $\dot{\sigma}_T = ((1 - \varepsilon) / \pi) (\omega_T - d_H \pi / dt)$.

The relation that $\dot{\sigma} = 0$ at $\sigma = 0$, $\sigma = \varepsilon$ and $\sigma = 1$ should be used if necessary.

(3) Thermal equation

$$St \text{ and } Tr: \frac{d\theta}{dt} = \frac{1}{c_p} \left(\frac{p_{10}}{p} \right)^\kappa \frac{dQ}{dt}, \quad (3.6)$$

where $\kappa = R / c_p$, and dQ / dt shows non-adiabatic heating.

(4) Hydrostatic relation

$$St: \frac{\partial \phi}{\partial \sigma} = - \frac{RT}{\sigma} = - \frac{R}{\sigma} \left(\frac{p_T}{\varepsilon p_{10}} \right)^\kappa \theta, \quad (3.7.1)$$

$$Tr: \frac{\partial \phi}{\partial \sigma} = - \frac{\pi RT}{\sigma \pi + p_T - \varepsilon p_s} \\ = - \frac{\pi R}{\sigma \pi + p_T - \varepsilon p_s} \left(\frac{\sigma \pi + p_T - \varepsilon p_s}{(1 - \varepsilon) p_{10}} \right)^\kappa \theta. \quad (3.7.2)$$

(5) Energy equation

5.1) Kinetic energy

$$St: \frac{\partial}{\partial t} (p_T K) + \nabla \cdot (p_T K \mathbf{V} + p_T \phi \mathbf{V}) \\ + \frac{\partial}{\partial \sigma} (p_T K \dot{\sigma} + p_T \phi \dot{\sigma}) + \frac{\partial (\sigma \phi)}{\partial \sigma} \frac{\partial p_T}{\partial t} \\ + p_T \omega \alpha = p_T \mathbf{V} \cdot \mathbf{F}, \quad (3.8.1)$$

where $K = \frac{1}{2} \mathbf{V}^2$ and $\alpha = -(\varepsilon / p_T) (\partial \phi / \partial \sigma)$ is a specific volume.

$$Tr: \frac{\partial}{\partial t} (\pi K) + \nabla \cdot (\pi K \mathbf{V} + \pi \phi \mathbf{V}) \\ + \frac{\partial}{\partial \sigma} (\pi K \dot{\sigma} + \pi \phi \dot{\sigma}) + \frac{\partial (\sigma \phi)}{\partial \sigma} \frac{\partial \pi}{\partial t} + \pi \omega \alpha \\ + \left(\frac{\partial \phi}{\partial \sigma} \right) \frac{\partial}{\partial t} (p_T - \varepsilon p_s) = \pi \mathbf{V} \cdot \mathbf{F}, \quad (3.8.2)$$

where $\alpha = -((1 - \varepsilon) / \pi) (\partial \phi / \partial \sigma)$.

5.2) Potential energy

$$St: \frac{\partial}{\partial t} (p_T c_p T) + \nabla \cdot (p_T c_p T \mathbf{V}) \\ + \frac{\partial}{\partial \sigma} (p_T c_p T \dot{\sigma}) - p_T \omega \alpha = p_T \frac{dQ}{dt}, \quad (3.9.1)$$

$$Tr: \frac{\partial}{\partial t} (\pi c_p T) + \nabla \cdot (\pi c_p T \mathbf{V}) \\ + \frac{\partial}{\partial \sigma} (\pi c_p T \dot{\sigma}) - \pi \omega \alpha = \pi \frac{dQ}{dt}. \quad (3.9.2)$$

(6) Total energy equations

Integration of energy equations for total mass in the stratosphere and the troposphere leads to, under the assumption of vanishing of energy transfer at the lateral boundary,

$$St: \frac{\partial}{\partial t} \int_{\varepsilon g}^1 p_T K dV = - \int_{\varepsilon g}^1 \phi_T \frac{\partial p_T}{\partial t} dS \\ - \int_{\varepsilon g}^1 p_T \omega \alpha dV + \int_{\varepsilon g}^1 p_T \mathbf{V} \cdot \mathbf{F} dV, \quad (3.10.1)$$

$$Tr: \frac{\partial}{\partial t} \int \frac{1}{(1 - \varepsilon) g} \pi K dV \\ = \int \frac{1}{g} \left(\phi_T \frac{\partial p_T}{\partial t} - \phi_s \frac{\partial p_s}{\partial t} \right) dS \\ - \int \frac{1}{(1 - \varepsilon) g} \pi \omega \alpha dV \\ + \int \frac{1}{(1 - \varepsilon) g} \pi \mathbf{V} \cdot \mathbf{F} dV, \quad (3.10.2)$$

$$St: \frac{\partial}{\partial t} \int_{\varepsilon g}^1 p_T c_p T dV = \int_{\varepsilon g}^1 p_T \omega \alpha dV \\ + \int_{\varepsilon g}^1 p_T \frac{dQ}{dt} dV, \quad (3.11.1)$$

$$Tr: \frac{\partial}{\partial t} \int \frac{1}{(1 - \varepsilon) g} \pi c_p T dV \\ = \int \frac{1}{(1 - \varepsilon) g} \pi \omega \alpha dV$$

$$+ \int \frac{1}{(1-\varepsilon)g} \pi \frac{dQ}{dt} dV, \quad (3.11.2)$$

where $\int ( ) dV \equiv \int ( ) dx dy d\sigma$ and $\int ( ) dS \equiv \int ( ) dx dy$ and the assumption that $\dot{\sigma}_T = 0$ at $\sigma = \varepsilon$ is used.

Needless to say, terms including $\omega\alpha$ play roles in the exchange between kinetic and potential energies, and $(\phi_T/g)(\partial p_T/\partial t)$ and $(\phi_s/g)(\partial p_s/\partial t)$ are related to the exchange of energy between the upper and the lower layers around the tropopause and the ground surface respectively. As we can easily see from (3.2), the term $(\phi_T/g)(\partial p_T/\partial t)$ is a part of the kinetic energy transfer, $(\omega\phi)/g$ at the tropopause in the x, y, p -system. Thus, we may say that the variation of p_T with time, $\partial p_T/\partial t$, contributes to the kinetic energy transfer through the tropopause, which will be shown in the next section. At the same time, the presence of the pressure change with time or ω at the tropopause is identical with the fact that the vertical mean divergence in the troposphere does not vanish. In this meaning, the non-zero $\partial p_T/\partial t$ acts just like as an anchor or block for the spurious westward retrogression of the ultra-long waves.

4. Kinetic energy transfer through the tropopause

The energy transfer through the tropopause was investigated theoretically by Charney and Drazin (1961). With the use of an atmospheric model depicted here, let us examine the transfer of energy in more detail. We separate $\phi_T(\partial p_T/\partial t)$ into two parts, *i.e.*, $\bar{\phi}_T(\partial \bar{p}_T/\partial t)$ and $\phi_T'(\partial p_T'/\partial t)$, where bar denotes space mean at $\sigma = \sigma_T$ and primed quantity is the deviation from its average. If we take the area sufficiently large, $(\partial \bar{p}_T/\partial t)$ becomes almost zero. $\phi_T'(\partial p_T'/\partial t)$ has close connection with the motion of tropospheric disturbances. Namely, when cyclone in the troposphere develops, the height of the tropopause associated with the trough tends to lower and accordingly the pressure at the tropopause increases. In the case of development of anticyclone like formation of blocking situation, almost inverse tendency is observed. Since these observational facts indicate a negative correlation between ϕ_T' and $\partial p_T'/\partial t$, it may be said from (3.10) that

kinetic energy is transferred from the troposphere toward the stratosphere; that is, the troposphere acts as an energy source to the stratosphere, when large scale troughs or ridges develop. Fig. 2 illustrates the schematic relation of ϕ_T' and $\partial p_T'/\partial t$ which can be expressed as

Fig. 2. Schematic configuration of ϕ_T' and $\partial p_T'/\partial t$ in sinusoidal pattern.

$$\phi_T' = A \sin kx, \quad (A > 0),$$

$$\frac{\partial p_T'}{\partial t} = -B \sin k(x - \delta), \quad (B > 0).$$

Hence

$$\phi_T' \frac{\partial p_T'}{\partial t} = -C\Omega$$

where $C = AB$ and $\Omega = \cos k\delta - \cos k(2x - \delta)$.

(i) *The case of trough* ($\pi \leq kx \leq 2\pi$)

Occlusion: $\delta \sim 0$ and $\Omega(>0)$ is large. $A(>0)$ is maximum but $B(>0)$ becomes small and then changes sign (*i.e.*, upward and then downward transports). (Fig. 2a)

Development of migrating trough: $\delta \neq 0$ and $\Omega(>0)$ decreases. Both $A(>0)$ and $B(>0)$ increase (*i.e.*, upward transport). (Fig. 2b). Hence, considering the fact that a short wave moves in general more rapidly than a long wave, and thus Ω and C of the former are smaller than those of the latter, we can im-

agine that the amplifying long wave, especially stationary wave, plays more important role on the strato-tropospheric interaction than the short wave.

(ii) *The case of ridge* ($0 \leq kx \leq \pi$)

Blocking: $\delta \sim 0$ and $\Omega(>0)$ is large. $A(>0)$ is maximum but becomes small and afterwards changes sign (*i.e.*, upward and afterwards downward transfer). (Fig. 2c).

Formative stage of blocking or amplification of moving ridge: $\delta \neq 0$ and $\Omega(>0)$ is small. Both $A(>0)$ and $B(>0)$ increase (*i.e.*, upward transfer). (Fig. 2d).

5. Numerical estimation of the energy transfer

Using the results of the aerological study of Defant and Taba (1958), we computed the correlation of ϕ_T' or p_T' and $\partial p_T' / \partial t \approx \Delta p_T' / \Delta t$. In Fig. 3 shown are the profiles of pressure at the tropopause and their time changes for 24 hours along a specified latitudinal arc. The

Fig. 3a.

former are interpolated, by consideration of the movement of disturbances, from two successive maps of the tropopause pattern illustrated by Defant and Taba. The latter are computed as the difference between two successive profiles of p_T . Figs. 3 a, b, c are zonal crosssections at 45°, 40° and 35°N, for the dates from 5–6 January, 1956 in (a) to 8–9 in (c). Thus, we can follow the profiles of p_T and Δp_T with time. The trough near 80°W begins to develop on 6–7 January and moves to south-eastwards at the speed of about 5° lat. per day. On 8–9 January, the development ceases and tends to decay. The assumption that the tropopause is a material surface and it has a constant value of potential temperature seems to be almost valid so far as this example is concerned. (Defant and Taba, 1957).

Table 1. Examples of the correlation $(\phi_T'/g)\partial p_T'/\partial t \simeq (\phi_T'/g)\Delta p_T'/\Delta t$ calculated from the profiles of p_T and Δp_T illustrated in Fig. 3 after Defant and Taba (1958). Minus sign shows upward propagation and plus sign downward. (Unit: joule m⁻² sec⁻¹)

	Jan. 5-6	6-7	7-8	8-9	Mean
45°N	4.3	-94.8			-45.3
40		-56.7	-29.3		-43.0
35			-83.9	65.7	-9.1

In Table 1, given is the numerical value of the vertical propagation of kinetic energy through the tropopause, which is calculated from Fig. 3 and averaged along latitudinal arc. Computation was carried out with grid interval of 5 degree longitude. We can presume from this result that the correlation $|\bar{p}_T' \cdot (\partial p_T'/\partial t)|$ has large values at developing stage, and that as soon as deepening stops, the correlation coefficient becomes small or sometimes even the sign changes. It is noted that the magnitude of the propagation of kinetic energy in vertical is of the order of 10 joule m⁻² sec⁻¹, but it often changes its sign. The time mean value may be in the order of 1 joule m⁻² sec⁻¹.

So far as the short term variation of an atmospheric motion is concerned, the interaction of kinetic energy between the troposphere and the lower stratosphere is rather active and the direction of energy propagation in vertical changes frequently. We can, however,

say nothing, from these limited informations, about the long term features of the propagation and of the interaction between the troposphere and the stratosphere.

6. Dynamical aspects of the tropospheric motion under the variable tropopause with special emphasis on very long wave

In numerical prediction with a quasi-geostrophic model in p -coordinate we usually pose the following boundary conditions in vertical, *i.e.*,

$$\omega=0 \text{ at } p=p_U \text{ and } p_L,$$

where p_U and p_L refer to pressure at the top and the lowest levels of the atmosphere, and the effect of orography is disregarded. When this method is used with a rough vertical resolution, one can not expect to include successfully an interaction of the troposphere and the layer above it. That is because the vertical mean divergence vanishes mostly in the troposphere. Because of this difficulty such an atmospheric model in p -system fails to describe accurately the motion of the ultra-long wave and induces very rapid westward spurious movement as the result of the so-called β -effect. To avoid this shortcoming an artificial non-vanishing mean divergence was introduced in the model. (Cressman, 1958; Wiin-Nielsen, 1959).

Concerning these points, the atmospheric model mentioned in the preceding section is the one which is favorable to avoid the difficulty and is probably capable of simulating the real situation even with a coarse vertical resolution. To see these circumstances in more detail, let us solve the perturbation solution superposed on a zonal current in barotropic and baroclinic atmosphere incorporating the existence of the tropopause.

(1) Equivalent barotropic model

We shall in the following make an assumption for the vertical distribution of height Z :

$$\bar{Z}(x, y, p, t) = Z_0(p) + A(p) \cdot \bar{Z}(x, y, t), \quad (6.1)$$

where

$$Z_0(p) = \frac{T_0}{\Gamma} \left[1 - \left(\frac{p}{p_0} \right)^{\Gamma/\Gamma_0} \right],$$

$$A(p) = a_1 p^2 + a_2 p + a_3,$$

$$a_1 = -\frac{3}{64 \times 10^4} \text{mb}^{-2}, \quad a_2 = \frac{1}{200} \text{mb}^{-1}, \quad a_3 = -\frac{5}{16},$$

Po _____

Fig. 4. Vertical resolution in a barotropic model.

T_0 is the temperature at the ground surface (288°K), Γ is the constant lapse rate ($0.65^\circ\text{C}/100\text{m}$), and $A(p)$ becomes 0 at $p=p_0=1000\text{mb}$, 1.0 at $p=p_0-(p_0-\bar{p}_T)/2=600\text{mb}$ and 0.5 at $p=\bar{p}_T=200\text{mb}$ (Fig. 4). The functional form of $A(p)$ mentioned above is determined by the observed values of disturbance in the troposphere, this form may not be adequate for the basic zonal flow, but for the sake of simplicity we use the same formula of $A(p)$ both for the disturbance and the zonal current, since the form of $A(p)$ for the zonal flow is not essential here.

Thus, we can get the vertical distribution of potential temperature θ from (6.1) as

$$\theta(x, y, p, t) = \frac{g\bar{p}}{R} \left(\frac{p_0}{p} \right)^\kappa \left[\frac{RT_0}{gp_0} \left(\frac{p}{p_0} \right)^{R\Gamma/g-1} - (2a_1\bar{p} + a_2) \bar{Z}(x, y, t) \right]. \quad (6.2)$$

Since we consider the tropopause as a material surface, $\theta=\theta_T=\text{constant}$ at $p=p_T$, where p_T is the pressure distribution at the tropopause. For the sake of simplicity, we assume that

$$p_T(x, y, t) = \bar{p}_T (= \text{const.}) + p_T'(x, y, t).$$

Inserting $\theta=\theta_T$ and $p=p_T$ in (6.2) and using an inequality $|p_T'/\bar{p}_T| < 1$, we have the first approximation for $p_T'(x, y, t)$, after expanding (6.2) with $p_T'/\bar{p}_T$ and ignoring higher order terms, as follows.

$$p_T' = \frac{\bar{p}_T [b_1 - b_2' - b_3 \bar{p}_T \cdot \bar{Z}(x, y, t)]}{\left[b_1(1-\kappa) + b_2' \left(\frac{R\Gamma}{g} - 1 \right) - 2a_1 \bar{p}_T \cdot \bar{Z}(x, y, t) \right]}, \quad (6.3)$$

where

$$b_1 = \frac{R\bar{T}_T}{g}, \quad \bar{T}_T = \theta_T \left(\frac{\bar{p}_T}{p_0} \right)^\kappa \sim 216^\circ\text{K}, \quad b_2 = \frac{RT_0}{g},$$

$$b_2' = b_2 \left(\frac{\bar{p}_T}{p_0} \right)^{R\Gamma/g}, \quad \text{and } b_3 = -(2a_1\bar{p}_T + a_2).$$

As an amplitude of pressure at the tropopause is not so small compared with $\bar{p}_T$ itself, the inequality is more or less crude but the essential characteristics would still be kept. If we disregard the existence of tropopause, *i.e.*, $\theta_T \rightarrow \infty$, (6.3) states that p_T' must vanish.

From the assumption that the tropopause is material, it follows that an air parcel should not depart from the surface; in other words, $\dot{\sigma}=0$ at $p=p_T$. We thus have

$$\omega_T = -\frac{\partial p_T'}{\partial t} + \mathbf{V}_T \cdot \nabla p_T'. \quad (6.4)$$

Now, let us consider the quasi-geostrophic vorticity equation in p -system which reads

$$\frac{\partial \zeta_g}{\partial t} + \mathbf{V}_g \cdot \nabla (\zeta_g + f) - f_0 \frac{\partial \omega}{\partial p} = 0, \quad (6.5)$$

where $\zeta_g = (g/f_0) \nabla^2 Z(x, y, p, t)$. In the following, we put $f = f_0 + \beta y$. f_0 is Coriolis parameter at 45°N and $\beta = \partial f / \partial y$ is assumed to be constant. It is known that assumption of β -plane becomes inaccurate as wave length tends to increase. Nevertheless we use it here for the sake of simplicity, so that meridional wave length should be small.

Inserting (6.1) into (6.5), integrating with respect to pressure from p_T to p_0 and linearizing under the assumption that

$$\bar{Z}(x, y, t) = \bar{Z}_D(x, t) - \frac{g}{f_0} \bar{U}y,$$

we get

$$d_1 \left(\frac{\partial}{\partial t} \frac{\partial^2}{\partial x^2} + \beta \frac{\partial}{\partial x} \right) \bar{Z}_D + d_2 \bar{U} \frac{\partial}{\partial x} \frac{\partial^2}{\partial x^2} \bar{Z}_D$$

$$+ d_3 \frac{\partial \bar{Z}_D}{\partial x} + f_0 \omega_T = 0, \quad (6.6)$$

where d_1 , d_2 and d_3 are combinations of a_1 ,

$a_2, a_3, p_0, p_T, g, f_0$ and $\partial \bar{p}_T / \partial y$ (see Appendix A).

From (6.3) and (6.4), we get

$$\begin{aligned} \omega_T &= \left(\frac{\partial}{\partial t} + \frac{1}{2} \bar{U} \frac{\partial}{\partial x} \right) p_T' \\ &= -\nu \left(\frac{\partial}{\partial t} + \frac{1}{2} \bar{U} \frac{\partial}{\partial x} \right) \bar{Z}_D, \end{aligned} \quad (6.7)$$

where ν is approximately 0.28 mb/m for parameters adopted here and we ignored the third term in the denominator of (6.3) considering an average state.

Therefore we finally have the following prognostic equation for the equivalent barotropic model from (6.6) and (6.7).

$$\begin{aligned} \left(\frac{\partial^2}{\partial x^2} - \mu^2 \right) \frac{\partial}{\partial t} \bar{Z}_D + \left(\beta + \gamma - \frac{\bar{U}}{2} \mu^2 \right) \frac{\partial}{\partial x} \bar{Z}_D \\ + \bar{U}^* \frac{\partial^3}{\partial x^3} \bar{Z}_D = 0, \end{aligned} \quad (6.8)$$

where $\mu^2 = (f_0/d_1)\nu \simeq 0.47 \times 10^{-12} \text{ m}^{-2}$, $\gamma = d_3/d_1 = (1/600) \cdot (f_0/2) \cdot (\partial \bar{p}_T / \partial y) \sim 0(10^{-12})$ in M. Gr. Sec. unit and $\bar{U}^* = (d_2/d_1) \bar{U} \simeq (2/3) \bar{U}$.

Equation (6.8) may be satisfied by a trigonometric wave solution of the form: $\bar{Z}_D \sim \exp ik(x - Ct)$, provided

$$C = \frac{\bar{U}^* k^2 + \frac{\bar{U}}{2} \mu^2 - \beta - \gamma}{k^2 + \mu^2} = \frac{\bar{U}^* + \frac{\bar{U}}{2} \frac{\mu^2}{k^2} - \frac{\beta}{k^2} - \frac{\gamma}{k^2}}{1 + \mu^2/k^2} \quad (6.9)$$

and $k = 2\pi/L$ (L is a wave length).

Except additional terms in the numerator, the formula of the phase velocity obtained here has the similar form to that may be available when we give an artificial mean divergence in the vorticity equation described in p -coordinate by Wiin-Nielsen (1959). Physical implications are, however, somewhat different from each other. We have not so far made any assumption about the vertical distribution of static stability in the troposphere as Wiin-Nielsen did, but we have taken into account the variation of the tropopause pressure. In other words, the motion of the ultra-long wave was treated in a self-adjusting system of the interaction between the stable stratosphere and the troposphere. The feasibility of the vertical transfer of kinetic energy through the tropopause simultaneously results

in the non-vanishing vertical mean divergence in the troposphere, which is indirectly controlled by the vertical distributions of static stability in both the stratosphere and the troposphere.

Mean meridional northward inclination of the tropopause pressure $\partial \bar{p}_T / \partial y$ acts just like as β -effect on phase speed of disturbance but γ seems to be less by one order of magnitude than β for the characteristic value of $\partial \bar{p}_T / \partial y$ in (6.9). As we can easily see in (6.9), phase speed C becomes finite in the limit, $k \rightarrow 0$ ($L \rightarrow \infty$), i.e., $C = -(\beta + \gamma - U_0 \mu^2/2)/\mu^2$. But if we disregard the existence of tropopause and make μ^2 to be zero, C becomes infinite and there will appear the rapid retrogression or the false westward movement of the very long wave.

(2) 2-level baroclinic model

Under the incorporation of a self-adjusting control of the ultra-long waves in a way similar to the one used in the barotropic case treated in the preceding section, we shall next solve the perturbation equations for the simple baroclinic model and observe whether or not the existence of the tropopause changes the phase velocity of disturbance and/or the instability criterion.

Fig. 5. Vertical resolution in a baroclinic model.

For the sake of simplicity we shall again assume that the tropopause pressure p_T may be expressed as $\bar{p}_T + p_T'$, where $\bar{p}_T$ is constant and $p_T' = p_T'(x, y, t)$, and a vertical p -velocity at $p = \bar{p}_T$ may be equal to ω_T in (6.7). These assumptions are rather crude but do not distort the essential feature of the present dis-

cussion. $\bar{p}_T=200$, $p_3=400$, $p_4=600$, $p_5=800$ and $p_0=1000$ mb surfaces are denoted with subscripts T , 3, 4, 5 and 0.

The vorticity equation for the mean flow $Z_m=(Z_3+Z_5)/2$ and the shear flow $Z_s=(Z_3-Z_5)/2$, and the thermal equation for level 4 may read

$$\left(\frac{\partial}{\partial t} + \mathbf{V}_m \cdot \nabla\right) \zeta_m + \mathbf{V}_s \cdot \nabla \zeta_s + \beta v_m + \frac{f_0^2}{2\Delta p \cdot g} \omega_T = 0 \quad (6.10)$$

$$\left(\frac{\partial}{\partial t} + \mathbf{V}_m \cdot \nabla\right) \zeta_s + \mathbf{V}_s \cdot \nabla \zeta_m + \beta v_s - \frac{f_0^2}{\Delta p \cdot g} \omega_4 + \frac{f_0^2}{2\Delta p \cdot g} \omega_T = 0 \quad (6.11)$$

$$\omega_4 = \frac{2g}{S\Delta p} \left(\frac{\partial}{\partial t} + \mathbf{V}_m \cdot \nabla\right) Z_s \quad (6.12)$$

and $\omega_T = -\nu \left(\frac{\partial}{\partial t} + \frac{1}{2} \mathbf{V}_m \cdot \nabla\right) Z_m$, where $U_m = \frac{1}{2}(U_3 + U_5)$, $U_s = \frac{1}{2}(U_3 - U_5)$, $U_4 \simeq U_m$ and S , the static stability, is constant.

Inserting wave solution into (6.10)–(6.12) and linearizing them, we obtain the frequency determining equation (see Appendix B), which gives the phase speed as

$$C = \frac{2+1}{2} \frac{\mu^2}{k^2} U_m - \frac{2+\frac{\mu^2}{k^2} + \frac{\varepsilon}{k^2}}{2(1+\mu^2/k^2)(1+\varepsilon/k^2)} \frac{\beta}{k^2} - \frac{\frac{k^2}{k^2}}{2(1+\mu^2/k^2)(1+\varepsilon/k^2)} U_s \pm \frac{\sqrt{D}}{2(1+\mu^2/k^2)(1+\varepsilon/k^2)}, \quad (6.13)$$

where

$$D = \left[\frac{1}{2} \left(\frac{\mu^2}{k^2} \right) \left(1 + \frac{\varepsilon}{k^2} \right) U_m + (\varepsilon - \mu^2) \frac{\beta}{k^4} \right]^2 - 4U_s^2 \left(1 + \frac{\mu^2}{k^2} \right) \left(\left(\frac{\varepsilon}{k^2} \right)^2 - 1 \right) - 2 \left(1 + \frac{\varepsilon}{k^2} \right) \left(\frac{\mu^2}{k^2} \right) \left(1 + \frac{1}{2} \frac{\mu^2}{k^2} \right) U_m U_s + 2U_s \left(\frac{\mu^2}{k^2} \right) \left(\frac{\beta}{k^2} \right) \left[2 + \left(\frac{\mu^2}{k^2} \right) + \left(\frac{\varepsilon}{k^2} \right) \right] + U_s^2 \left(\frac{\mu^2}{k^2} \right)^2, \quad (6.14)$$

$$\varepsilon = \frac{2f_0^2}{S\Delta p^2}, \text{ and } \mu^2 = \nu f_0^2 / (2\Delta p \cdot g).$$

In the case of $\omega_T \ll \omega_4$, we can ignore ω_T in (6.11) and then term including U_s in (6.13) and the third, fourth and fifth terms in the right hand side of (6.14) will disappear.

Again, results in the baroclinic case are formally parallel to those obtained in the case of artificial non-vanishing mean divergence, for example, solutions by Wiin-Nielsen (1959). Therefore we will briefly observe results. In (6.13), if we make $k \rightarrow 0$ ($L \rightarrow \infty$) to see how the phase velocity becomes for very long wave, we obtain

$$\lim_{k \rightarrow 0} C = \begin{cases} \frac{5}{4} U_m - \frac{\beta}{\varepsilon} & \text{for plus sign,} \\ \frac{1}{4} U_m - \frac{\beta}{\mu^2} & \text{for minus sign.} \end{cases} \quad (6.15)$$

It should be noted that in (6.16) the ultra-long wave in the baroclinic case again has finite phase speed if the variation of tropopause pressure is incorporated. Appearance of $\frac{1}{4} U_m$ in (6.16) is not essential, for it depends on the functional form of $A(p)$.

For a baroclinic unstable wave, the assumption $\omega_T \ll \omega_4$ is permissible and the instability condition $D < 0$ reduces to

$$4U_s^2 \left(1 + \frac{\mu^2}{(2\pi)^2} L^2 \right) \left(\frac{\varepsilon^2}{(2\pi)^4} L^4 - 1 \right) > \left[\frac{1}{2} \left(\frac{\mu^2}{(2\pi)^2} L^2 \right) \left(1 + \frac{\varepsilon}{(2\pi)^2} L^2 \right) U_m + \frac{(\varepsilon - \mu^2)\beta}{(2\pi)^4} L^4 \right]^2. \quad (6.17)$$

In the case of $\mu^2=0$, the mean zonal velocity U_m is irrelevant of the instability condition. In order to satisfy (6.17) the wave length of an unstable wave L^* must be

$$L^* > \sqrt{\frac{2\pi}{\varepsilon}},$$

where for $S=2.5 \times 10^{-2} \text{ m}^2 \text{ sec}^{-2} \text{ mb}^{-2}$, $L^*=2810$ km and for $S=5 \times 10^{-2} \text{ m}^2 \text{ sec}^{-2} \text{ mb}^{-2}$, $L^*=3966$ km.

From (6.17) we can see that,

- (a) if U_m happens to be zero,
  - (1) waves become unstable for smaller U_s than for that of non-zero U_m and
  - (2) for a fixed wave length, U_s^* which is the minimum wind shear necessary for

baroclinic instability becomes smaller for $\mu^2 \neq 0$ than that for $\mu^2 = 0$.

Also we observe that,

(b) if $\mu^2 \neq 0$, the larger the value of U_m , the larger the value of U_s^* and that,

(c) if $\mu^2 = 0$, U_s^* is smaller than that for $\mu^2 \neq 0$ for wave lengths less than a certain critical value which depends on U_m , on the other hand, for wave length beyond the critical value U_s^* is smaller in the case of $\mu^2 \neq 0$ than that of $\mu^2 = 0$.

7. Conclusions

In order to incorporate the effect of the variable tropopause and of the stratosphere and to examine the influence on the motion of tropospheric disturbance, a computational atmospheric model is designed and its properties are investigated. The tropopause is treated as a material surface.

The energy equation shows that the vertical transfer of kinetic energy through the tropopause is expressed in the terms of $(\overline{\phi_T'}/\bar{g}) \cdot \partial p_T' / \partial t$ which makes it simple to estimate the transport of energy from observational data. Heuristic consideration on this term reveals that the amplifying long wave, especially stationary wave, plays more important role in the strato-tropospheric interaction than the short wave.

Using aerological materials presented by Defant and Taba (1958), we computed $(\overline{\phi_T'}/\bar{g}) \cdot \partial p_T' / \partial t$. The results indicate strong interaction through kinetic energy between the troposphere and the stratosphere, though the term of computation is rather short.

σ -coordinate system is not convenient to treat in the form of vorticity and divergence equation and also complicated in the process of filtering meteorological noises. So we solved the quasi-geostrophic equations expressed with p -coordinate which retains the physical characteristics of σ -system, and obtained wave solutions of linearized forms of prognostic equations in the barotropic and baroclinic model. The resulting prognostic equations are formally similar to those obtained with respect to the equations of Cressman's model. This model contains the non-vanishing mean divergence and is favorable to suppress the westward spurious retrogression of the ultralong wave in the troposphere. The existence

of the vertical transfer of kinetic energy through the tropopause is important because it is related to the non-vanishing vertical mean divergence in the troposphere, which is indirectly controlled by the vertical distribution of static stability in both the stratosphere and the troposphere.

Hence we obtain the formula of the phase velocity of wave and curves of baroclinic instability which are modified by the existence of the stratosphere and similar to those obtained by Wiin-Nielsen (1959). Generally speaking the baroclinic instability criterion is shifted in the same sense as β -effect is weakened.

The numerical prediction model described in σ -coordinate system presented in this paper is favorable to take into account the variation of tropopause pressure and the existence of the stable stratosphere for forecasting the motion of the tropospheric disturbance.

Acknowledgments

The author wishes to express his hearty thanks to Prof. S. Syōno, Tokyo University, for his constant encouragement throughout this work. The author is also greatly indebted to Dr. A. Arakawa, Electronic Computation Center of Japan Meteorological Agency, for his very useful suggestions and constructive criticism, to staff members of our Center and particularly to Dr. K. Gambo for their kind encouragements.

It is a pleasure to thank Miss K. Endo for typewriting the manuscript and Miss Y. Atsuki for drafting figures.

Appendix

A. Derivation of the linearized equation (6.6)

Inserting (6.1) into (6.5) and integrating with respect to pressure from p_T to p_0 , we obtain

$$\begin{aligned} & \left(\frac{\partial \bar{\zeta}_g}{\partial t} + \left(\frac{\partial f}{\partial y} \right) \bar{v}_g \right) \int_{p_T}^{p_0} A(p) dp \\ & + (\bar{V}_g \cdot \nabla \bar{\zeta}_g) \int_{p_T}^{p_0} A^2(p) dp + f_0 \omega_T \\ & + A(p_T) \left(\bar{\zeta}_g \frac{\partial p_T}{\partial t} + \bar{v}_g f \frac{\partial p_T}{\partial y} \right) \\ & + A^2(p_T) (\bar{V}_g \cdot \bar{\zeta}_g \nabla p_T) = 0 \end{aligned} \quad (\text{A.1})$$

where $\omega = 0$ at $p = p_0$ is used. Although equation (5.7) is described in p -coordinate system,

it includes the physical effect of the tropopause and is essentially equivalent to the vorticity equation for the large scale motion in σ -coordinate system. If we linearize (A.1) under the assumption that

$$\bar{Z}(x, y, t) = \bar{Z}_D(x, t) - \frac{g}{f_0} \bar{U}_y,$$

and take into account the functional form for $A(p)$, (A.1) is reduced to (6.6), i.e.,

$$d_1 \left(\frac{\partial}{\partial t} \frac{\partial^2}{\partial x^2} + \beta \frac{\partial}{\partial x} \right) \bar{Z}_D + d_2 \bar{U} \frac{\partial}{\partial x} \frac{\partial^2}{\partial x^2} \bar{Z}_D + d_3 \frac{\partial \bar{Z}_D}{\partial x} + f_0 \omega_T = 0, \quad (6.6)$$

where

$$\begin{aligned} d_1 &= \frac{g}{f_0} \left[\frac{a_1}{3} (p_0^3 - \bar{p}_T^3) + \frac{a_2}{2} (p_0^2 - \bar{p}_T^2) + a_3 (p_0 - \bar{p}_T) \right] \simeq \frac{g}{f_0} \times 600 \text{ mb}, \\ d_2 &= \frac{g}{f_0} \left[\frac{a_1^2}{5} (p_0^5 - \bar{p}_T^5) + \frac{a_1 a_2}{2} (p_0^4 - \bar{p}_T^4) + \frac{1}{3} (a_2^2 + 2a_1 a_3) (p_0^3 - \bar{p}_T^3) + a_2 a_3 (p_0^2 - \bar{p}_T^2) + a_3^2 (p_0 - \bar{p}_T) \right] \simeq \frac{g}{f_0} \times 400 \text{ mb}, \\ d_3 &= \frac{g}{f_0} \left(\frac{f_0}{2} \frac{\partial \bar{p}_T}{\partial y} \right). \end{aligned}$$

B. Derivation of the frequency determining equation

Inserting wavy solution

$$Z_m = -\frac{f_0}{g} U_m y + \hat{Z}_m e^{ik(x-Ct)} \quad (B.1)$$

and

$$Z_s = -\frac{f_0}{g} U_s y + \hat{Z}_s e^{ik(x-Ct+\delta)} \quad (B.2)$$

into (6.10)–(6.12) and linearizing them, we obtain the frequency determining equation

$$\begin{aligned} & (1 + \mu^2/k^2) (1 + \varepsilon/k^2) C^2 + [(1 + \mu^2/k^2) (\beta/k^2 - U_m - (\varepsilon/k^2) U_m) + (1 + \varepsilon/k^2) (\beta/k^2 - U_m - (\mu^2/2k^2) U_m) + U_s (\mu^2/k^2)] C + [(\beta/k^2 - U_m - (\mu^2/2k^2) U_m) \times (\beta/k^2 - U_m - (\varepsilon/k^2) U_m) + U_s^2 (\varepsilon/k^2 - 1) - (U_m U_s/2) (\mu^2/k^2)] = 0. \end{aligned} \quad (B.3)$$

C. Treatment of (6.10)–(6.12) as an initial value problem

It is of interest to treat (6.10)–(6.12) as an initial value problem in a way similar to the one presented by Ogura (1957). Assuming the form of particular solution of (6.10)–(6.12) as

$$Z_3(x, y, t) = -\frac{f_0}{g} U_3 y + \alpha_3(t) e^{ikx + i\delta_3(t)} \quad (C.1)$$

$$Z_5(x, y, t) = -\frac{f_0}{g} U_5 y + \alpha_5(t) e^{ikx + i\delta_5(t)} \quad (C.2)$$

we can obtain the initial tendency for the amplification and the phase velocity at level 3 and 5 as follows.

$$\begin{aligned} \frac{d\alpha_3}{dt} &= -\frac{k\alpha_5}{(\gamma+2)(1+\mu^*/k^2)} \left[\frac{\beta}{k^2} + V - \left(\frac{\mu^2}{\varepsilon} \right) \times \left(\frac{\gamma+1}{\gamma} \frac{U_m}{2} + \frac{\beta}{k^2} - U_5 - \frac{U_3}{\gamma} \right) \right] \\ &\times \sin(\delta_5 - \delta_3), \end{aligned} \quad (C.3)$$

$$\begin{aligned} C_3 &= -\frac{1}{k} \frac{d\delta_3}{dt} = \frac{U_3 - \beta/k^2}{1 + \mu^*/k^2} - \frac{1}{(\gamma+2)(1+\mu^*/k^2)} \\ &\times \left[V \left\{ 1 + \frac{\alpha_5}{\alpha_3} \cos(\delta_5 - \delta_3) \right\} - \frac{\beta}{k^2} \left\{ 1 - \frac{\alpha_5}{\alpha_3} \cos(\delta_5 - \delta_3) \right\} - \left(\frac{\mu^2}{\varepsilon} \right) \times \left\{ \left(\frac{\gamma+1}{\gamma} \frac{U_m}{2} + \frac{U_5}{\gamma} \right) + \frac{\alpha_5}{\alpha_3} \left(\frac{\gamma+1}{\gamma} \frac{U_m}{2} + \frac{\beta}{k^2} - U_5 - \frac{U_3}{\gamma} \right) \cos(\delta_5 - \delta_3) \right\} \right], \end{aligned} \quad (C.4)$$

$$\begin{aligned} \frac{d\alpha_5}{dt} &= -\frac{k}{\gamma+2} \alpha_3 \left[V - \frac{\beta}{k^2} + \frac{\mu^2/\varepsilon}{\gamma(1+\mu^*/k^2)} \right] \\ &\times \left(\frac{U_m}{2} - U_3 + \frac{\beta}{k^2} \right) \sin(\delta_5 - \delta_3), \end{aligned} \quad (C.5)$$

$$\begin{aligned} C_5 &= -\frac{1}{k} \frac{d\delta_5}{dt} = \frac{(\gamma+1)(1+\mu^*/k^2) + 1}{(\gamma+2)(1+\mu^*/k^2)} \left(U_5 - \frac{\beta}{k^2} \right) \\ &+ \left[\left\{ \left(\frac{U_3}{1+\mu^*/k^2} - U_5 \right) + V \frac{\alpha_3}{\alpha_5} \cos(\delta_3 - \delta_5) \right\} + \frac{\beta}{k^2} \left\{ 1 - \frac{\alpha_3}{\alpha_5} \cos(\delta_3 - \delta_5) \right\} + \frac{\mu^2/\varepsilon}{\gamma(1+\mu^*/k^2)} \right] \\ &\times \left\{ \left(\frac{U_m}{2} - U_3 \right) + \left(\frac{U_m}{2} - U_3 + \frac{\beta}{k^2} \right) \times \frac{\alpha_3}{\alpha_5} \cos(\delta_3 - \delta_5) \right\}, \end{aligned} \quad (C.6)$$

where $\mu^* = \mu^2/2$, $\mu^2 = \nu f_0^2 / (2\Delta p \cdot g)$, $\varepsilon = 2f_0^2 / (S\Delta p^2)$, $\gamma = 2k^2/\varepsilon$ and $V = U_3 - U_5$. If $\mu^2 = 0$, (C.3) and (C.4) become the same as those derived by

Ogura. In barotropic case, *i.e.*, $\gamma \rightarrow \infty$, we get

$$\frac{d\alpha_3}{dt} = 0$$

and

$$C_3 = \frac{U_3 - \beta/k^2}{1 + \mu^*/k^2}$$

The latter is same as (6.9). Equation (C.3) and (C.4) state that the initial tendency for the amplification at upper level is reduced by a factor μ^*/k^2 (>0) in the denominator, but at lower level the effect of μ^*/k^2 varies and decreases. Physical interpretation of baroclinicity appeared in (C.3)–(C.6) may not be needed to be explained here. Concerning the phase speed, control of the very long wave at upper level is clearly shown in (C.4) but, at lower level, we observe in (C.6) that control is not enough at least as far as the initial trend is concerned. As to the mean phase velocity in the whole troposphere, the control of ultra-long wave can be performed as is shown in (6.16). In our analytical argument we have disregarded the pressure change with respect to time on the ground surface since the magnitude of the change is smaller than that at tropopause. In the exact treatment of σ -system, this pressure variation on the ground should be incorporated and may contribute to suppress the false retrogression, especially in the lower part of the troposphere.

References

- Burger, A.P., 1958: Scale consideration of planetary motions of the atmosphere. *Tellus*, **10**, 196–205.
- Charney, J.G. and P.G. Drazin, 1961: Propagation of planetary-scale disturbances from the lower into the upper atmosphere. *J. geophys. Res.*, **66**, 83–109.
- Cressman, G.P., 1958: Barotropic divergence and very long atmospheric waves. *Mon. Wea. Rev.*, **86**, 293–297.
- Eliassen, A., 1960: On the use of a material layer model of the atmosphere in numerical prediction. *Proceedings of international symposium on numerical weather prediction in Tokyo*. 207–212.
- Gates, W.L., 1957: A dynamical model for large-scale tropospheric and stratospheric motions. *Quart. J. Roy. meteor. Soc.*, **83**, 141–160.
- Murakami, T., 1964: Multiple-level model for the p - and s -coordinates. *J. meteor. Soc. Japan*, **42**, 1–13.
- Ogura, Y., 1957: Wave solutions of the vorticity equation for the $2\frac{1}{2}$ -dimensional model. *J. meteor.*, **14**, 60–64.
- Defant, Fr. and H. Taba, 1957: The threefold structure of the atmosphere and the characteristics of the tropopause. *Tellus*, **9**, 259–274.
- , 1958: The strong index change period from January 1 to January 7, 1956. *Tellus*, **10**, 225–242.
- , 1958: The break down of zonal circulation during the period January 8 to 13, 1956, the characteristics of temperature field and tropopause and its relation to the atmospheric field of motion. *Tellus*, **10**, 430–450.
- Wiin-Nielsen, A., 1959: On barotropic and baroclinic models, with special emphasis on ultra-long waves. *Mon. Wea. Rev.*, **87**, 171–183.

Burger, A.P., 1958: Scale consideration of plane-

時間的に変動する圏界面の力学的効果

新 田 尚*

気象庁電子計算室

大規模運動に対する大気モデルを成層圏まで拡張することを考え、圏界面と地表面を座標に含む、次のような σ 座標を鉛直方向にもつ x, y, σ -系を採用した。即ち、

$$\sigma = \varepsilon \frac{p}{p_T} : \text{成層圏}, \quad \sigma = (1 - \varepsilon) \frac{p - p_T}{p_S - p_T} + \varepsilon : \text{対流圏}.$$

* 現在：米気象局国立気象センター，米国メリーランド州ストランド

ここに ε は $0 < \varepsilon < 1$ なる定数で圏界面の座標を示すから、層の鉛直方向のわけ方によってきめられるべきものである。(例えば 2 層モデル (成層圏と対流圏) に対しては $\varepsilon = 1/2$, 3 層モデルに対しては成層圏 1 層, 対流圏 2 層なら $\varepsilon = 1/3$ がよい。) 従ってこの σ 座標では $\sigma = \varepsilon$ が内部境界面となる。

このモデルは対流圏と成層圏との力学的相互作用を扱うのに便利にできている。基本的仮定として圏界面は物質面であるとし、その結果内部境界面としての圏界面 ($\sigma = \varepsilon$) では $d\sigma/dt = 0$ とおいた。基礎運動方程式と熱の式及びこれから導かれるエネルギー方程式を示した。成層圏と対流圏との相互作用をあらわす運動エネルギー交換の項 $\overline{(\phi_T'/g) \cdot \partial p_T'/\partial t}$ (ϕ_T' , p_T' はそれぞれ圏界面のジオポテンシャルと気圧) の役割を簡単な $\sin$ 型の波について調べた。短期間の、対流圏と下部成層圏との運動エネルギーのやりとりには、発達期にある長波の谷、峯が重要な役割を果していることがわかった。

更に実況についてみるために、Defant と Taba (1958) の解析結果から $\overline{(\phi_T'/g) \cdot \partial p_T'/\partial t}$ の値を計算した。限られた資料から一般的な結論は出せないが、その日々の値は、 $10 \text{ joule m}^{-2} \text{ sec}^{-1}$ のオーダーで、その向きは擾乱が発達しているときは上向きに、衰弱しているときは下向きにとはげしく変る。従って時間平均してみると $1 \text{ joule m}^{-2} \text{ sec}^{-1}$ のオーダーになる。

圏界面の気圧の時間変動を許すことは、同時に、対流圏全体について鉛直積分しても消えない発散をもたらすことになるから、このようなモデルで記述される対流圏の擾乱は、Cressman (1958) のモデルの場合と同じ特性をもつ。即ち、超長波のみかけ上の西進がおさえられ、パロクリニックの場合には、更に不安定度曲線があたかも β -効果が減じたのと同じように変る。このことを線型化した場合について示した。