NOTES AND CORRESPONDENCE

Observational Implication for the Tropospheric NO₂ Burden

By K. Shibasaki and T. Ogawa

*Geophysics Research Laboratory, University of Tokyo, Bunkyo-ku, Tokyo 113**(Manuscript received 10 October 1980, in revised form 9 February 1981)*

We have developed a spectroscopic technique to measure the atmospheric NO₂ burden. It is based upon the NO₂ absorption of the solar radiation in the wavelength regime 430-450 nm, and was applied for a balloon sunset observation to obtain the vertical profile of the stratospheric NO₂ (Ogawa *et al.*, 1981). From the same balloon flight, we got a control spectrum which is free from absorption by the atmospheric NO₂ and is essential in our data analysis. Since we could recover the spectrometer flown on the balloon, we wanted to use it for the ground observation.

The tropospheric NO₂ concentration is of considerable interest with respect to the global diffusion of the polluted air and the natural cycle of odd nitrogen compounds. Recent observations by Noxon (1978, 1980) show the tropospheric NO₂ burden is less than $1 \times 10^{15} \text{ cm}^{-2}$ in the natural background. If a constant mixing ratio is applicable for the height distribution of the tropospheric NO₂, then the vertical column density $1 \times 10^{15} \text{ cm}^{-2}$ is equivalent to a mixing ratio of 0.05 ppbv. We are interested in elucidating whether his result is also the case in our country.

The sunrise or sunset observation which looks at the sun along a slant path in the atmosphere is necessary to detect the NO₂ absorption in the solar spectra, unless the atmosphere is highly polluted with NO₂. Consequently, a mountain or seaside is chosen for the observing site. One observation was made at Mt. Norikura (location: 36°07'N, 137°33'E; elevation: 2876 m) on August 8/9, 1979. The other was at the tracking station of Sanriku Balloon Center (location: 39°10'N, 141°49'E; elevation: 450 m) on June 3 through 7, 1980.

Figure 1 represents the apparent variation of the NO₂ column density with solar zenith angle as observed at Mt. Norikura. The procedure to

deduce the column density was the same as by Ogawa *et al.* (1981). The morning data points are dispersed because of a low signal to noise ratio, but it is evident that they are smaller than the evening ones. This phenomenon is already known and explained by the diurnal variation of the stratospheric NO₂: *i.e.*, conversion of NO₂ to N₂O₅ at night (*e.g.*, Noxon *et al.*, 1979).

The solid curve in Figure 1 is the column density as calculated from the stratospheric NO₂ profile obtained in our balloon sunset observation on May 29, 1978. According to Noxon (1979), the monthly average of the atmospheric NO₂ burden changes little in May through August, so that our balloon profile may be applicable to these ground-based observation. The agreement between the calculation and the observation seems good. This infers almost all the observed column density comes from the stratosphere. Accepting the maximum probable errors, we can take the

Fig. 1 Apparent variation of the NO₂ column density with solar zenith angle as observed at Mt. Norikura on August 8/9, 1979. Solid curve is for a model NO₂ distribution only in the stratosphere, and dashed curve for a model with a tropospheric mixing ratio of 0.1 ppbv together with the stratosphere.

model with a tropospheric mixing ratio of 0.1 ppbv together with the stratospheric distribution. Therefore, it may be suggested that the tropospheric mixing ratio is less than 0.1 ppbv at this observation. A constant mixing ratio of 0.1 ppbv is equivalent to a vertical column density of $1 \times 10^{15} \text{ cm}^{-2}$. Note that our present observation was made at a height of 2876 m. This particular observation was made above the boundary layer, and must have been free from the pollution. Again, this value may be reasonable for the natural troposphere, as it agrees well with Noxon's.

The same analysis was made for the observation at Sanriku Balloon Center. An upper limit of 0.2 ppbv was obtained. Because the optical condition was not good in this observation, the limit was larger than the previous one. Kawamura and Fushimi (1978) measured the surface NO_2 concentration at the adjacent places to obtain a background concentration of about 0.5 ppbv both at Mt. Norikura and Ryori Meteorological Rocket Station which is 14 km south of Sanriku Balloon Center. The agreement between their and our results is not bad, in view of the fact that the tropospheric NO_2 is highly variable and that the experimental uncertainty is appreciable.

In conclusion, our present analysis infers that the tropospheric NO_2 concentration is likely to be very small compared to that of the stratosphere: a few tenths of ppbv or less. However, this value is not always applicable for the atmosphere in rural Japan. Kawamura and Fushimi detected NO_2 rich air mass with a few ppbv during their observation at Ryori. According to our continual observations at GRL Kakioka Field Station, which is 70 km northeast from Tokyo,

the tropospheric concentration could become as low as the natural background, but sometimes it became as high as those in Tokyo. Further observations should be necessary to depict the average feature and variability of the tropospheric NO_2 .

Acknowledgements

The authors wish to thank Professor F. Moriyama, Tokyo Astronomical Observatory, University of Tokyo, and Professor J. Nishimura, Institute of Space and Aeronautical Science, University of Tokyo, for affording us the observational facilities of Norikura Corona Observatory and Sanriku Balloon Center, respectively. This research was financially supported by the Ministry of Education, Science and Culture under Grant-in-Aid No. 454105.

References

- Kawamura, K. and K. Fushimi, 1978: On the concentrations of atmospheric SO_2 , NO_2 , NO , CO and O_3 as observed at Ryori, Mt. Norikura, Minamitorishima Is. and Chichijima Is. *Tech. Rep. Meteorolog. Res. Inst. Japan*, 117-123.
- Noxon, J. F., 1978: Tropospheric NO_2 . *J. Geophys. Res.*, **83**, 3051-3057.
- , 1979: Stratospheric NO_2 , 2. Global behavior. *J. Geophys. Res.*, **84**, 5067-5076.
- , 1980: Correction. *J. Geophys. Res.*, **85**, 4560-4561.
- , E. C. Whipple and R. S. Hyde, 1979: Stratospheric NO_2 , 1. Observational method and behavior at mid-latitude. *J. Geophys. Res.*, **84**, 5047-5065.
- Ogawa T., K. Shibasaki and K. Suzuki, 1981: Balloon observation of the stratospheric NO_2 profile by visible absorption spectroscopy. *J. Met. Soc. Japan*, **59**, 412-418.

対流圏 NO_2 全量について

柴 崎 和 夫, 小 川 利 紘

東京大学理学部地球物理研究施設