—Original—

Evaluation of Glycosaminoglycans Levels in Normal Joint Fluid of the Knee

Yoshihito Nakayama, Yasumasa Shirai, Kiyoshi Yoshihara and Shinji Uesaka

Department of Orthopaedic Surgery, Nippon Medical School

Abstract

Glycosaminoglycans (GAGs) level in joint fluid have been investigated in various joint diseases as a joint maker. However, there are very few studies of normal joint fluid providing a baseline for a better understanding of altered GAGs level in pathological joint fluid. We investigated GAGs level in knee joint fluid for 25 healthy young volunteers with a mean age of 27.5 years (range: 18 to 36 years). Biochemical evaluations included the concentration of hyaluronic acid (HA), chondroitin 4-sulfate (C4S) and chondroitin 6-sulfate (C6S), and the C6S/C4S ratio. The unsaturated disaccharides derived from HA by Morgan-Elson methods, and those from CS were measured with high performance liquid chromatography. The mean HA concentration was 3.4 ± 0.6 mg/ml. The mean concentration of C4S and C6S, and C6S/C4S ratio were 19.0 ± 4.9 nmol/ml, 125.5 ± 44.4 nmol/ml and 6.5 ± 1.1 respectively. (J Nippon Med Sch 2000; 67: 92–95)

Key words: hyaluronic acid, chondroitin sulfate, normal joint fluid, biochemical analysis

Introduction

Glycosaminoglycans (GAGs) in joint fluid consist of hyaluronic acid (HA), chondroitin sulfate (CS) and keratan sulfate (KS) isomers. These GAGs have an important role in the joint lubrication and cartilaginous nutrition. Recently their concentrations have been thought to reflect joint tissue metabolism in normal and diseased states. GAGs levels have been investigated in various joint diseases¹⁻³, and considerable interest has been developing as a joint marker for rheumatoid arthritis (RA) and osteoarthritis (OA).

However, there have been some debates^{4,5} about whether GAGs levels in joint fluid increase or decrease in RA and OA, compared with those in normal subjects, because there are few studies of normal joint fluid. Our review of current published studies on nor-

mal joint fluid reveals the following limitations: studies are small, few description regarding to subject selection, inadequate study population and discrepancy between the results of HA and CS concentration. To our knowledge, is no detailed study evaluating normal GAGs levels, especially CS levels in joint fluid. The aim of this study is to clarify normal GAGs levels of joint fluid.

Materials and Methods

We examined joint fluid from healthy young volunteers with no history of knee symptoms and injuries. Moreover athlete participating high level sports was excluded from the present study. Sufficient volumes of joint fluid for assay (0.2 ml) were obtained from 25 of 59 volunteers. The mean age of these 25 volunteers was 27.5 years (range: 18 to 36 years).

Corresponding to, Yoshihito Nakayama M.D., Department of Orthopaedic Surgery, Nippon Medical School, 1-1-5 Sendagi, Bunkyo-ku, Tokyo 113-8603, Japan

Journal Website (<http://www.nms.ac.jp/jnms/>)

Table 1 The results of glycosaminoglycans levels of synovial fluid in 25 healthy young volunteers

Case	Age	Sex	Volume	HA	C4S	C6S	C6S/C4S
			ml	mg/ml	nmol/ml	nmol/ml	
1	26	M	0.5	3.3	15.2	75.8	5.0
2	31	F	0.5	4.4	16.3	100.7	6.2
3	24	M	0.5	3.5	15.7	86.7	5.5
4	36	M	0.5	4.2	19.1	102.2	5.4
5	18	F	0.8	3.9	21.7	157.6	7.3
6	35	M	1.0	2.6	16.9	134.4	7.7
7	35	M	0.3	3.6	25.2	139.7	5.5
8	26	F	0.5	3.4	18.1	80.3	4.4
9	34	M	0.4	3.1	12.0	86.0	7.2
10	35	F	0.8	3.9	14.6	96.4	6.6
11	26	F	0.6	3.6	14.1	92.0	6.5
12	21	F	1.0	3.2	12.7	83.8	6.6
13	26	M	0.3	3.4	18.4	134.3	7.3
14	20	M	0.6	3.3	16.6	104.0	6.3
15	22	M	1.0	3.8	20.9	153.1	7.3
16	29	M	0.9	3.0	16.7	108.6	6.5
17	22	M	1.3	3.5	29.7	214.4	7.2
18	25	M	0.3	2.5	26.0	181.4	7.0
19	28	M	0.3	2.7	22.9	219.1	9.6
20	26	M	1.1	2.0	15.8	102.6	6.5
21	28	F	0.3	3.8	16.4	99.4	6.1
22	23	M	0.4	4.4	27.7	218.3	7.9
23	29	F	0.3	3.4	17.6	107.6	6.1
24	32	M	0.6	3.9	27.8	161.5	5.8
25	31	F	0.2	3.2	16.9	97.0	5.7

M : male, F : female, HA : hyalunonic acid, C4S : chondroitin 4-sulfate, C6S : chondroitin 6-sulfate, C6S/C4S : a ratio of C6S to C4S

Joint fluid samples were collected in sterile plastic tubes and centrifuged at $10,000 \times g$ for 15 minutes at 4°C to remove cells and joint debris. The supernatants were stored at -80°C until biochemical assay. The unsaturated disaccharides derived from hyaluronic acid (HA) by Morgan-Elson methods⁶, and those from CS were measured by high performance liquid chromatography (HPLC)⁷. Biochemical evaluations included the concentration of HA, chondroitin 4-sulfate (C4S), and chondroitin 6-sulfate (C6S), and C6S/C4S ratio.

Results

The results of GAGs levels of joint fluid in 25 healthy volunteers were shown in **Table 1**. The mean volume of joint fluid was $0.6 \pm 0.3 \text{ ml}$ (range: 0.2 to 1.3 ml). The mean HA concentration was $3.4 \pm 0.6 \text{ mg/ml}$ (range: 2.0 to 4.4 mg/ml). The mean concentration of C4S and C6S were $19.0 \pm 4.9 \text{ nmol/ml}$ (range: 12.0 to

29.7 nmol/ml) and $125.5 \pm 44.4 \text{ nmol/ml}$ (range: 75.8 to 219.1 nmol/ml) respectively. The mean C6S/C4S ratio was 6.5 ± 1.1 (range: 4.4 to 9.6).

Discussion

The determination of GAGs in human joint fluid has been difficult due to diversity of structures and limited amount of unsaturated disaccharides. Recently the development of HPLC allows more specific and sensitive determination of HA and CS isomers derived from GAG chains^{8,9}.

HA is a main component of synovium and joint fluid. The viscoelastic property of joint fluid is mainly due to HA with a high capacity for binding water. Detection and quantitation of HA is of interest in many fields, especially in clinical chemistry. Recent studies demonstrated that HA has anti-inflammatory functions including inhibition of prostaglandin E2 synthesis^{10,11},

suppression of leucocyte migration and proliferation¹², and protection against cytotoxicity caused by reactive oxygen species¹³.

In normal condition, HA concentration in joint fluid is maintained at a higher level, comparing with blood level. In inflammatory diseases¹⁻³ such as RA, gout and traumatic arthritis, HA concentration has been reported to be lower than normal. Balazs et al.¹⁴ stated that a considerable variation was found in HA concentration of joint fluid collected from various joints. They reported that HA concentration in joint fluid of the knee was 2.5~3.8 mg/ml in normal subjects aged 18 to 35 years. Our results for HA concentration agree well with their study.

CS contain a repeating disaccharides of N-acetyl-galactosamine and glucuronate. In aggrecan¹⁵, most galactosamine residues carry a single sulphate group in 4 or 6 position, and a small portion are non-sulphated. Articular cartilage are rich in CS¹⁶, especially in C6S. Bayliss et al.¹⁷ reported that C6S makes up 72~95% of CS in normal articular cartilage. C4S is distributed widely not only in articular cartilage, but also in synovium and ligament. Shinmei et al.¹⁸ stated that C6S in joint fluid derives mostly from articular cartilage because of its large tissues volume in the joint, and that C6S concentration and C6S/C4S ratio may reflect cartilage metabolism.

Asari et al.¹⁹ reported in a study of 6 young volunteers that the mean concentrations of C4S and C6S, C6S/C4S ratio were 11.8 ± 1.98 nmol/ml, 58.0 ± 19.5 nmol/ml and 4.9 respectively. However no subject selection and detailed back ground data of study population were described in their study. Namiki²⁰ noted in a study of 12 young athletes that the mean concentrations of C4S and C6S, C6S/C4S ratio and HA concentration were $14.8 \pm$ nmol/ml, 63.3 ± 19.9 nmol/ml, 5.5 ± 3.5 and 2.7 ± 0.7 mg/ml respectively. These authors stated that questions remain about whether athletes are adequate for study population, and found that the results of HA concentration was lower compared with a study of Balazs et al. In comparison with their results, our results showed higher values for all three parameters. The reason for the discrepancy between these results is unclear, but may be mainly differences in subject selection or study population.

Change in CS concentration of joint fluid have been

reported in various joint diseases. In recent studies, C4S and C6S concentrations and C6S/C4S ratio in RA^{21,22} were reported to be 26.4~29.7 nmol/ml, 28.9~43.2 nmol/ml and 1.1~1.9 respectively, and those in OA²³⁻²⁵ 15.4~20.4 nmol/ml, 56.8~81.1 nmol/ml and 3.8~4.0. Judging from normal level obtained in the present study, in RA, C4S concentration considerably increased and C6S concentration and C6S/C4S ratio markedly decreased. In contrast, in OA, C4S concentration is similar to normal and C6S concentration C6S/C4S ratio are considerably decreased. Shinmei et al.²⁶ noted that the proliferated synovium may accelerate the production and release of C4S into joint fluid, causing increased C4S concentration in RA, and loss of articular cartilage or an altered cartilage metabolism in RA and OA relate to decreased C6S concentration.

The present study demonstrates that GAGs level in normal joint fluid. Measurements of GAGs levels in joint fluid may provide useful information in monitoring joint tissue metabolism, and in assessing severity and the prognosis of joint diseases.

References

1. Lohmander LS: Markers of cartilage metabolism in arthrosis: a review. *Acta Orthop Scand* 1991; 62: 623-632.
2. Caterson B, Hughes CE, Roughley P, Mort JS: Anabolic and catabolic markers of proteoglycan metabolism in osteoarthritis. *Acta Orthop Scand* 1995; 66: 121-124.
3. Hardingham T: Changes in chondroitin sulphate structure induced by joint disease. *Acta Orthop Scand* 1995; 66: 107-110.
4. Caterson B, Mahmoodian F, Sorrell JM, Hardingham TE, Bayliss MT, Carney SL: Modulation of native chondroitin sulphate structure in tissue development and in disease. *Journal Cell Science* 1990; 97: 411-417.
5. Brown MP, West LA, Merritt KA, Plaas AHK: Changes in sulfation patterns of chondroitin sulfate in equine articular cartilage and synovial fluid in response to aging and osteoarthritis. *AJVR* 1998; 59: 786-791.
6. Huang Y, Toyoda H, Toida T, Imanari T: Determination of chondroitin sulphates in human whole blood, plasma and blood cells by high-performance liquid chromatography. *Biomedical Chromatography* 1995; 9: 102-105.
7. Bayliss MT, Davidson C, Woodhouse SM, Osborne DJ: Chondroitin sulphation in human joint tissues varies with age, zone and topography. *Acta Orthop Scand*

- 1995; 66: 22–25.
8. Sandy JD, Flannery CR, Neame PJ, Lohmander LS: The structure of aggrecan fragments in human synovial fluid. *J Clin Invest* 1992; 89: 1512–1516.
9. Mourao PAS: Distribution of chondroitin 4-sulfate and chondroitin 6-sulfate in human articular and growth cartilage. *Arthritis Rheum* 1988; 31: 1028–1033.
10. Murakami K, Shibata T, Kubota E, Maeda H: Intra-articular levels of prostaglandin E₂, hyaluronic acid, and chondroitin-4 and -6 sulfates in the temporomandibular joint synovial fluid of patients with internal derangement. *J Oral Maxillofac Surg* 1998; 56: 199–203.
11. Yasui T, Akatsuka M, Tobetto K, Hayaishi M, Ando T: The effect of hyaluronan on interleukin-1 α -induced prostaglandin E₂ production in human osteoarthritic synovial cells. *Agents Actions* 1992; 37: 155–156.
12. Smith GN, Myers SL, Brandt KD, Mickler EA: Effect of intraarticular hyaluronan injection in experimental canine osteoarthritis. *Arthritis and Rheumatism* 1998; 41: 976–985.
13. Engstrom-Laurent A, Hallgren R: Circulating hyaluronic acid levels vary with physical activity in healthy subjects and in rheumatoid arthritis patients: Relationship to synovitis mass and morning stiffness. *Arthritis and Rheumatism* 1987; 30: 1333–1338.
14. Balazs EA: The physical properties of synovial fluid and the special role of hyaluronic acid. "Disorders of the Knee" (Helfet AJ), 1982; pp 61–74 J.B. Lippincott.
15. Rizkalla G, Reiner A, Bogoch E, Poole AR: Studies of the articular cartilage proteoglycan aggrecan in health and osteoarthritis. *J Clin Invest* 1992; 90: 2268–2277.
16. Plaas AHK, West LA, Wong-Palms S, Nelson FRT: Glycosaminoglycan sulfation in human osteoarthritis 1998; 273: 12642–1269.
17. Bayliss MT, Osborne D, Woodhouse S, Davidson C: Sulfation of chondroitin sulfate in human articular cartilage. *J Biol Chem* 1999; 274: 892–900.
18. Shinmei M, Miyauchi S, Machida A, Miyazaki K: Quantitation of chondroitin 4-sulfate and chondroitin 6-sulfate in pathologic joint fluid. *Arthritis Rheum* 1992; 35: 1304–1308.
19. Asari A, Miyauchi S, Sekiguchi T, Machida A, Kuriyama S, Miyazaki K: Hyaluronan, cartilage destruction and hydrarthrosis in traumatic arthritis. *Osteoarthritis and cartilage* 1994; 2: 79–89.
20. Namiki O: Composition and metabolism of normal synovial fluid. *Joint Surgery* 1996; 15: 31–39. (in Japanese).
21. Sato M, Kasai T, Takemura M, Noma A, Takeuti S: Analysis of synovial fluid in patients with osteoarthritis and rheumatoid arthritis. *Cental Jpn J Orthop Surg Traumatol* 1995; 38: 199–220. (in Japanese).
22. Ueyoshi A, Kiura Y: Quantitation of synovial fluid in the rheumatoid arthritis and osteoarthritis: analysis of chondroitin sulfate content. *Joint Surgery* 1997; 24: 124–127. (in Japanese).
23. Lewis S, Crossman M, Flannelly J, Doherty M, Bayliss M, Mason R: Measurement of chondroitin sulphate disaccharides in pathological joint fluids. *Acta Orthop Scand* 1995; 66: 157–158.
24. Miyamoto K, Kozuka K, Kikuike A, Itozu M, Matsunaga T: Analysis of the synovial fluid in patients with osteoarthritis of knee joint in the point of OA-grade and daily activity patterns. *Cental Jpn J Orthop Surg Traumatol* 1995; 38: 121–122. (in Japanese).
25. Uesaka S, Nakayama Y, Fujii N, Kitta M, Yoshihara K, Shirai Y, Kaji K: Chondrosulfate level in synovial fluid on osteoarthritis of the knee. *Joint Surgery* 1998; 25: 5–8. (in Japanese).
26. Shinmei M, Kobayashi T, Yoshihara Y, Samura A: Significance of the levels of carboxy terminal typeII procollagen peptide, chondroitin sulfate isomers, tissue inhibitor of metalloproteinases, and metalloproteinases in osteoarthritis joint fluid. *J Rheumatol* 1995; 22: 78–81.

(Received, November 8, 1999)

(Accepted, December 8, 1999)