

Circulating Anti-Nuclear Antibodies in Systemic Sclerosis: Utility in Diagnosis and Disease Subsetting

Masataka Kuwana

Department of Allergy and Rheumatology, Nippon Medical School, Tokyo, Japan

The presence of circulating anti-nuclear antibodies (ANAs) is a hallmark of immune dysregulation in patients with systemic sclerosis (SSc). Currently, a variety of SSc-specific ANAs, including anticen-tromere, anti-topoisomerase I, and anti-RNA polymerase III antibodies, have been well characterized, and their commercial kits are available worldwide. Since these autoantibodies are specifically detected in SSc patients and are associated with unique sets of disease manifestations, they are widely used in routine clinical practice for diagnosis, clinical subgrouping, and prediction of future organ involvements and prognosis. In addition, SSc-specific ANAs are also useful in predicting future development of SSc in patients with Raynaud's phenomenon without any scleroderma skin changes, because their produc-tion often precedes onset of SSc symptoms. Application of circulating SSc-specific ANA measurement to clinical practice has greatly improved patient care, but utility of the autoantibody testing could be maxi-mized by combining other clinical information, such as degree and extent of skin thickness and disease duration. (J Nippon Med Sch 2017; 84: 56–63)

Key words: anti-nuclear antibody, biomarker, diagnosis, scleroderma, systemic sclerosis

Introduction

Systemic sclerosis (SSc) or scleroderma is a connective tissue disease characterized by excessive fibrosis, mi-croangiopathy, and immune dysregulation, including autoimmunity and chronic low-grade inflammation¹. Multiple genes contribute to disease susceptibility, but environmental exposures are likely to play a major role in causing and progressing the disease. Since clinical presentation in patients with SSc is highly variable, dis-ease subgrouping as well as prediction of future organ involvement and prognosis are extremely important in clinical setting. One of the distinctive hallmarks of the immune dysregulation is the presence of circulating autoantibodies reactive with various cellular components. The majority of disease-associated autoantibodies in SSc patients are anti-nuclear antibodies (ANAs) that target proteins that play essential roles in transcription, splic-ing, and cell division. It has been shown that ANA speci-ficities specifically detected in SSc patients are associated with unique disease manifestations. Therefore, individual ANAs are attractive biomarkers in routine rheumatology

practice. This review features the spectrum of SSc-specific ANAs and their clinical utility.

SSc-specific and SSc-related ANAs

ANAs detected by indirect immunofluorescence (IIF) technique are a hallmark of SSc, and are found in >95% of the patients. The majority of nuclear autoantigens rec-ognized by SSc sera have been already identified to date. ANAs detected in SSc sera can be divided into two groups, including SSc-specific ANAs, which are specifi-cally detected in SSc patients, and SSc-associated ANAs, which are found in patients with SSc but also in those with non-SSc connective tissue diseases. Currently, at least 10 ANA specificities specific to SSc have been re-ported and well characterized. These include anticen-tromere, anti-topoisomerase I (topo I), anti-RNA polym-erase (RNAP) III, anti-U3 ribonucleoprotein (RNP), anti-Th/To, anti-U11/U12 RNP, anti-PM-Scl, anti-Ku, anti-RuvBL1/2, anti-U1 RNP antibodies. These SSc-specific ANAs are detectable in >80% of SSc patients. Two classic autoantibodies discovered in the late 1970's are anti-topo

Correspondence to Masataka Kuwana, MD, PhD, Department of Allergy and Rheumatology, Nippon Medical School, 1-1-5 Sendagi, Bunkyo-ku, Tokyo 113-8602, Japan
E-mail: kuwanam@nms.ac.jp
Journal Website (<http://www2.nms.ac.jp/jnms/>)

Table 1 Staining patterns on IIF and clinical correlation of SSc-specific ANAs

ANA specificity	Staining pattern on IIF	Disease subset	Organ manifestations
Anticentromere (ACA)	Discrete speckled	lcSSc	PAH, DU in late disease
Anti-topo I	Speckled (with or without nucleolar)	dcSSc	ILD, DU in early disease
Anti-RNA polymerase III	Speckled (with or without nucleolar)	dcSSc	Rapid progression of skin thickening, SRC, GAVE, Malignancy at diagnosis
Anti-U3 RNP	Nucleolar	dcSSc/lcSSc	ILD, PAH, SRC, Lower GI involvement in early disease, Myopathy
Anti-Th/To	Nucleolar	lcSSc	ILD, PAH
Anti-U11/U12 RNP	Speckled	dcSSc/lcSSc	ILD
Anti-PM-Scl	Nucleolar	lcSSc (myositis overlap)	Myositis, DM rash
Anti-Ku	Speckled	lcSSc (myositis overlap)	Myositis
Anti-RuvBL1/2	Speckled	dcSSc (myositis overlap)	Myositis
Anti-U1 RNP	Speckled	lcSSc (‘MCTD’)	Inflammatory arthritis, Myositis, PAH

DM: dermatomyositis, DU: digital ulcer, GAVE: gastric antral vascular ectasia, GI: gastrointestinal tract, ILD: interstitial lung disease, MCTD: mixed connective tissue disease, PAH: pulmonary arterial hypertension, SRC: scleroderma renal crisis

I antibody and anticentromere antibody (ACA). Another group of antibodies, including anti-U1 RNP, anti-Ku, anti-PM-Scl antibodies, were first identified using double immunodiffusion technique. The remaining ANA specificities were discovered using RNA or protein immunoprecipitation (IP) assay. SSc-specific ANAs target various nuclear components involved in essential cellular processes, such as cell division, splicing, and transcription. These autoantibodies are rarely seen in patients with other connective tissue diseases without SSc features and thus are important diagnostic markers. In addition, detection of SSc-specific ANAs is clinically useful in classifying SSc patients into subtypes that are almost exclusively associated with characteristic clinical phenotypes (Table 1). SSc-specific ANAs are usually present at diagnosis of SSc or even precede appearance of SSc-related clinical manifestations such as Raynaud’s phenomenon. In addition, these autoantibodies do not switch from one antibody to another, and typically remain detectable throughout the disease course, regardless of the treatment regimen. Patients rarely have two or more SSc-specific ANAs together, indicating mutual exclusiveness.

On the other hand, SSc-associated ANAs include anti-nucleolar organizing region 90 (NOR-90) and anti-B23 antibodies. Anti-NOR-90, also termed the human upstream binding factor of RNAP I, was first reported in patients with cancer, but later in patients with connective tissue diseases, including SSc, Sjögren’s syndrome, and

rheumatoid arthritis². B23 is a nucleolar phosphoprotein overexpressed in many cancer cells, and is also targeted by cancer sera. Anti-B23 antibody is detected in <11% of SSc patients, and is associated with pulmonary arterial hypertension (PAH), often coexisting with anti-U3 RNP or anti-U1 RNP antibodies³. Anti-SSA/Ro and anti-SSB/La can occur in SSc patients, but are usually associated with concomitant Sjögren’s syndrome. Recently, it has been shown that tripartite motif family of protein 21 (TRIM21) or Ro52, which is one of components of the SSA/Ro autoantigen, is targeted by sera from patients with various connective tissue diseases in the absence of anti-SSA/Ro antibody. A recent multicenter cohort study involving 963 patients with SSc has found that anti-TRIM21 antibodies are present in 20% of the patients and are associated with interstitial lung disease (ILD) and overlapping features with other connective tissue diseases⁴.

There are a small proportion of SSc patients who are negative for ANA by IIF. In a recent cohort study involving 3,249 patients with SSc from a multicenter registry in North America, only 6.4% were negative for ANA⁵. ANA-negative SSc patients constitute a distinct subset with a greater proportion of males, less frequent and less severe vasculopathy, and more frequent lower gastrointestinal (GI) involvement.

Clinical Correlations of Individual SSc-specific ANAs

1. Anticentromere Antibody (ACA)

Moroi and colleagues first described ACA, which was originally associated with the CREST (Calcinosis, Raynaud's phenomenon, Esophageal hypomotility, Sclerodactyly, and Telangiectasia) variant of SSc⁶. ACA recognizes the kinetochore lesion of individual metaphase chromosomes, and the autoimmune targets are now identified as centromere protein (CENP)-A, B, and C. The frequency of ACA in SSc patients has been reported to be 20–30% in many ethnic groups. ACA typically produces discrete speckled staining pattern on IIF, but use of chromosomal spreads as the substrate is necessary to confirm its presence. ACA is sometimes detected in patients with primary Sjögren's syndrome or primary biliary cirrhosis as well as in individuals without an apparent connective tissue disease, who are almost always elderly⁷. The natural history of ACA-positive SSc patients includes long-standing Raynaud's phenomenon followed by appearance of puffy fingers after a variable period of time, ranging from a couple of months to 10 or more years. Patients with ACA are almost always classified as having limited cutaneous SSc (lcSSc)⁸, a term which has now replaced CREST syndrome. Severe ILD, cardiomyopathy or scleroderma renal crisis (SRC) almost never occur in ACA-positive patients, but 10–20% develop PAH later in the course of the disease. In this regard, evidence-based DETECT algorithm for identification of SSc patients at high risk for developing PAH includes ACA as one of the useful characteristics⁹.

2. Anti-topoisomerase I (Topo I) Antibody

Anti-topo I antibody was first discovered by double immunodiffusion assay as an antibody reactive with a 70-kDa protein (Scl-70)¹⁰, which was later identified as an enzyme that catalyzes relaxation of supercoiled double-stranded DNA, termed topo I¹¹. Anti-topo I antibody is detected in 20–30% of SSc patients in many ethnic groups, and is highly specific to SSc. The coexistence of anti-topo I and other SSc-related ANAs is rare (~1%). Almost two-thirds of anti-topo I-positive patients have diffuse cutaneous SSc (dcSSc), but progression of skin thickening is slower than those with anti-RNAP III antibody¹². ILD occurs in >70% of patients with anti-topo I antibody during the disease course, and ~25% develop severe disease which requires oxygen supplementation. The patients with severe ILD die of this complication at an average of 10 years after onset of SSc. Since ILD is now the major cause of death in SSc patients, anti-topo I antibody is considered to be a marker for poor prognosis. Never-

theless, progression and prognosis of ILD in patients with anti-topo I antibody are variable, although anti-topo I antibody titer is not useful in predicting a risk for developing severe ILD. Patients with anti-topo I antibody rarely develop PAH, but pulmonary hypertension secondary to ILD is fairly common later in the course of the disease. Other important organ involvements associated with anti-topo I antibody include cardiomyopathy and peripheral vascular complications, such as digital ulcer (DU) and gangrene, particularly early in the disease course⁸.

3. Anti-RNA Polymerase III (RNAP III) Antibody

A series of autoantibodies to RNAPs were identified by protein-IP assay. SSc sera contain autoantibodies directed against the RNAP I, II, and III in different combinations¹³, which fall into 4 groups: those reactive with RNAP I, II and III, those with RNAP I and III, those with RNAP III alone, and those with RNAP II alone¹⁴. Antibodies to RNAP II alone, which specifically target an active form, is considered as an SSc-associated ANA, because they are detected in patients without SSc as well as in those with SSc in association with anti-topo I antibody¹⁵. Instead, antibodies reactive with RNAP III, which are recognized commonly by the remaining groups, are most often regarded as the SSc-specific ANA. Sera positive for anti-RNAP III antibody produce a speckled staining pattern on IIF, but also produce nucleolar staining if anti-RNAP I antibodies coexist. The frequency of anti-RNAP III antibody in SSc patients varies among ethnic groups: higher frequency in North American Caucasian and UK patients (20–25%) in comparison with French or Japanese patients (5%)^{16,17}. Age at onset is older and the proportion of males tends to be higher in patients with anti-RNAP III. Nearly all patients with this antibody have dcSSc with rapidly progressive skin thickening. Despite that, many patients experience rapid regression of skin thickening over time even without treatment.

Patients with anti-RNAP III antibody have the highest risk for developing SRC, but they seldom develop severe ILD¹³. In Northern Europe and North America, approximately 60% of all patients experiencing SRC carry anti-RNAP III antibody¹⁸. In anti-RNAP III-positive SSc patients, coexistence of anti-RNAP II antibody and a higher antibody titer measured by enzyme immunoassays are independently associated with the SRC risk¹⁹. Lower incidence of SRC in Japan, French, and Mediterranean countries is thought to reflect the lower prevalence of anti-RNAP III antibody within these populations when compared with the UK and North America²⁰. Despite the

lower prevalence of anti-RNAP III antibody in French SSc patients, this antibody remains the strongest serological marker for SRC²⁰. Anti-RNAP III antibody has been detected in patients complicating acute renal failure mimicking SRC in the absence of sclerotic skin disease, but these patients eventually develop rapidly progressive skin thickening²¹. The presence of anti-RNAP III antibody should alert the managing physician to the increased risk of SRC and appropriate monitoring of blood pressure for early detection of this complication since the prompt introduction of angiotensin converting enzyme inhibitor can be life-saving. In recent cohorts, survival in patients with anti-RNAP III is better than those with anti-topo I, since SRC is more easily treated with angiotensin converting enzyme inhibitors than ILD. Finally, a case-control study using the large database identified anti-RNAP III antibody as one of risk factors for gastric antral vascular ectasia (GAVE; "watermelon stomach")²².

It has been recently highlighted that there is an association with cancer among SSc patients with anti-RNAP III antibody in close temporal relationship to SSc onset^{23,24}. The prevalence of cancer was almost similar among SSc patients with anti-RNAP III, ACA, and anti-topo I, but the median disease duration at cancer diagnosis was quite different: 13.4, 11.1, and 1.2 years in anti-topo I-positive, ACA-positive, and anti-RNAP III-positive patients, respectively²³. Another large-scale cohort including 1,044 patients with SSc found anti-RNAP III is an independent marker of coincident cancer at SSc diagnosis²⁵. The temporal relationship between cancer and SSc among anti-RNAP III-positive patients suggests that SSc is a para-neoplastic disorder in this patient subset²⁶.

4. Anti-U3 RNP Antibody

Anti-U3 RNP antibody was first identified by RNA-IP assay as an autoantibody that precipitated the U3 RNP particle, which consists of 34-kD fibrillarin complexed with U3 RNA²⁷. This antibody produces bright nucleolar staining on IIF. Anti-U3 RNP antibody is found in 4–10% of patients with SSc, and is the most frequent in African Americans²⁸. Two thirds of the patients have dcSSc, but others are classified as typical lcSSc. Nearly one-third of the patients develop non-inflammatory skeletal myopathy, which is an SSc-related condition distinct from myositis overlap. It has been reported that severe GI involvement, including pseudo-obstruction and small bowel malabsorption, is associated with anti-U3 RNP antibody, especially in early course of the disease (<2 years)²⁹. Severe internal organ involvement, including ILD, PAH, cardiomyopathy, and SRC are common in pa-

tients with anti-U3 RNP antibody, irrespective of dcSSc or lcSSc. Anti-U3 RNP antibody was identified as an independent risk factor for development of PAH in the large case-control study³⁰. In fact, PAH is the most common cause of death, leading to an increased mortality in this group of patients²⁸. An unusual combination of SRC followed later by PAH is occasionally found in patients with dcSSc and anti-U3 RNP antibody³¹. Survival in anti-U3 RNP-positive patients is comparable to that in patients with anti-topo I antibody.

5. Anti-Th/To Antibody

Anti-Th/To antibody was identified by RNA-IP assay as an antibody that precipitated both 7-2 and 8-2 RNAs. Later, it has been shown that this antibody directs against subunits of mitochondrial RNA processing and ribonuclease P complexes³². Of at least 6 subunits consisting of these complexes, a 120-kDa protein contains the major autoantibody epitope³³. Anti-Th/To antibody almost exclusively occurs in patients with lcSSc, although its frequency overall in SSc patients is only 2–5%. Like ACA-positive patients, anti-Th/To-positive patients are predominantly Caucasians, but tend to have a shorter duration of Raynaud's phenomenon before onset of other symptoms such as puffy fingers. Severe ischemic complications are infrequent, but this antibody is associated with significant ILD or PAH, which often occurs early in the disease course³⁴. In addition, patients with anti-Th/To can also develop pulmonary hypertension due to ILD. This increased frequency and severity of pulmonary complications result in a decreased survival compared with lcSSc patients without this antibody.

6. Anti-U11/U12 RNP Antibody

This autoantibody was identified by RNA-IP assay as the antibody that precipitated U11 RNA and U12 RNA together³⁵. It is a rare antibody specificity found in 1–3% of patients with SSc. This antibody produces speckled nuclear staining on IIF, and sometimes is associated with low-titer anti-U1 RNP antibodies. Patients with this antibody are classified as having either dcSSc or lcSSc. A characteristic feature of patients with anti-U11/U12 RNP antibody is a high frequency of ILD (~80%), which is often severe and rapidly progressive, and associated with a 2.25-fold greater risk of death in comparison with anti-U11/U12 RNP-negative patients with ILD.

7. Anti-PM-Scl Antibody

Anti-PM-Scl antibody was first identified by double immunodiffusion assay. The PM-Scl complex is composed of several subunits, of which the 100-kD and 75-kD proteins carry the main autoantigenic determinants.

Anti-PM-Scl antibody produces a homogenous nucleolar pattern, and is rarely found in non-Caucasian patients³⁶. Anti-PM-Scl-positive patients often present with the subacute myositis, but also have typical Raynaud's phenomenon and scleroderma skin changes, usually lcSSc³⁷. They are most frequently diagnosed as having SSc-polymyositis (PM) overlap, but a significant proportion of the patients have rashes consistent with dermatomyositis (DM). This antibody is found in more than 25% of SSc patients with myositis overlap, but in only 2% of SSc patients overall. Serious internal organ involvement is rare, leading to a favorable prognosis. Myositis is usually mild and shows a good response to corticosteroids. In a recent large cross-sectional study of patients with idiopathic inflammatory myopathies, 9% had anti-PM-Scl antibody in the absence of any scleroderma skin changes³⁸.

8. Anti-Ku Antibody

Anti-Ku antibody was first identified by double immunodiffusion assay. The Ku autoantigen is now recognized as a heterodimer of 70-kD and 80-kD subunits. The prevalence of anti-Ku antibody in SSc patients is ~2%³⁹. Anti-Ku antibody is primarily detected in patients with SSc in overlap. The majority of patients have typical Raynaud's phenomenon and scleroderma skin changes, usually lcSSc. Concomitant inflammatory myopathy is common, but some have additional features of lupus. Anti-Ku antibody is occasionally detected in patients with systemic lupus erythematosus (SLE) without SSc features, but additional lupus-associated autoantibodies such as anti-DNA antibodies are always positive. Disease onset in anti-Ku-positive patients is usually younger than 40 years. Internal organ involvement is infrequent and usually mild if present, but arthritis is common. Anti-Ku antibody is associated with fewer vascular manifestations, such as DU or telangiectasia⁴⁰. Myositis is usually mild and shows a good response to corticosteroids, leading to favorable prognosis.

9. Anti-RuvBL1/2 Antibody

This newly identified antibody recognizes a double hexamer consisting of RuvBL1 and RuvBL2, which is located in the nucleoplasm, by protein-IP assay⁴¹. Anti-RuvBL1/2 antibody is a rare antibody specificity detected in 1–2% of patients with SSc. This antibody produces speckled nuclear staining with a high antibody titer on IIF. Patients with this antibody are mostly males and have a unique combination of clinical features, including dcSSc and myositis overlap. Internal organ involvement is mild in general, but some develop signifi-

cant myocardial involvement.

10. Anti-U1 RNP Antibody

Anti-U1 RNP antibody was first identified by double immunodiffusion assay, in association with anti-Sm antibody, a lupus-specific ANA. Anti-U1 RNP antibodies are directed against the 70 K, A and C proteins associated with U1 RNA, while anti-Sm antibodies are directed against the B/B' and D proteins that are core components of the U series small nuclear RNAs involved in pre-messenger RNA splicing (U1, U2, U4/U6, U5, and others)⁴². Anti-U1 RNP antibody, which produces a pure speckled pattern with a high antibody titer, is primarily detected in patients with SSc in overlap. This antibody is preferentially found in African Americans and Orientals. Anti-U1 RNP antibody was first described as a serologic marker for mixed connective tissue disease (MCTD)⁴³, but is also found in sera from patients with SSc, SLE, PM/DM, or primary Sjögren's syndrome. Disease onset is at a relatively young age with mid 30's. SSc patients with this antibody usually present with inflammatory symptoms, such as myositis and arthritis. Raynaud's phenomenon and puffy fingers occur early in the disease, but later these patients develop typical manifestations of SSc. Most of them have lcSSc, although approximately 20% develop dcSSc. Serious complications are relatively uncommon, but pulmonary complications, including PAH and ILD, are sometimes life-threatening. Prognosis is favorable in general, but PAH is the most common cause of death.

Screening and Detection of SSc-specific ANAs in Routine Clinical Practice

A conventional method for ANA detection is IIF on cultured HEp-2 cell slides. This technique is recommended as the autoantibody screening test because it is highly sensitive and provides additional information on the antibody titer and staining pattern. Speckled staining is often detected in patients with dcSSc and suggests the presence of anti-topo I in case of a high ANA titer ($\geq 1 : 320$) or anti-RNAP III in case of a low ANA titer ($< 1 : 160$). Anti-U1 RNP and anti-RuvBL1/2 antibodies, which are associated with SSc in overlap, also produce a high-titer speckled pattern. A nucleolar pattern is fairly specific to SSc, and three major SSc-related anti-nucleolar antibodies are anti-U3 RNP, anti-Th/To, and anti-PM-Scl antibodies. Anti-RNAP III antibody also produces a nucleolar pattern when anti-RNAP I antibody coexists, but a concomitant speckled staining is always present. A discrete speckled pattern is often detected in patients with lcSSc and suggests the presence of ACA.

Identification of individual SSc-specific ANAs requires additional techniques, such as double immunodiffusion assay and/or IP assay, but convenient immunoassays such as enzyme immunoassays (EIA) are widely used in routine clinical practice, because of simplicity, reproducibility, speed, and ability to handle many samples at the same time. The majority of commercially available immunoassays utilize recombinant autoantigens, which are expressed in the bacterial or eukaryotic system, while some assays still use native proteins purified from cellular extracts. It is necessary to use validated kits, since false-positive and false-negative results can occur. If the results obtained from EIA were inconsistent with clinical presentation, re-evaluation using original detection methods, such as IP assay, is highly recommended.

Clinical Utility of SSc-specific ANAs

1. SSc Diagnosis

SSc-specific ANAs are useful in diagnosis of SSc because of their highly specific nature. In fact, the presence of ACA, anti-topo I antibody, or anti-RNAP III antibody is included in one of the classification criteria for SSc, which has been proposed by the American College of Rheumatology (ACR) and the European League against Rheumatism (EULAR)⁴⁴. In addition, the presence of SSc-specific ANA and nailfold capillary abnormalities are identified as independent predictors for future development of SSc in patients with Raynaud's phenomenon, but without any features suggestive of connective tissue diseases⁴⁵, indicating that SSc-specific ANAs are also useful in identifying patients with pre-SSc efficiently.

2. Disease Subsetting

Classification of patients into dcSSc and lcSSc subsets is useful in predicting future organ involvement and prognosis, but more precise subgrouping is feasible by considering SSc-specific ANAs because of their strong correlations with main organ manifestations (**Table 1**). For example, three major dcSSc-specific ANAs, including anti-topo I, anti-RNAP III, and anti-U3 RNP antibodies, cover >80% of dcSSc patients, but major organ involvements linking to poor prognosis are apparently different. The presence of anti-topo I antibody in dcSSc patients provides independent prognostic information with regard to a risk for progression of ILD and a greater risk of mortality⁸. In contrast, anti-RNAP III antibody has also been associated with increased mortality, primarily due to an increased prevalence of SRC⁸, but with improvement in the management of SRC, such associations may not be replicated in the future. On the other hand, PAH

still remains intractable condition with poor long-term survival even after introduction of a series of pulmonary vasodilators, and can occur in dcSSc patients with anti-U3 RNP antibody.

In addition, it is often difficult to classify patients into dcSSc or lcSSc early in the course of the disease when skin thickness is restricted to the distal portion of extremities. In this case, information on SSc-specific ANAs is useful in predicting the future extent of skin involvement. If one of dcSSc-associated ANAs, including anti-topo I, anti-RNAP III, and anti-U3 RNP antibodies, is present in circulation, disease modifying treatment should be indicated for preventing future organ damage at an early stage of dcSSc.

Conclusions

In summary, circulating ANA specificities are the best biomarkers for diagnosis and clinical subgrouping of SSc patients. Characterization of autoantibody status is an essential tool in clinical practice in the evaluation of patients who have SSc or are suspected to have this disease. In addition, because of their strong clinical associations, they allow clinicians to predict future disease manifestations and help to decide treatment strategy in individual patients. It is important to note, however, that clinical associations of autoantibodies are not absolute, and organ-specific manifestations can occur in the presence of any autoantibody reactivity.

Conflict of Interest: Kuwana M holds a patent on anti-RNAP III antibody-measuring kit.

References

1. Varga J, Abraham D: Systemic sclerosis: a prototypic multisystem fibrotic disorder. *J Clin Invest* 2007; 117: 557–567.
2. Rodriguez-Sanchez JL, Gelpi C, Juarez C, Hardin JA: A new autoantibody in scleroderma that recognizes a 90-kDa component of the nucleolus-organizing region of chromatin. *J Immunol* 1987; 139: 2579–2584.
3. Ulanet DB, Wigley FM, Gelber AC, Rosen A: Autoantibodies against B23, a nucleolar phosphoprotein, occur in scleroderma and are associated with pulmonary hypertension. *Arthritis Rheum* 2003; 49: 85–92.
4. Hudson M, Pope J, Mahler M, Tatibouet S, Steele R, Baron M, Canadian Scleroderma Research Group (CSRG), Fritzler MJ: Clinical significance of antibodies to Ro52/TRIM21 in systemic sclerosis. *Arthritis Res Ther* 2012; 14: R50.
5. Salazar GA, Assassi S, Wigley F, Hummers L, Varga J, Hinchcliff M, Khanna D, Schiopu E, Phillips K, Furst DE, Steen V, Baron M, Hudson M, Taillefer SS, Pope J, Jones N, Docherty P, Khalidi NA, Robinson D, Simms RW, Silver RM, Frech TM, Fessler BJ, Molitor JA, Fritzler MJ, Se-

- gal BM, Al-Kassab F, Perry M, Yang J, Zamanian S, Rev-eille JD, Arnett FC, Pedroza C, Mayes MD: Antinuclear antibody-negative systemic sclerosis. *Semin Arthritis Rheum* 2015; 44: 680–686.
6. Moroi Y, Peebles C, Fritzler MJ, Steigerwald J, Tan EM: Autoantibody to centromere (kinetochore) in scleroderma sera. *Proc Natl Acad Sci USA* 1980; 77: 1627–1631.
7. Gelber AC, Pillemer SR, Baum BJ, Wigley FM, Hummers LK, Morris S, Rosen A, Casciola-Rosen L: Distinct recognition of antibodies to centromere proteins in primary Sjogren's syndrome compared with limited scleroderma. *Ann Rheum Dis* 2006; 65: 1028–1032.
8. Kuwana M, Kaburaki J, Okano Y, Tojo T, Homma M: Clinical and prognostic associations based on serum antinuclear antibodies in Japanese patients with systemic sclerosis. *Arthritis Rheum* 1994; 37: 75–83.
9. Coghlan JG, Denton CP, Grünig E, Bonderman D, Distler O, Khanna D, Müller-Ladner U, Pope JE, Vonk MC, Doelberg M, Chadha-Boreham H, Heinzl H, Rosenberg DM, McLaughlin VV, Seibold JR, DETECT study group: Evidence-based detection of pulmonary arterial hypertension in systemic sclerosis: the DETECT study. *Ann Rheum Dis* 2014; 73: 1340–1349.
10. Douvas AS, Achten M, Tan EM: Identification of a nuclear protein (Scl-70) as a unique target of human antinuclear antibodies in scleroderma. *J Biol Chem* 1979; 254: 10514–10522.
11. Shero JH, Bordwell B, Rothfield NF, Earnshaw WC: High titers of autoantibodies to topoisomerase I (Scl-70) in sera from scleroderma patients. *Science* 1986; 231: 737–740.
12. Domsic RT, Rodriguez-Reyna T, Lucas M, Fertig N, Medsger TA Jr: Skin thickness progression rate: a predictor of mortality and early internal organ involvement in diffuse scleroderma. *Ann Rheum Dis* 2011; 70: 104–109.
13. Kuwana M, Kaburaki J, Mimori T, Tojo T, Homma M: Autoantibody reactive with three classes of RNA polymerases in sera from patients with systemic sclerosis. *J Clin Invest* 1993; 91: 1399–1404.
14. Kuwana M, Okano Y, Kaburaki J, Medsger TA Jr, Wright TM: Autoantibodies to RNA polymerases recognize multiple subunits and demonstrate cross-reactivity with RNA polymerase complexes. *Arthritis Rheum* 1999; 42: 275–284.
15. Satoh M, Kuwana M, Ogasawara T, Ajmani AK, Langdon JJ, Kimpel D, Wang J, Reeves WH: Association of autoantibodies to topoisomerase I and the phosphorylated (IIO) form of RNA polymerase II in Japanese scleroderma patients. *J Immunol* 1994; 153: 5838–5848.
16. Kuwana M, Okano Y, Kaburaki J, Tojo T, Medsger TA Jr: Racial differences in the distribution of systemic sclerosis-related serum antinuclear antibodies. *Arthritis Rheum* 1994; 37: 902–906.
17. Meyer OC, Fertig N, Lucas M, Somogyi N, Medsger TA Jr: Disease subsets, antinuclear antibody profile, and clinical features in 127 French and 247 US adult patients with systemic sclerosis. *J Rheumatol* 2007; 34: 104–109.
18. Penn H, Howie AJ, Kingdon EJ, Bunn CC, Stratton RJ, Black CM, Burns A, Denton CP: Scleroderma renal crisis: patient characteristics and long-term outcomes. *QJM* 2007; 100: 485–494.
19. Hamaguchi Y, Koder M, Matsushita T, Hasegawa M, Inaba Y, Usuda T, Kuwana M, Takehara K, Fujimoto M: Clinical and immunological predictors of scleroderma renal crisis for Japanese systemic sclerosis patients with anti-RNA polymerase III autoantibodies. *Arthritis Rheumatol* 2015; 67: 1045–1052.
20. Sobanski V, Dauchet L, Lefèvre G, Lambert M, Morell-Dubois S, Sy T, Hachulla E, Hatron PY, Launay D, Dubucquoi S: Prevalence of anti-RNA polymerase III antibodies in systemic sclerosis: new data from a French cohort and a systematic review and meta-analysis. *Arthritis Rheumatol* 2014; 66: 407–417.
21. Bhavsar SV, Carmona R: Anti-RNA polymerase III antibodies in the diagnosis of scleroderma renal crisis in the absence of skin disease. *J Clin Rheumatol* 2014; 20: 379–382.
22. Ghrénassia E, Avouac J, Khanna D, Derk CT, Distler O, Suliman YA, Airo P, Carreira PE, Foti R, Granel B, Berezne A, Cabane J, Ingegnoli F, Rosato E, Caramaschi P, Hesselstrand R, Walker UA, Alegre-Sancho JJ, Zarrouk V, Agard C, Riccieri V, Schiopu E, Gladue H, Steen VD, Allanore Y: Prevalence, correlates and outcomes of gastric antral vascular ectasia in systemic sclerosis: a EUSTAR case-control study. *J Rheumatol* 2014; 41: 99–105.
23. Shah AA, Rosen A, Hummers L, Wigley F, Casciola-Rosen L: Close temporal relationship between onset of cancer and scleroderma in patients with RNA polymerase I/III antibodies. *Arthritis Rheum* 2010; 62: 2787–2795.
24. Moinzadeh P, Fonseca C, Hellmich M, Shah AA, Chighizola C, Denton CP, Ong VH: Association of anti-RNA polymerase III autoantibodies and cancer in scleroderma. *Arthritis Res Ther* 2014; 16: R53.
25. Shah AA, Hummers LK, Casciola-Rosen L, Visvanathan K, Rosen A, Wigley FM: Examination of autoantibody status and clinical features that associate with cancer risk and cancer-associated scleroderma. *Arthritis Rheumatol* 2015; 67: 1053–1061.
26. Joseph CG, Darrah E, Shah AA, Skora AD, Casciola-Rosen LA, Wigley FM, Boin F, Fava A, Thoburn C, Kinde I, Jiao Y, Papadopoulos N, Kinzler KW, Vogelstein B, Rosen A: Association of the autoimmune disease scleroderma with an immunologic response to cancer. *Science* 2014; 343: 152–157.
27. Lischwe MA, Ochs RL, Reddy R, Cook RG, Yeoman LC, Tan EM, Reichlin M, Busch H: Purification and partial characterization of a nucleolar scleroderma antigen (Mr=34,000; pI, 8.5) rich in NG,NG-dimethylarginine. *J Biol Chem* 1985; 260: 14304–14310.
28. Aggarwal R, Lucas M, Fertig N, Oddis CV, Medsger TA Jr: Anti-U3 RNP autoantibodies in systemic sclerosis. *Arthritis Rheum* 2009; 60: 1112–1118.
29. Nishimagi E, Tochimoto A, Kawaguchi Y, Satoh T, Kuwana M, Takagi K, Ichida H, Kanno T, Soejima M, Baba S, Kamatani N, Hara M: Characteristics of patients with early systemic sclerosis and severe gastrointestinal tract involvement. *J Rheumatol* 2007; 34: 2050–2055.
30. Steen V, Medsger TA Jr: Predictors of isolated pulmonary hypertension in patients with systemic sclerosis and limited cutaneous involvement. *Arthritis Rheum* 2003; 48: 516–522.
31. Gündüz OH, Fertig N, Lucas M, Medsger TA Jr: Systemic sclerosis with renal crisis and pulmonary hypertension: a report of eleven cases. *Arthritis Rheum* 2001; 44: 1663–1666.
32. Van Eenennaam H, Vogelzangs JH, Lugtenberg D, Van Den Hoogen FH, Van Venrooij WJ, Pruijn GJ: Identity of the RNase MRP- and RNase P-associated Th/To autoantigen. *Arthritis Rheum* 2002; 46: 3266–3272.
33. Kuwana M, Kimura K, Hirakata M, Kawakami Y, Ikeda Y: Differences in autoantibody response to Th/To between systemic sclerosis and other autoimmune diseases. *Ann Rheum Dis* 2002; 61: 842–846.

34. Mitri GM, Lucas M, Fertig N, Steen VD, Medsger TA Jr: A comparison between anti-Th/To-and anticentromere antibody-positive systemic sclerosis patients with limited cutaneous involvement. *Arthritis Rheum* 2003; 48: 203–209.
35. Fertig N, Domsic RT, Rodriguez-Reyna T, Kuwana M, Lucas M, Medsger TA Jr, Feghali-Bostwick CA: Anti-U11/U12 RNP antibodies in systemic sclerosis: a new serologic marker associated with pulmonary fibrosis. *Arthritis Rheum* 2009; 61: 958–965.
36. Reimer G, Scheer U, Peters JM, Tan EM: Immunolocalization and partial characterization of a nucleolar autoantigen (PM-Scl) associated with polymyositis/scleroderma overlap syndromes. *J Immunol* 1986; 137: 3802–3808.
37. D'Aoust J, Hudson M, Tatibouet S, Wick J, Canadian Scleroderma Research Group, Mahler M, Baron M, Fritzler MJ: Clinical and serologic correlates of anti-PM/Scl antibodies in systemic sclerosis: a multicenter study of 763 patients. *Arthritis Rheumatol* 2014; 66: 1608–1615.
38. Brouwer R, Hengstman GJ, Vree Egberts W, Ehrfeld H, Bozic B, Ghirardello A, Grøndal G, Hietarinta M, Isenberg D, Kalden JR, Lundberg I, Moutsopoulos H, Roux-Lombard P, Vencovsky J, Wikman A, Seelig HP, van Engelen BG, van Venrooij WJ: Autoantibody profiles in the sera of European patients with myositis. *Ann Rheum Dis* 2001; 60: 116–123.
39. Mimori T, Akizuki M, Yamagata H, Inada S, Yoshida S, Homma M: Characterization of a high molecular weight acidic nuclear protein recognized by autoantibodies in sera from patients with polymyositis-scleroderma overlap. *J Clin Invest* 1981; 68: 611–620.
40. Rozman B, Cucnik S, Sodin-Semrl S, Czirkjak L, Varju C, Distler O, Huscher D, Aringer M, Steiner G, Matucci-Cerinic M, Guiducci S, Stamenkovic B, Stankovic A, Kveder T: Prevalence and clinical associations of anti-Ku antibodies in patients with systemic sclerosis: a European EUSTAR-initiated multi-centre case-control study. *Ann Rheum Dis* 2008; 67: 1282–1286.
41. Kaji K, Fertig N, Medsger TA Jr, Satoh T, Hoshino K, Hamaguchi Y, Hasegawa M, Lucas M, Schnure A, Ogawa F, Sato S, Takehara K, Fujimoto M, Kuwana M: Autoantibodies to RuvBL1 and RuvBL2: a novel systemic sclerosis-related antibody associated with diffuse cutaneous and skeletal muscle involvement. *Arthritis Care Res* 2014; 66: 575–584.
42. Pettersson I, Hinterberger M, Mimori T, Gottlieb E, Steitz JA: The structure of mammalian small nuclear ribonucleoproteins. Identification of multiple protein components reactive with anti-(U1)ribonucleoprotein and anti-Sm autoantibodies. *J Biol Chem* 1984; 259: 5907–5914.
43. Sharp GC, Irvin WS, Tan EM, Gould RG, Holman HR: Mixed connective tissue disease-an apparently distinct rheumatic disease syndrome associated with a specific antibody to an extractable nuclear antigen (ENA). *Am J Med* 1972; 52: 148–159.
44. van den Hoogen F, Khanna D, Fransen J, Johnson SR, Baron M, Tyndall A, Matucci-Cerinic M, Naden RP, Medsger TA Jr, Carreira PE, Riemekasten G, Clements PJ, Denton CP, Distler O, Allanore Y, Furst DE, Gabrielli A, Mayes MD, van Laar JM, Seibold JR, Czirkjak L, Steen VD, Inanc M, Kowal-Bielecka O, Müller-Ladner U, Valentini G, Veale DJ, Vonk MC, Walker UA, Chung L, Collier DH, Csuka ME, Fessler BJ, Guiducci S, Herrick A, Hsu VM, Jimenez S, Kahaleh B, Merkel PA, Sierakowski S, Silver RM, Simms RW, Varga J, Pope JE: 2013 classification criteria for systemic sclerosis: an American College of Rheumatology/European League against Rheumatism collaborative initiative. *Arthritis Rheum* 2013; 65: 2734–2747.
45. Koenig M, Joyal F, Fritzler MJ, Roussin A, Abrahamowicz M, Boire G, Goulet JR, Rich E, Grodzicky T, Raymond Y, Sénécal JL: Autoantibodies and microvascular damage are independent predictive factors for the progression of Raynaud's phenomenon to systemic sclerosis: a twenty-year prospective study of 586 patients, with validation of proposed criteria for early systemic sclerosis. *Arthritis Rheum* 2008; 58: 3902–3912.

(Received, December 28, 2016)

(Accepted, February 10, 2017)