* To whom correspondence should be addressed.

Table 1 Synergistic activity of a variety of imidazoles with permethrin (0.01 µg/fly) against susceptible houseflies.

No.	Compound (5 µg/fly)	Mortality (%) ^{a)}
1		60
2		90
3		0
4		100
5		100
6		0
7		0
8		0
9	KK-42	10
10	TH-27	0
11	KK-110	0
	Piperonyl butoxide	100

^{a)} Permethrin and each imidazole were not toxic alone at 0.01 µg and 5 µg, respectively.

susceptible houseflies. Permethrin alone was not toxic at 0.01 µg. When applied alone, all imidazole compounds examined were non-toxic at 5 µg. 1-Decyl- and 1-dodecylimidazoles (**1** and **2**), which are effective synergists against carbaryl,¹⁰⁾ exhibited some synergistic activity with permethrin. The phenoxypropyl analog **3**, which induces precocious metamorphosis in *B. mori* larvae as well as compounds **1** and **2**,¹³⁾ was inactive as a synergist. 1-(4-Octyloxyphenyl)- and 1-(4-benzyloxyphenyl)imidazoles (**4** and **5**) were as potent as piperonyl butoxide in the synergistic activity. It is noteworthy that compound **6** with one extra carbon atom between the imidazole and the benzene ring did not synergize the action of permethrin. 4-(4-Benzyloxyphenyl)imidazole (**7**), a structural isomer of **5**, was inactive as a synergist, suggesting that the location of a 4-benzyloxyphenyl group was important for the activity. 1,5-Diphenylimidazole (**8**), which has been reported to

Table 2 Synergistic activity of 1-substituted imidazoles with permethrin against susceptible houseflies.

No.	Compound	Synergistic ratio ^{a)}
2		1.9
4		4.1
5		3.6
12		2.7
13		2.4
14		2.4
15		2.8
	Piperonyl butoxide	4.0

^{a)} Ratio between LD₅₀ of permethrin alone to LD₅₀ of permethrin applied with synergist.

inhibit the epoxidation of aldrin in rat liver microsomes and armyworm guts preparations,¹¹⁾ did not show any synergistic activity with permethrin. 1,5-Disubstituted imidazoles such as **KK-42**, **TH-27** and **KK-110** were not effective synergists of permethrin, indicating some selectivity for the specific P450s.

Since the benzyloxyphenyl analog **5** showed considerably high synergistic activity, the analogs having a halogen atom at the benzyl group were synthesized and evaluated. Table 2 presents the synergistic ratios of the representative imidazoles. Compound **4** exhibited activity comparable to that of piperonyl butoxide which was twice as active as compound **2**. Compound **5** was slightly less active than piperonyl butoxide. The introduction of a fluoro substituent into the benzyl group irrespective of its position decreased the activity in comparison with that of compound **5**. There was little difference in activity between the 2-fluorobenzyl analog **12** and 2-chlorobenzyl analog **15**.

Based on the strong capacity of 1-substituted imidazoles to bind to the prosthetic heme iron of P450,¹⁰⁾ it would appear that 1-alkyloxyphenylimidazoles in this article inhibit P450s involved in the metabolism of permethrin. The activity of imidazoles as P450 inhibitors seems to be dependent on the

ability to bind to the hydrophobic regions close to the active site, the heme iron, of the enzyme. It might be possible to develop P450 inhibitors with reasonable selectivity by structural modification of the substituents on the imidazole ring.

ACKNOWLEDGMENTS

We thank Dr. M. Kamei of Earth Biochemical Co. for providing permethrin. This work was supported in part by Grant-in-Aid for Scientific Research 07556081 from the Ministry of Education, Science, Sports and Culture of Japan.

REFERENCES

- 1) E. Hodgson: *Insect Biochem.* **13**, 237 (1983)
- 2) G. C. Unnithan, J. F. Andersen, T. Hisano, E. Kuwano & R. Feyereisen: *Pestic. Sci.* **43**, 13 (1995)
- 3) O. Yamashita, K. Kadono-Okuda, E. Kuwano & M. Eto: *Agric. Biol. Chem.* **51**, 2295 (1987)
- 4) J. P. Roussel, M. Kiuchi, H. Akai & E. Kuwano: *C.R. Acad. Sci. Paris, Serie III* **305**, 141 (1987)
- 5) T. D. Jarvis, F. G. P. Earley & H. H. Rees: *Pestic. Biochem. Physiol.* **48**, 153 (1994)
- 6) E. Kuwano, M. Kikuchi & M. Eto: *Agric. Biol. Chem.* **52**, 1619 (1988)
- 7) B. Darvas, E. Kuwano, M. Eto, M. H. Tag El-Din & T. Timar: *Agric. Biol. Chem.* **54**, 3045 (1990)
- 8) B. Darvas, H. H. Rees, N. Hoggard, M. H. Tag El-Din, E. Kuwano, I. Belai & T. Timar: *Pestic. Sci.* **36**, 135 (1992)
- 9) C. F. Wilkinson, K. Hetnarski & L. J. Hicks: *Pestic.*

Biochem. Physiol. **4**, 299 (1974)

- 10) C. F. Wilkinson, K. Hetnarski, G. P. Cantwell & F. J. Di Carlo: *Biochem. Pharmac.* **23**, 2377 (1974)
- 11) T. D. Rogerson, C. F. Wilkinson & K. Hetnarski: *Biochem. Pharmac.* **26**, 1039 (1977)
- 12) E. Kuwano, T. Hisano, M. Eto, K. Suzuki, G. C. Unnithan & W. S. Bowers: *Pestic. Sci.* **34**, 263 (1992)
- 13) K. Shimazu, M. Shimizu, K. Suzuki & E. Kuwano: *J. Pesticide Sci.* **21**, 337 (1996)
- 14) R. M. Sacher, R. L. Metcalf & T. R. Fukuto: *J. Agric. Food Chem.* **16**, 779 (1968)

要 約

感受性イエバエに対するイミダゾール誘導体とペルメトリンとの共力活性

渡辺康一郎, 武谷立子, 山田直隆, 桑野栄一
昆虫成育制御活性を示すイミダゾール化合物とその関連化合物について, 感受性イエバエに対するペルメトリンとの殺虫共力効果を検討した。幼若ホルモン及び脱皮ホルモン生成に関与するチトクローム P450 を阻害する 1,5-二置換イミダゾール類には共力作用は認められなかった。共力作用を示したのは 1-置換イミダゾール類で, 特に, 1-(4-オクチルオキシフェニル)イミダゾール (4) と 1-(4-ベンジルオキシフェニル)イミダゾール (5) が高い活性を示した。化合物 4 の共力活性はピペロニルブトキシドと同等で, カルバリルに対して強い共力作用を示す 1-ドデシルイミダゾールの約 2 倍の強さであった。化合物 5 のベンジル基にフッ素あるいは塩素原子を導入すると活性は低下した。