Note

Population Dynamics of *Xanthomonas campestris* pv. *poae*, a Biological Control Agent for Annual Bluegrass, in Turfgrass Fields after Application

Tomoki NISHINO and Takane FUJIMORI

Plant Protection Research Laboratory, Japan Tobacco Inc., 6-2 Umegaoka, Aoba-ku, Yokohama, Kanagawa 227-8512, Japan

(Received September 22, 1997; Accepted December 17, 1997)

INTRODUCTION

Xanthomonas campestris pv. *poae*¹⁾ strain JT-P482²⁾ is a causal agent of bacterial wilt in annual bluegrass (*Poa annua* L.) and has been investigated as a biological control agent for annual bluegrass in intensively managed turfgrass fields. Strain JT-P482 was isolated from apparently asymptomatic annual bluegrass in Kanagawa prefecture in Japan in 1993.²⁾ Strain JT-P482 is able to destroy annual bluegrass when the inoculum concentration is over 1.0×10^6 colony forming unit/ml (cfu/ml).³⁾ High bacterial concentration of inocula,^{3,4)} repeated application,⁵⁾ high air temperature^{3,4)} and low air humidity³⁾ improve control of annual bluegrass. Seed production of annual bluegrass is also suppressed by strain JT-P482.⁵⁾ Strain JT-P482 does not cause any symptoms on 9 species of major crops or 8 species of other turfgrasses, however, rough bluegrass (*P. trivialis* L.) shows moderate wilt in greenhouse conditions.^{2,6)} *X. c.* pv. *poae*

JT-P482 and Rif-482,⁷⁾ which is a rifampicin tolerant mutant of strain JT-P482, do not show antagonistic activity to bacteria and fungi on solid media in a dual culture.⁷⁾ Populations of soil bacteria and soil fungi in mascarenegrass (*Zoysia tenuifolia* Willd. ex Trin.) turfgrass fields did not change drastically after application of strain Rif-482.⁷⁾ In the soil in a natural ecosystem, populations of many phytopathogenic bacteria are unable to survive for a long period.⁸⁾ The authors previously reported that the population of *X. c.* pv. *poae* strain Rif-482 in soil rapidly declined after bacterial suspensions of the strain had been applied to turfgrass fields.⁷⁾ This decline of the population in soil has been considered to be due to the fact that many phytopathogenic bacteria are overcome by indigenous microorganisms in soil in a natural ecosystem.⁷⁻⁹⁾ In this report, we examine the population dynamics of strain JT-P482 within turfgrasses after inoculation, and the persistence of strain Rif-482 within plant material and soil sections after application of the bacteria to creeping bentgrass (*Agrostis stolonifera* L.) turfgrass fields.

MATERIALS AND METHODS

1. Greenhouse Tests

To prepare inocula, a culture of *X. c.* pv. *poae* JT-P482 in YNB²⁾ liquid medium was placed on a rotary shaker at 28°C for 2 days and subsequently centrifuged. The collected cells were re-suspended with sterile distilled water and the bacterial concentration was adjusted to 1.0×10^8 cfu/ml by comparison with an absorption curve at 575 nm. Seeds of test plants were sown in organic soil in plastic pots and grown at 25°C/20°C (day/night) in a greenhouse for 3-4 weeks (Table 1). Plants were inoculated with the prepared inoculum by means of leaf clipping-inoculation.²⁾ The inoculated plants were

Table 1 Test plants and their susceptibility to *Xanthomonas campestris* pv. *poae* JT-P482.

Test plants	Cultivar/Origin	Susceptibility ^{b)}
Annual bluegrass (<i>Poa annua</i> L.)	Originating in UK ^{a)}	++
Rough bluegrass (<i>P. trivialis</i> L.)	Originating in UK	+
Kentucky bluegrass (<i>P. pratensis</i> L.)	Cultivar KenBlue	—
do	Cultivar Merit	—
Creeping bentgrass (<i>Agrostis stolonifera</i> L.)	Cultivar Pencross	—
do	Cultivar Seaside	—

^{a)} Plants originating in UK were purchased from Herbiseed (Wokingham, England, UK); the other plants were provided by the Mycogen Corporation (California, US).

^{b)} ++, plant was highly susceptible, inoculated plants died; +, plant was susceptible, inoculated plants wilted but a few plants regenerated; —, plant was resistant.

kept in the same greenhouse in order to observe the symptoms and the population dynamics of strain JT-P482 in weeds and turfgrasses. The method for counting the number of *X. c. pv. poae* JT-P482 in the tissue samples (5 mm length) of the inoculated plants was described in a previous report.⁴⁾

2. Field Tests

Two 2 m² plots of creeping bentgrass turf were used for investigating the persistence *X. c. pv. poae* Rif-482 in plant materials and in soil in the turfgrass field. Annual bluegrass seedlings grown in organic soil in plastic pots at 25°C/20°C (day/night) for 5–6 weeks in a greenhouse were transplanted at intervals of 20 cm into one plot. The other plot contained only creeping bentgrass. Both plots were mowed to a height of 1 cm every week. A bacterial suspension (1.0×10^8 cfu/ml) of *X. c. pv. poae* Rif-482⁷⁾, which is a rifampicin tolerant mutant of strain JT-P482, was sprayed at a rate of 400 ml/m² to both plots on April 15 in 1994. Spraying was carried out immediately after mowing.

To collect sod samples, the test plots were bored (40 mm in diameter) from the land surface to a depth of 5 cm. In the plot the annual bluegrass had been placed in, the sod samples were also taken from the turfgrass fields containing the weed. Each sod sample was separated into four parts: the topmost part of the land surface (A-part, consisting of plant leaves and culms); from the surface to a depth of 15 mm (B-part, consisting of plant roots, sand, and clay soil); from 15 mm to 30 mm below the surface (C-part, consisting of mostly clay soil); and from 30 mm to 45 mm below the surface (D-part, consisting of clay soil). Zero point five grams of the A-part and 5.0 g of the B-part, the C-part and the D-part, were each shaken separately on a rotary shaker with 30 ml of sterile distilled water to isolate the bacteria. To remove soil particles and plant debris, the suspensions were filtrated with No. 5A filter paper (Kiriya, Tokyo) by vacuum filtration. The filtrates were diluted serially and were spread onto YNA²⁾ plates containing 30 µg/ml of rifampicin and 200 µg/ml of cycloheximide. The plates were incubated at 28°C and the number of single colonies was counted 4 days after incubation.

RESULTS AND DISCUSSION

1. Growth and Migration of *X. c. pv. poae* JT-P482 in Grasses

Multiplication of *X. c. pv. poae* JT-P482 in inoculated leaf tips was observed in all test grasses, annual bluegrass, rough bluegrass, creeping bentgrass and Kentucky bluegrass (*P. pratensis* L.) (Fig. 1). In the lowest part of the culm of plants, multiplication was observed in annual bluegrass and rough bluegrass. However, the multiplication of strain JT-P482 in the lowest part of the culm was not observed in creeping bentgrass and Kentucky bluegrass. Wilt was observed in inoculated annual bluegrass and rough bluegrass but inoculated creeping bentgrass and Kentucky bluegrass showed no symptoms from *X. c. pv. poae* JT-P482 (Table 1). In our experiments, the bacterial multiplication in the lowest

Fig. 1 Multiplication of *Xanthomonas campestris* pv. *poae* JT-P482 in 5 mm lengths of tissue from the inoculation leaf tips (●) and the lowest part of the culm (○) of weeds and turfgrasses.

Creeping bentgrass (cv. Pencross (A), cv. Seaside (B)), Kentucky bluegrass (cv. Kenblue (C), cv. Merit (D)), annual bluegrass (E) and rough bluegrass (F) were inoculated with 1.0×10^8 cfu/ml of bacterial suspension of *X. c. pv. poae* JT-P482 by leaf clipping-inoculation. Inoculated plants were grown in the greenhouse which was kept at 25°C/20°C (day/night temperature). Bars indicate standard error ($n=3$).

part of the culm bears a relationship to the bacterial migration through the inoculated leaf. These results suggest that *X. c. pv. poae* JT-P482 is able to multiply in the cutting site of host grasses, annual bluegrass and rough bluegrass, and non-host grasses, creeping bentgrass and Kentucky bluegrass, however the spread through leaves of non-host grasses is limited. *X. c. pv. poae* JT-P482 is able to migrate throughout the leaves of host grasses only. This pathogenic trait was confirmed in the results of Robinson and Callow¹⁰⁾ who used *X. c. pv. poae* and *X. oryzae* pv. *oryzae*/rough bluegrass and rice (*Oryza sativa* L. cv. IR8) (pathogen/plant combinations).

In the case of the combinations of strain JT-P482/annual bluegrass or rough bluegrass (pathogen/plant combinations), strain JT-P482 multiplied in inoculated leaf tips till 7 days after inoculation and then the population declined. This decline in population was because the inoculated leaf tips had wilted and dried out (data not shown).

2. Persistence of *X. c. pv. poae* Rif-482 in Turfgrass Fields

In the plots containing transplanted annual bluegrass, *X. c. pv. poae* Rif-482 multiplied in the A-part of the sod sample (consisting of plant materials) from under 1.0×10^8 cfu/g of wet tissue to over 1.0×10^9 cfu/g of wet tissue about 1 month after application, and then the population declined to the

Fig. 2 Population dynamics of *Xanthomonas campestris* pv. *poae* Rif-482 in sod samples.

Samples were separated into 4 parts: A-part (consisting of plant leaves and culms) the topmost part of the landsurface; B-part (plant roots, sand and clay soil) land surface to a depth of 15 mm; C-part (mostly clay soil) from 15 mm to 30 mm below the surface; D-part (clay soil) from 30 mm to 45 mm below the surface. Bacterial suspensions (1.0×10^8 cfu/ml) of *X. c. pv. poae* Rif-482 were sprayed at the rate of 400 ml/m² on two 2 m² plots of creeping bentgrass turf, one containing annual bluegrass (●) and one containing creeping bentgrass only (○) on April 15 in 1994. Broken lines, an arrow, and bars indicate the detection limits of this method, the time of annual bluegrass death and standard error ($n=3$), respectively.

detection limit (3.0×10^3 cfu/g of wet tissue) about 3 months after application (Fig. 2). It is concluded that the population dynamics of strain Rif-482 in the A-part of the sod sample from the plots containing the annual bluegrass, was under the control of the number of strain Rif-482 within infected annual bluegrass, since the population of strain Rif-482 declined linearly in the A-part sample from the creeping bentgrass plots after application. In the annual bluegrass plots, the weeds showed wilt from about 2 weeks after application and died about 1 month after application. These results suggest that the decline in number of strain Rif-482 in the A-part of the sod sample 1 month after application is due to the death and desiccation of the host plants, annual bluegrass. The relationship between the disease development of annual bluegrass and the population

dynamics of *X. c. pv. poae* JT-P482 in annual bluegrass was confirmed in the results from the greenhouse experiments.³⁾

In the soil of the sod samples (B-part, C-part and D-part) from both test plots, viable cells of strain Rif-482 were observed (Fig. 2). However, we previously reported that *X. c. pv. poae* JT-P482 and Rif-482 were not able to increase their population over 1.0×10^3 cfu/g dry soil in a natural ecosystem.⁷⁾ The source of strain Rif-482 existing in the soil of the sod samples was supposed to be residue of plant materials of the A-part introduced by rainfall and/or watering, since the population dynamics of strain Rif-482 in soil parts showed a similar curve to those in the A-part and the population of the strain in the soil declined with the depth (Fig. 2).

These results indicate that *X. c. pv. poae* JT-P482 persists at the cutting sites of the non-host turfgrasses, e.g. creeping bentgrass and Kentucky bluegrass, longer than in the soil of the turfgrass fields. However, the population of *X. c. pv. poae* in the turfgrasses of the turfgrass fields declines linearly below under the detection limit (3.0×10^3 cfu/g wet tissue) about 2 months after application in the spring season. In the case of existing the host plant, annual bluegrass, after the death of any annual bluegrass plants in turfgrass fields, the population of *X. c. pv. poae* declines below under the detection limit without any adverse effects on desirable turfgrasses.

ACKNOWLEDGMENTS

The authors wish to thank Dr. M. Yamada and Ms. J. Nishino for their valuable advice to our research. We also thank Ms. R. Hokabe for her assistance.

REFERENCES

- 1) T. Egli & D. Schmidt: *Phytopathol. Z.* **104**, 138 (1982)
- 2) T. Nishino, S. Imaizumi, K. Miyabe, M. Yamada & M. Goto: *Ann. Phytopathol. Soc. Jpn.* **61**, 555 (1995)
- 3) T. Nishino & T. Fujimori: *Ann. Phytopathol. Soc. Jpn.* **62**, 492 (1996)
- 4) S. Imaizumi, T. Nishino, K. Miyabe, T. Fujimori & M. Yamada: *Biol. Control* **8**, 7 (1997)
- 5) S. Imaizumi, A. Tatenos & T. Fujimori: *J. Weed Sci. Tech.* **42**, 8 (1997)
- 6) S. Imaizumi, T. Nishino, M. Yamada & T. Fujimori: 15th Asian-Pacific Weed Science Society Conference (proceedings I(B)), pp. 693-697 (1995)
- 7) T. Nishino, K. Morita & T. Fujimori: *J. Pesticide Sci.* **22**, 326 (1997)
- 8) M. L. Schuster & D. P. Coyne: *Annu. Rev. Phytopathol.* **12**, 199 (1974)
- 9) M. Habte & M. Alexander: *Appl. Microbiol.* **29**, 159 (1975)
- 10) J. N. Robinson & J. A. Callow: *Plant Pathol.* **35**, 169 (1986)

要 約

芝地に散布されたスズメノカタビラの微生物防除剤である *Xanthomonas campestris* pv. *poae* の散布後の菌数変動

西野友規, 藤森 嶺

スズメノカタビラ (*Poa annua* L.) に対して除草活性を持つ植物病原細菌 *Xanthomonas campestris* pv. *poae* JT-P482 の散布後の運命を明らかにするために、宿主・非宿主植物体内における JT-P482 株の増殖・移行、そしてペントグラス芝生における Rif-482 株 (JT-P482 株のリファンピシン耐性変異株) の菌体数の変動を調査した。その結果、宿主植物であるスズメノカタビラとラフブルーグラス (*P. trivialis* L.) においては、剪葉接種された JT-P482 株は接種部位から地際部まで増殖しつつ移行す

るが、非宿主植物であるケンタッキーブルーグラス (*P. pratensis* L.)、ペントグラス (*Agrostis stolonifera* L.) においては剪葉接種された JT-P482 株は接種部位に局在し、植物体内を移行せず、病徴も現れないことが明らかとなった。また、4月中旬にペントグラス芝地に散布された Rif-482 株はスズメノカタビラが芝地中に存在しない場合、その芝地中の菌体数は散布直後から徐々に減少し、散布2カ月後には検出限界以下にまで減少した。しかし、その芝地にスズメノカタビラが存在した場合は、菌体数は散布直後から増加し、芝生中のスズメノカタビラが枯死する散布1カ月後を境として減少を始め、散布3カ月後には検出限界以下にまで減少した。以上の結果から、芝生上に散布後の *X. c.* pv. *poae* JT-P482 (Rif-482) は、スズメノカタビラを枯死させた後には速やかにその菌体数を減少させ、芝地中に高濃度に残留しないことが示された。