Full Paper

Dictyostelium Differentiation-Inducing Factor-1 Binds to Mitochondrial Malate Dehydrogenase and Inhibits Its Activity

Tomoko Matsuda¹, Fumi Takahashi-Yanaga^{1,*}, Tatsuya Yoshihara¹, Katsumi Maenaka², Yutaka Watanabe⁴, Yoshikazu Miwa¹, Sachio Morimoto¹, Yuzuru Kubohara⁵, Masato Hirata³, and Toshiyuki Sasaguri¹

¹Department of Clinical Pharmacology, Faculty of Medical Sciences, ²Medical Institute of Bioregulation,

³Laboratory of Molecular and Cellular Biochemistry, Faculty of Dental Sciences, Kyushu University, Fukuoka 812-8582, Japan

⁴Department of Applied Chemistry, Faculty of Engineering, Ehime University, Matsuyama 790-8577, Japan

⁵Biosignal Research Center, Institute for Molecular and Cellular Regulation, Gunma University, Maebashi 371-8512, Japan

Received December 11, 2009; Accepted January 8, 2010

Abstract. We have reported that the differentiation-inducing factors (DIFs) DIF-1 and DIF-3, morphogens secreted from *Dictyostelium discoideum*, inhibit proliferation of several cancer cells via suppression of the Wnt/ β -catenin signaling pathway. However, the target molecules of DIFs involved in the anti-proliferative effects are still unknown. In the present study, DIF-1–tethered resins were synthesized to explore the target molecules of DIFs, and mitochondrial malate dehydrogenase (mMDH) was identified as one of the target molecules. In the in vitro assay, DIF-1 and other analogs including 2-MIDIF-1, DIF-3, and 6-MIDIF-3 were found to be capable of binding to mMDH but not to cytoplasmic MDH. However, only DIF-1 and 2-MIDIF-1 inhibited the enzymatic activity of mMDH. The effects of DIF analogs on ATP content and cell proliferation were then analyzed using HeLa cells. DIF-1 and 2-MIDIF-1 were found to lower the ATP content and both chemicals inhibited HeLa cell proliferation, suggesting that inhibition of mMDH activity affected cell energy production, probably leading to the inhibition of proliferation. These results suggest that the inhibition of mMDH activity by DIF-1 and 2-MIDIF-1 could be one of the mechanisms to induce anti-proliferative effects, independent of the inhibition of the Wnt/ β -catenin signaling pathway.

Keywords: differentiation-inducing factor, mitochondrial malate dehydrogenase (mMDH), energy production, ATP content, cell proliferation

Introduction

Differentiation-inducing factors (DIFs), first identified in *Dictyostelium discoideum* as putative morphogens required for stalk cell differentiation (1 – 4), also affect mammalian cells (5 – 8). We reported that DIF-1 and DIF-3 [1-(3,5-dichloro-2,6-dihydroxy-4-methoxyphenyl)hexan-1-one and 1-(3-chloro-2,6-dihydroxy-4-methoxyphenyl)hexan-1-one, respectively] inhibited mammalian cell proliferation by suppressing the Wnt/ β -catenin sig-

naling pathway via the activation of glycogen synthase kinase-3 β (GSK-3 β) (9 – 13). Since the constant activation of the Wnt/ β -catenin signaling pathway has frequently been demonstrated in several types of human malignant neoplasms (14, 15), it could be suggested that DIFs are potent antitumor agents and identification of the target molecule(s) for DIFs may offer ideas for the design of new anticancer drugs. However, the target molecule(s) of DIFs involved in the anti-proliferative effect and/or the inhibition of the Wnt/ β -catenin signaling pathway are unknown. Shimizu et al. first reported that calmodulin-dependent cyclic nucleotide phosphodiesterase could be a pharmacological target molecule for DIF-1, whereas a specific inhibitor for phosphodiesterase 1 failed to mimic

*Corresponding author. yanaga@clipharm.med.kyushu-u.ac.jp
Published online in J-STAGE on February 20, 2010 (in advance)
doi: 10.1254/jphs.09348FP

the anti-proliferative effect of DIF-1 (16).

In the present study, the target molecules of DIFs were explored. For this purpose, DIF-1–tethered beads for affinity chromatography were synthesized. Upon further investigation, we found that the mitochondrial isoform of malate dehydrogenase (mMDH) was one of the target molecules of DIFs.

Malate dehydrogenase (MDH; EC 1.1.1.37) is an essential enzyme in the tricarboxylic acid cycle and malate-aspartate shuttle (17, 18), oxidizing malate to oxaloacetate and reducing oxaloacetate to malate, respectively. Two isoenzymes of MDH are found in animal tissues: one is in the mitochondria (mMDH, 314 amino acids) and the other is in the cytoplasm (cMDH, 332 amino acids) (19). The major role of mMDH is to oxidize malate to oxaloacetate in the tricarboxylic acid cycle, and that of cMDH is to reduce oxaloacetate to malate in the malate-aspartate shuttle. Since both the tricarboxylic acid cycle and malate-aspartate shuttle play essential roles in the energy production in cells, we analyzed the effect of DIF-1 and other DIF analogs on the activities of MDH in relation to ATP production and cell proliferation using HeLa cells.

Materials and Methods

Materials

DIF-1 and DIF-3 (1-(3,5-dichloro-2,6-dihydroxy-4-methoxyphenyl)hexan-1-one and 1-(3-chloro-2,6-dihydroxy-4-methoxyphenyl)hexan-1-one, respectively) were synthesized according to the procedure of Masento et al. (20). The other DIF analogs (2-MIDIF-1, 2-methoxy isomer of DIF-1, and 6-MIDIF-3, 6-methoxy isomer of DIF-3) were synthesized by Toyama Chemical Co. (Tokyo). Mitochondrial MDH (M2634) and cMDH (M7383) prepared from porcine heart were purchased from Sigma (St. Louis, MO, USA).

Preparation of DIF-1–tethered resin

To a solution of DIF-1 (20 mg, 0.065 mmol) and 2,6-lutidine (11 μ L, 0.078 mmol) in CH_2Cl_2 (200 μ L) was added 2-chlorethanesulfonyl chloride (8 μ L, 0.078 mmol), and the resulting mixture was stirred for 2 h at room temperature. After work-up, silica-gel chromatography (AcOEt/hexane 1:4) gave mono-*O*-ethenesulfonylated DIF-1 (11 mg, 42%) together with DIF-1 (58% recovered). AffiGel 10 (chemical capacity: 0.01 mmol/gel mL; Bio-Rad, Hercules, CA, USA,) was modified by treatment with 2-aminoethanethiol in isopropanol. The modified resin (1 mL) was treated with sulfonyl-DIF-1 (8.0 mg, 0.02 mmol) and triethylamine (3.3 μ L, 0.024 mmol) and ethanol (0.5 mL) for 1 h. The resin was obtained by filtration of the mixture and washing with ethanol. It was then similarly treated with *N*-phenylmaleimide

(3.5 mg, 0.02 mmol), triethylamine (3.3 μ L, 0.024 mmol), and ethanol (0.5 mL) for 3 h in order to cap the residual SH function. After filtration, the DIF-1–tethered affinity resin was obtained.

Cell culture

The human cervical carcinoma cell line HeLa was grown in Dulbecco's modified Eagle's medium (Sigma) supplemented with 10% fetal bovine serum, 100 U/mL penicillin G, and 0.1 mg/mL streptomycin.

Affinity chromatography using DIF-1–tethered resins

Cytoplasmic proteins were prepared from cells cultured in 100-mm dishes using NE-PER™ nuclear and cytoplasmic extraction reagents (Pierce, Rockford, IL, USA). Just before use, 10 μ L of DIF-1–tethered-beads were washed with 1 mL of buffer A containing 10 mM Tris-HCl pH 7.4, 1 mM EDTA, 50 mM KCl, and 10% glycerol and then resuspended with 200 μ L of the same buffer. The beads were incubated in the presence or absence of 100 μ M of DIF-1 at 4°C for 1 h, followed by the addition of the cell extract (1 mg of cytoplasmic proteins) and incubation for an additional 2 h. Proteins bound to the beads were extracted in Laemmli's sample buffer after extensive washing with buffer A.

Identification of binding protein

The extracted protein from the beads was separated by 10% SDS-PAGE, followed by silver staining. The gel pieces with bands only observed in the absence of DIF-1 were treated with trypsin, and the resulting peptides were analyzed by liquid chromatography/mass/mass spectrometry (LC-MS/MS) as described in the previous report (21). Mass spectroscopy data were analyzed by the MASCOT search engine (Matrix Science, Boston, MA, USA) using the International Protein Index (European Bioinformatics Institute, Hinxton, UK) as references.

Quartz crystal microbalance (QCM) assay

To measure the affinity of DIF and its analogs for MDH, a QCM assay was performed as described previously (22) with minor modifications. Briefly, 2 μ L of 4 mg/mL mMDH, cMDH or bovine serum albumin (BSA) was fixed on the sensor-tip of the QCM and placed in the chamber filled with 2 mL of phosphate-buffered saline. After the frequency of the sensor-tip was stable, ethanol (vehicle), DIF-1 or the analogs (2 μ L; final concentration of 30 μ M) were injected into the chamber. The frequency of the chip was measured at 25°C with constant stirring.

Measurement of enzymatic activities of mMDH and cMDH

Activities of mMDH or cMDH were assayed either in

the forward direction (malate + NAD⁺ → oxaloacetate + NADH) or reverse direction. Mitochondrial MDH (10.1 mg/mL) or cMDH (7.4 mg/mL) were diluted 1,000-times with a solution containing 0.1 mg/mL BSA and 1 μ M dithiothreitol and were pre-incubated with or without DIF-1 (final concentration: 30 μ M) for 10 min on ice. Then, 10 μ L of the enzyme solution and 40 μ L of β -NAD⁺ (final concentration: 10 mM) was mixed in a 96-well plate, followed by monitoring at the wavelength of 340 nm using a Flex Station3 micro-plate reader (Molecular Devices, Sunnyvale, CA, USA). Three minutes later, 50 μ L of malate (final concentration: 500 mM) was added to the each well as substrate. One unit was defined as the activity that can increase NADH by 1.0 μ mol per min.

Intracellular ATP measurement and cell proliferation assay

HeLa cells (2.0×10^4 cells/well) were seeded into 24-well plates and treated with or without 30 μ M DIF-1 or the analogs for given periods. Cells were harvested by the trypsin/EDTA treatment and enumerated using a Coulter Counter (Beckman Coulter, Fullerton, CA, USA). Intracellular ATP content was determined with a luciferin–luciferase method using an ATP assay kit (TOYO INK, Tokyo). Luciferase activity was determined with a luminometer (Lumat LB 9507; Berthold Technologies, Bad Wildbad, Germany).

Results

DIF-1 modification and immobilization

Ligand-immobilized beads were used as a tool to identify the target molecules of DIF-1. Therefore, we first synthesized the DIF-1 analog, DIF-1-sulfonate shown in Fig. 1A (see also Fig. 5A for the structure of DIF-1), which could be connected to reactive beads. Assays of cell proliferation was performed using HeLa cells, since we reported that DIF-1 and DIF-3 inhibited mammalian cell proliferation. As shown in Fig. 1B, DIF-1-sulfonate clearly exhibited inhibitory effects on cell proliferation, albeit with a weaker potency than DIF-1, encouraging us to further synthesize DIF-1–tethered beads using the analog.

DIF-1 binds to mMDH and the effects on the enzyme activity

HeLa cell extracts were mixed with immobilized-DIF-1 and affinity chromatography was performed as described in Materials and Methods. To identify the proteins that bind specifically to DIF-1, the cell extracts were incubated with the immobilized beads in the presence or absence of DIF-1 at 4°C for 2 h. Binding proteins were separated by 10% SDS-PAGE and two silver-

Fig. 1. Chemical structure and effects of DIF-1-sulfonate. A) Structure of DIF-1-sulfonate. The sulfonate group acting as a bridge for connection to the beads is marked by a rectangle. B) Effect of DIF-1-sulfonate on the cell proliferation. HeLa cells were treated with the indicated chemicals for 48 h and enumerated. The reported data are each the mean $\pm$ S.D. of three independent experiments ($n = 3$). ** $P < 0.01$ vs. control (by Student's t -test).

Fig. 2. DIF-1–tethered resin-binding proteins in HeLa cells. The extract from HeLa cells was loaded onto DIF-1–tethered resin with or without excess amount of DIF-1. The bound protein was analyzed by SDS-PAGE followed by silver staining. Arrows indicate two bands that specifically bound to DIF-1–tethered resin.

stained bands were identified as the proteins that bind to DIF-1–tethered beads, but not in the presence of the soluble ligand (Fig. 2). The upper band was identified as mMDH by mass spectrometry, while the lower band was

Fig. 3. DIF-1 binds with mMDH. QCM assay was performed as described in Materials and Methods. A difference of frequency was measured before and after injection. The arrows in the upper figure indicate the injection points of vehicle (ethanol: EtOH) or DIF-1. The lower graph shows the difference in frequency between before and after injection. The reported data are each the mean $\pm$ S.D. of three independent experiments. * $P < 0.05$ vs. control (by Student's *t*-test).

not identified. To confirm whether DIF-1 binds directly to mMDH in vitro, we performed QCM analysis. As shown in Fig. 3, the frequency was significantly reduced by adding DIF-1 when mMDH was fixed on the sensor tip. No significant changes were observed when cMDH, the cytosolic isoenzyme of MDH or BSA was fixed on the tip, indicating that DIF-1 directly binds to mMDH but not to cMDH. Subsequently, the effect of DIF-1 on the activities of mMDH and cMDH were analyzed. Mitochondrial MDH or cMDH, obtained from a commercial source was assayed for NAD⁺-dependent oxidation (forward direction). DIF-1 inhibited mMDH activity in a dose-dependent manner (Fig. 4A), whereas cMDH activity was not affected (Fig. 4B). The result agreed well with the result shown in Fig. 3, indicating that DIF-1 inhibited mMDH activity by direct binding.

Effects of DIF-1 analogs on mMDH

The interactions between mMDH and DIF analogs (2-MIDIF-1, DIF-3, and 6-MIDIF-3) were analyzed by the QCM assay. As shown in Fig. 5A, all DIF analogs examined were revealed to directly bind to mMDH and the positive signals with 2-MIDIF-1 was highest among the analogs examined, suggesting that 2-MIDIF-1 had the highest affinity. The effects of DIF analogs on the enzyme activity were then analyzed (Fig. 5B). Interestingly, of the 3 analogs examined, only 2-MIDIF-1 was inhibitory, although the effect of 2-MIDIF-1 appeared weaker than that of DIF-1 (Figs. 4A and 5B).

Fig. 4. Effects of DIF-1 on MDH activity. The mMDH (A) and cMDH (B) activities were assayed in the forward direction using different concentrations of DIF-1. Upper panels show the change of absorption and lower panels show the graph for MDH activity calculated from the upper panels as described in Materials and Methods. Each value is the mean $\pm$ S.E.M. of three independent experiments done in triplicate. ** $P < 0.01$ vs. control (by Student's *t*-test).

Fig. 5. Effect of DIF analogs on mMDH. A) Upper panel shows the chemical structure of DIF analogs and lower panel shows the binding property of DIF analogs to mMDH. Difference of frequency was measured before and after injection. Data are shown as the difference of frequency between before and after injection. Each value is the mean $\pm$ S.E.M. of three independent experiments. * $P < 0.05$ vs. control, ** $P < 0.01$ vs. control (by Student's *t*-test). B) Effect of DIF analogs on mMDH activity. mMDH activity was measured in the forward direction. Upper panels show the change of absorption and lower panel shows the graph for mMDH activity calculated from the upper panel as described in Materials and Methods. Each value is the mean $\pm$ S.E.M. of three independent experiments done in triplicate. * $P < 0.05$ vs. control (by Student's *t*-test).

Effects of DIF-1 and DIF analogs on ATP content and cell growth

To examine the relationship between the inhibition of mMDH activity, cell energy production, and cell proliferation, the effects of DIF analogs on intracellular ATP content and cell proliferation were examined using HeLa cells. The cellular content of ATP was gradually increased as the time of culture increased up to 72 h, and DIF-3 and 6-MIDIF-3, ineffective in the mMDH assay, did not impact the results (Fig. 6A). On the other hand, DIF-1 and 2-MIDIF-1 lowered the ATP content in the

same order of potency as that for the inhibition of mMDH activity.

Cell proliferation was also analyzed. As shown in Fig. 6B, DIF-1 and DIF-3 were the most potent inhibitors of the cell proliferation, while 6-MIDIF-3 was ineffective. In the case of 2-MIDIF-1, the inhibition of cell proliferation became evident at 72 h, when the cellular content of ATP was apparently lowered to a similar level as that with DIF-1. However, the ATP content of HeLa cells after 72-h treatment with 2-MIDIF-1 was still approximately 70% of that at time 0, suggesting that 2-MIDIF-

Fig. 6. Effects of DIF analogs on intracellular ATP content and cell growth. HeLa cells (2.0×10^4 cells/well) were seeded into 24-well plates and treated with or without 30 μ M of DIF analogs for the given periods. Cells were harvested by trypsin/EDTA treatment, followed by the measurement of the ATP content (A) and cell number (B). The reported data are each the mean $\pm$ S.E.M. of three independent experiments, each performed in duplicate. * $P < 0.05$ vs. control, ** $P < 0.01$ vs. control (by Student's *t*-test).

1-treated HeLa cells did not deplete ATP. This result was well correlated with the fact that 2-MIDIF-1 showed a weak inhibitory effect on cell proliferation.

Discussion

Previously, we have been reported that DIF-1 and DIF-3 inhibit cell proliferation via the suppression of the Wnt/ β -catenin signaling pathway in human cancer cells (9–13). Although the purpose of the study was to find the target molecule of DIFs related with the Wnt/ β -catenin signaling pathway, unfortunately we could not identify such a protein from mass spectrometry. Mitochondrial MDH, one of the DIFs target proteins found in this study, is not a component of the Wnt/ β -catenin signaling pathway and this enzyme was inhibited by only DIF-1 and not by DIF-3, whereas both chemicals showed an inhibitory effect on the Wnt/ β -catenin signaling pathway. These observations suggest that other target molecules related with the Wnt/ β -catenin signaling pathway must exist. Further studies are required to identify the target molecules of DIF analogs.

We analyzed the changes in cellular content of ATP and cell proliferation in HeLa cells treated with DIF analogs. DIF-1 lowered the ATP content and inhibited cell proliferation more robustly than 2-MIDIF-1. This might have resulted from the fact that DIF-1 exhibited stronger inhibition of mMDH activity than 2-MIDIF-1 and suggest that lowering the ATP content caused by the inhibition of mMDH is pharmacologically relevant to the inhibition of cell proliferation.

Alternatively, a recently reported finding that DIF-1, but not 2-MIDIF-1, was able to promote glucose uptake and consumption could explain the inhibition results (23). It is possible that DIF-1-treated cells deplete glucose in the medium during the culture period. Furthermore, it is well known that glycolysis rather than the

tricarboxylic acid cycle is the main energy production system in a number of cancer cell types (24–26), suggesting that inhibition of mMDH activity might not have a strong effect on ATP production and inhibition of glycolysis might be more effective to reduce ATP production in HeLa cells. Consequently, depletion of glucose in the culture medium by DIF-1 likely resulted in the decrease of the ATP content in the HeLa cells. This notion also might explain the delay of 2-MIDIF-1 on the ATP content. At the initial stage of the culture, 2-MIDIF-1-treated cells produced ATP mainly using glycolysis, uptaking glucose from the culture medium, but at the later stage, consumption of glucose triggered oxidation of fatty acids to yield acetyl-CoA for the tricarboxylic acid cycle, and gluconeogenesis from amino acids followed by glycolysis, thus producing an overall lower level of ATP.

In this report, we showed that mMDH, one of the essential enzymes in the tricarboxylic acid cycle and malate-aspartate shuttle for main energy production in normal cells, is one of the target molecules of DIFs. Indeed, all DIFs analogs examined in this study could bind to mMDH. However, among the DIF analogs examined, only DIF-1 and 2-MIDIF-1 inhibit mMDH activity, suggesting that the chlorine at the 5 position of the DIF ring is essential for inhibition. Although we have reported that both DIF-1 and DIF-3 inhibited proliferation of human cancer cells, this finding may suggest that DIF-3 is more suitable as a candidate for a new anti-cancer drug than DIF-1, since DIF-3 lacks the chlorine at the 5 position of the DIF ring and does not affect mMDH activity.

Acknowledgments

This work was supported by grants from the Ministry of Education, Culture, Sports, Science and Technology, Japan (a Grant-in-Aid

for Scientific Research). We thank M. Oda and E. Fujimoto in the Laboratory for Technical Support, Medical Institute of Bioregulation, Kyushu University for technical assistance.

References

- Morris HR, Taylor GW, Masento MS, Jermyn KA, Kay RR. Chemical structure of the morphogen differentiation inducing factor from *Dictyostelium discoideum*. *Nature*. 1987;328:811–814.
- Morris HR, Masento MS, Taylor GW, Jermyn KA, Kay RR. Structure elucidation of two differentiation inducing factors (DIF-2 and DIF-3) from the cellular slime mould *Dictyostelium discoideum*. *Biochem J*. 1988;249:903–906.
- Wurster B, Kay RR. New roles for DIF? Effects on early development in *Dictyostelium*. *Dev Biol*. 1990;140:189–195.
- Morandini P, Offer J, Traynor D, Nayler O, Neuhaus D, Taylor GW, et al. The proximal pathway of metabolism of the chlorinated signal molecule differentiation-inducing factor-1 (DIF-1) in the cellular slime mould *Dictyostelium*. *Biochem J*. 1995;306:735–743.
- Kubohara Y. DIF-1, Putative morphogen of *D. discoideum*, suppresses cell growth and promotes retinoic acid-induced cell differentiation in HL-60. *Biochem Biophys Res Commun*. 1997;236:418–422.
- Kubohara Y. Effect of differentiation-inducing factors of *Dictyostelium discoideum* on human leukemia K562 cells: DIF-3 is the most potent anti-leukemic agent. *Eur J Pharmacol*. 1999;381:57–62.
- Gokan N, Kikuchi H, Nakamura K, Oshima Y, Hosaka K, Kubohara Y. Structural requirements of *Dictyostelium* differentiation-inducing factors for their stalk-cell-inducing activity in *Dictyostelium* cells and anti-proliferative activity in K562 human leukemic cells. *Biochem Pharm*. 2005;70:676–685.
- Miwa Y, Sasaguri T, Kosaka C, Taba Y, Ishida A, Abumiya T, et al. Differentiation-inducing factor-1, a morphogen of dictyostelium, induces G(1) arrest and differentiation of vascular smooth muscle cells. *Circ Res*. 2000;86:68–75.
- Takahashi-Yanaga F, Taba Y, Miwa Y, Kubohara Y, Watanabe Y, Hirata M, et al. *Dictyostelium* differentiation-inducing factor-3 activates glycogen synthase kinase-3 β and degrades cyclin D1 in mammalian cells. *J Biol Chem*. 2003; 278:9663–9670.
- Yasmin T, Takahashi-Yanaga F, Mori J, Miwa Y, Hirata M, Watanabe Y, et al. Differentiation-inducing factor-1 suppresses gene expression of cyclin D1 in tumor cells. *Biochem Biophys Res Commun*. 2005;338:903–909.
- Mori J, Takahashi-Yanaga F, Miwa Y, Watanabe Y, Hirata M, Morimoto S, et al. Differentiation-inducing factor-1 induces cyclin D1 degradation through the phosphorylation of Thr²⁸⁶ in squamous cell carcinoma. *Exp Cell Res*. 2005;310:426–433.
- Takahashi-Yanaga F, Mori J, Matsuzaki E, Watanabe Y, Hirata M, Miwa Y, et al. Involvement of GSK-3 β and DYRK1B in differentiation-inducing factor-3-induced phosphorylation of cyclin D1 in HeLa cells. *J Biol Chem*. 2006;281:38489–38497.
- Matsuzaki E, Takahashi-Yanaga F, Miwa Y, Hirata M, Watanabe Y, Sato N, et al. Differentiation-inducing factor-1 alters canonical Wnt signaling and suppresses alkaline phosphatase expression in osteoblast-like cell lines. *J Bone Miner Res*. 2006;21:1307–1316.
- Barker N, Clevers H. Mining the Wnt pathway for cancer therapeutics. *Nat Rev Drug Discov*. 2006;5:997–1014.
- Fodde R, Brabletz T. Wnt/ β -catenin signaling in cancer stemness and malignant behavior. *Curr Opin Cell Biol*. 2007;19:150–158.
- Shimizu K, Murata T, Tagawa T, Takahashi K, Ishikawa R, Abe Y, et al. Calmodulin-dependent cyclic nucleotide phosphatase (PDE1) is a pharmacological target of differentiation-inducing factor-1, an antitumor agent isolated from *Dictyostelium*. *Cancer Res*. 2004;64:2568–2571.
- Goward CR, Nicholls DJ. Malate dehydrogenase: a model for structure, evolution, and catalysis. *Protein Sci*. 1994;3:1883–1888.
- LaNoue KF, Bryla J, Bassett DJ. Energy-driven aspartate efflux from heart and liver mitochondria. *J Biol Chem*. 1974;249:7514–7521.
- Joh T, Takeshima H, Tsuzuk T, Setoyama C, Shimada K, Tanase S, et al. Cloning and sequence analysis of cDNAs encoding mammalian cytosolic malate dehydrogenase. Comparison of the amino acid sequences of mammalian and bacterial malate dehydrogenase. *J Biol Chem*. 1987;262:15127–15131.
- Masento MS, Morris HR, Taylor G, Johnson SJ, Skapski AC, Kay RR. Differentiation-inducing factor from the slime mould *Dictyostelium discoideum* and its analogues. Synthesis, structure and biological activity. *Biochem J*. 1988;256:23–28.
- Matsumoto M, Hatakeyama S, Oyamada K, Oda Y, Nishimura T, Nakayama KI. Large-scale analysis of the human ubiquitin-related proteome. *Proteomics*. 2005;5:4145–4151.
- Lu Q-W, Morimoto S, Harada K, Du C-K, Takahashi-Yanaga F, Miwa Y, et al. Cardiac troponin T mutation R141W found in dilated cardiomyopathy stabilizes the troponin T-tropomyosin interaction and causes Ca²⁺ desensitization. *J Mol Cell Cardiol*. 2003;35:1421–1427.
- Omata W, Shibata H, Nagasawa M, Kojima I, Kikuchi H, Oshima Y, et al. *Dictyostelium* differentiation-inducing factor-1 induces glucose transporter 1 translocation and promotes glucose uptake in mammalian cells. *FEBS J*. 2007;274:3392–3404.
- Warburg O. On the origin of cancer cells. *Science*. 1956;123:309–314.
- Shaw RJ. Glucose metabolism and cancer. *Curr Opin Cell Biol*. 2006;18:598–608.
- Ganapathy V, Thangaraju M, Prasad PD. Nutrient transporters in cancer: relevance to Warburg hypothesis and beyond. *Pharmacol Ther*. 2009;121:29–40.