ETHYLALCOHOL INHIBITION OF BRAIN MITOCHONDRIAL RESPIRATION STIMULATED BY DINITROPHENOL

TADAATSU NUKADA* AND NORIO ANDOH**

*Department of Neuropharmacology and Neurochemistry, Institute of Higher Nervous Activity,
Faculty of Medicine, Osaka University, Kita-ku, Osaka*

Received for publication January 26, 1967

Beer and Quastel (1) showed that alcohol exercised marked inhibitions of potassium-stimulated rat brain cortex respiration at the concentration of 0.6 M which affected but little the unstimulated respiration of rat brain cortex slices incubated in glucose-phosphate-Ringer media. The alcohol, however, have little or no inhibitory effects on rat brain mitochondrial respiration incubated in the optimal media for oxidative phosphorylation. In the present paper it is shown that the alcohol has the inhibitory action on the brain mitochondrial respiration stimulated by 2, 4-dinitrophenol (DNP).

The mitochondrial fraction were prepared from rat whole brain as described previously (2). Oxygen consumption was measured by the conventional Warburg's method. DNP or ethylalcohol had no effects on the respiration of brain mitochondria when incubated in a complete medium containing 6.7 mM MgSO₄, 20 mM sodium phosphate buffer at pH 7.4, 2 mM disodium ATP, 33 mM KCl, 500 μg cytochrome c, 6.7 mM sodium pyruvate, 0.67 mM sodium malate and 0.3 mM NAD in a volume of 3 ml. On the other hand, when brain mitochondria were incubated in a simple medium which contained 50 mM KCl, 20 mM tris-buffer at pH 7.4, 6.7 mM MgSO₄, 6.7 mM sodium pyruvate and 0.67 mM sodium malate, the addition of DNP increased the respiration remarkably, although the control respiration in this medium was less than that in a complete medium. The respiration stimulated by the addition of DNP in a simple medium was inhibited significantly

TABLE 1. Effect of ethylalcohol on the respiration of rat brain mitochondria.

Incubation medium	Addition	μl O ₂ consumed per mitochondria equiv. to 500mg w.wt.			
		0-10 min	10-20 min	20-30 min	% control
Complete medium	None	86.4	76.5	72.6	100
	DNP(10 ⁻⁴ M)	96.7	67.3	62.3	96
	Ethanol (0.3 M)	89.2	69.1	67.7	96
	Ethanol (0.6 M)	81.8	68.4	67.0	93
Simple medium	None	37.0	35.7	34.9	100
	Ethanol (0.3 M)	37.0	35.6	34.4	100
	Ethanol (0.6 M)	38.7	38.7	37.5	108
	DNP(10 ⁻⁴ M)	72.0	67.9	65.2	100
	+ Ethanol (0.3 M)	59.6	59.6	49.7	82
	+ Ethanol (0.6 M)	44.6	41.9	35.4	59

額田 忠篤・安藤 憲夫

* Present address : 1st Division of Internal Medicine, Faculty of Medicine, Osaka University, Osaka.

** Present address : Department of Pharmacology, Faculty of Medicine, Osaka University, Osaka.

by the addition of ethylalcohol at the concentrations which affected but little the unstimulated brain mitochondrial respiration as shown in Table 1.

It is suggested that alcohols exercise their inhibitory effects on brain respiration not only at the brain cell membranes but also at the mitochondrial respiration.

REFERENCES

- 1) BEER, C.T. AND QUASTEL, J.H.: *Can. J. Biochem. Physiol.* **36**, 543 (1958); 2) NUKADA, T.: *Ibid.* **43**, 1119 (1965)

NEURONAL ADAPTATION TO BARBITURATES

YOSHIHIRO KINOSHITA, HIDEYO OHSHIKA AND KENGO NAKAI

Department of Pharmacology, Sapporo Medical College, Sapporo

Received for publication February 14, 1967

Although the mechanism involved in the development of tolerance to barbiturates is poorly understood, available evidence indicates that in experimental animals the tolerance often occurs as a result of induced synthesis of the hepatic microsomal enzymes concerned in the oxidation of barbiturates (1). However, it is well known that the sensitivity of various receptors in the central nervous system decreases after repeated administration of such drugs as narcotics, analgesics and other centrally acting drugs.

Attempts were, therefore, made to examine a predominantly causative factor in the development of tolerance to barbiturates. Female rats or mice were pretreated daily with intraperitoneal injections of 60 mg/kg phenobarbital for three days. Two days after the final injection, the animals were challenged with pentobarbital or hexobarbital and the sleep responses were examined.

As shown in Table 1, the treated rats were found to be strongly resistant to the challenge dose (35 mg/kg) of pentobarbital that produces moderate sleeping in nontolerant rats. Brain concentrations of pentobarbital at ten minutes after administration showed no significant difference between the control and treated rats. On the contrary, the median anesthetic dose in the treated rats was found to be about twice as much as that in nontreated rats.

TABLE 1. Decreased sensitivity to pentobarbital in rats after daily injections of 60 mg/kg phenobarbital for three days.

Treatment	Number of animals anesthetized*	Pentobarbital in brain** ($\mu\text{g/g} \pm \text{S.E.}$)		AD 50*** (mg/kg)
		10 min	180 min	
None	8 (8)	34.6 ± 5.1	18.4 ± 2.0	23.2
Phenobarbital	7(30)	33.5 ± 4.6	14.2 ± 2.3	47.7

* Values in parentheses indicate the number of experimental animals. Loss of righting reflex after intraperitoneal injection of pentobarbital (35 mg/kg) was used as criterion.

** Determined by the method of Brodie (2). Each value represents an average of 6 animals $\pm$ S.E.

*** The median anesthetic dose (AD 50) was estimated by up and down method of Dixon-Mood.