STUDIES ON PHARMACODYNAMIC ACTION OF
N-ETHYLMALIMIDE (NEM)
1ST REPORT: INFLUENCE OF NEM ON SYNAPTIC TRANSMISSION
OF SYMPATHETIC NERVES IN GUINEA PIGS

Hiroyuki YAMAMOTO, Takafumi KITANO, Hitoo NISHINO
and Tadashi MURANO

Department of Pharmacology, Wakayama Medical College, Wakayama, Japan

Accepted September 14, 1972

Abstract—In isolated guinea pig atrial preparations, NEM in the concentration of 5×10^{-5} M to 10^{-4} M produced an increase in the amplitude and rate of contraction at earlier stages, resulting in inhibition of contraction accompanied by an increase in resting tension and irregularity in rate at later stages, while the diphasic actions of NEM were completely antagonized by cysteine. The initial action of NEM was blocked by propranolol, guanethidine and failed to appear in the reserpinized atria, but reappeared after incubation of atria with norepinephrine indicating that it is mediated by norepinephrine. NEM augmented the contractile response of ductus deferens to hypogastric nerve stimulation at the earlier stages, but suppressed it at later stages. The augmenting effect of NEM on the response to sympathetic nerve stimulation was dependent on calcium concentrations in suspending medium. Effects of NEM on synaptic transmission of sympathetic nerves were discussed.

NEM has greater selectivity for accessible-SH groups on protein molecules and passes easily through cell membranes (1-2). For these reasons, pharmacological actions of NEM were expected to differ from those of p-chloromercuribenzoate (PCMB) or p-chloromercuribenzenesulphonate (PCMBs) (3-5). Utilizing the characteristic property of NEM several authors have verified that sulfhydryl-containing proteins play a vital role at least in the contraction of muscles (6-8).

The present work was undertaken to determine how the SH group is involved in the regulation or control of synaptic transmission of sympathetic nerve in guinea pigs atria and ductus deferens.

METHODS AND MATERIALS

1) Atrial preparation

Male guinea pigs weighing 300-500 g were used. The animals were sacrificed by exsanguination. The thorax was incised and the entire heart removed. The ventricular muscle, fat and connective tissue were discarded. Conventional atrial preparations were suspended in Krebs' bicarbonate solution aerated with a mixture of 95% O₂ and 5% CO₂ at $36 \pm 1^\circ\text{C}$.

The amplitude and rate of spontaneous atrial contraction were isometrically recorded via a force-displacement transducer. The concentration of compounds previously adjusted at pH 7.4 with NaOH and added to the bath was expressed in M.

2) *Ductus deferens-hypogastric nerve preparation*

Male guinea pigs weighing 300–500 g were used. The ductus deferens with hypogastric nerve was prepared by the method of Hukovic (9). The hypogastric nerve was placed on a pair of platinum electrodes located above the meniscus of the bath fluid and kept wet throughout the experiment. The hypogastric nerves were stimulated submaximally with the rectangular pulse, 2 msec in duration and 10, 20 and 40/sec in frequency and the ductus deferens was also stimulated transmurally with the rectangular pulse, 2 msec in duration and 10/sec in frequency. The stimuli were delivered for 5 sec every 3 min using a Nihon Kohden stimulator (MSE-3).

Preparations were suspended in Krebs' bicarbonate solution at $30 \pm 1^\circ\text{C}$ bubbled with 95% O_2 and 5% CO_2 .

3) *Isolated perfused bull-frog spinal cord preparations*

Bull-frogs weighing 100–200 g were used. The preparations were made according to the method of Matsuura (10).

The dorsal root of spinal cord was stimulated submaximally with rectangular pulse, 1 msec in duration and the spinal cord discharge was recorded by oscilloscope from the ventral root.

RESULTS

A) *Ino- and chronotropic responses of atria to NEM*

1) *Effect of NEM on isolated atria*

Although NEM in concentrations of less than 10^{-6} M showed no significant changes in the atrial contraction, $10^{-5} \sim 5 \times 10^{-5}$ M of NEM produced positive ino- and chronotropic effects lasting 20–30 min (Fig. 1). When the concentration was further increased to 10^{-4} M, the drug enhanced the atrial contraction approx. twice that of control immediately after application, followed by a progressive diminution in the amplitude of contraction 10 to 15 min later. There appeared an increase in resting tension and an irregularity of atrial rate in 30 min, at which time these unusual states were no longer restorable by washing.

On the other hand, the combined use of 10^{-4} M of cysteine with NEM (10^{-4} M) resulted in a complete suppression of NEM action, though cysteine itself did not reveal any significant effect on atrial motilities (Fig. 1). Furthermore, among NEM related compounds, maleimide had similar effects, but not so marked as NEM, while maleate had no effect on atria.

2) *Effects of propranolol and of reserpinization on the action of NEM*

Pretreatment with $10^{-6} \sim 10^{-5}$ M of the β -adrenergic blocker, propranolol, for 3 min not only significantly reduced the enhancement by NEM (10^{-4} M) of atrial contraction, but also prevented almost entirely, increases in resting tension and irregularities in the

FIG. 1. Positive ino- and chronotropic responses of guinea pigs atria to NEM.

normal atria

reserpinized atria

pre-incubation for 1hr with noradrenaline

FIG. 2. Effects of propranolol and reserpinization on action of NEM.

atrial rate, which had occurred at later stages (Fig. 2).

Furthermore, atria were obtained from guinea pigs which had been treated i.p. with 1 mg/kg of reserpine 24 hr previously. In such preparations, tyramine showed no significant effects.

Although the initial promoting effect of NEM on the amplitude of atrial contraction as above noted was no longer produced, the inhibitory effect occurring at later stages still appeared in the reserpinized atria. When incubating the reserpinized atria with 10^{-5} M of norepinephrine for 1 hr, then washing out the remaining norepinephrine from the suspension bath, initial promoting effect of NEM was reproduced.

From these findings, it can be presumed that norepinephrine in atria is necessary for exhibition of the promoting effect of NEM in earlier stages.

For further study of initial action of NEM on atria, effects of tyramine or of nicotine mediated by norepinephrine were compared with those of NEM.

FIG. 3. Enhancement by cocaine of initial action of NEM (5×10^{-5} M).
Numbers indicate atrial rate per min.
Time scale indicates 1 min.

FIG. 4. Influence of C_6 treatment on actions of NEM and nicotine.

3) Influence of cocaine, guanethidine and hexamethonium (C_6) on the initial action of NEM

In untreated atrial preparations, 10^{-5} M of tyramine produced an increase in amplitude and rate of atrial contraction, but the action was significantly reduced by pretreatment with 10^{-6} M of cocaine for 3 min (11, 12). On the contrary, the initial action of NEM was rather enhanced by the same pretreatment (Fig. 3).

Reversely, the promoting effect of tyramine was significantly diminished on guinea pigs atrial preparations, which had been administered guanethidine i.p. 30 mg/kg 1 hr prior to sacrifice. The action of NEM was not further reduced to any degree in these preparations than it was in the case of tyramine. The positive inotropic action of nicotine was almost completely blocked by pretreatment with 5×10^{-5} M of C_6 for 3 min, while the action of NEM was unaffected by the same pretreatment (Fig. 4).

B) Effects of NEM on responses of ductus deferens of guinea pig to hypogastric nerve stimulation

1) Effects of NEM on responses to nerve stimulation

Application of NEM in the concentration of 5×10^{-5} M gradually inhibited the response of ductus deferens to hypogastric nerve stimulation at the frequency of 40/sec, although the inhibition was easily reversed when the preparation was washed out (Fig. 5). When increased up to 10^{-4} M, the response was significantly suppressed in 10–20 min and then abolished in 30–40 min, small spontaneous movements being appeared. Contractions due to norepinephrine (10^{-5} M), Ach 5×10^{-5} M or KCl (10^{-2} M) could be evoked immediately after suppression by NEM of the response to nerve stimulation (Fig. 6).

Next, effects of NEM on the responses to nerve stimulation at the frequencies of 10,

FIG. 5. Effects of NEM on responses of isolated guinea pigs ductus deferens to hypogastric nerve stimulation.

FIG. 6. Response of ductus deferens to norepinephrine, acetylcholine and potassium chloride immediately after suppression by NEM of ductus deferens responses to nerve stimulation.

FIG. 7. Effects of NEM on ductus deferens responses to nerve stimulation at different frequencies.

● ; 10/sec, × ; 20/sec, △ ; 40/sec

20 and 40/sec were comparatively studied. At the initial stage of application, NEM in the concentration of 7.5×10^{-5} M augmented the contractile response of the preparation to nerve stimulation at a lower frequency (10/sec) by approx. 30%, but at 20/sec, the degree of the augmentation was slight, while at 40/sec, augmentation was unobservable (Fig. 7). The depressive effect of NEM on contraction was clearer when a higher frequency stimulation was administered.

2) Influence of calcium on the action of NEM

Effect of NEM on co-relationship between magnitude of the response at lower frequency (10/sec) and concentration of calcium in suspending medium was observed. The result showed that NEM potentiated the calcium enhancement of response in proportion to the concentration of calcium, in the range from 0.63 to 5 mM; NEM augmented the contractile response to nerve stimulation at a lower frequency by approx. 15% and 30% in 1.25 mM and 2.5 mM of calcium concentration, respectively, although an augmentation by the drug was not seen in 0.63 mM of calcium (Fig. 8).

FIG. 8. Influence of NEM on enhancement by calcium of ductus deferens responses to hypogastric nerve stimulation.

(○—○ NEM free, ●—● NEM 7.5×10^{-5} M. Vertical bar represents standard error)

FIG. 9. Effects of NEM on ductus deferens responses to transmural or hypogastric nerve stimulations.

N : response to nerve stimulation

TM : response to transmural stimulation

C) *Effects of NEM on the responses of ductus deferens of guinea pig to transmural stimulation*

At the initial stage of application, NEM in the concentration of 7.5×10^{-5} M produced a less enhancement of contractile response to transmural stimulation (10/sec). In the case of transmural stimulation, the enhancement was followed by progressive diminution of the response, milder than that seen in nerve stimulation (Fig. 9).

Response to transmural stimulation was depressed 10–20% by C_6 and NEM enhancement of transmural stimulation was also diminished slightly by C_6 .

From these findings (sections B and C), the theory is favored that NEM acts not only on sympathetic nerve terminals but also on ganglion cells (13–16).

D) *Effects of NEM on isolated bull-frog spinal cord in perfusion*

Although NEM below a concentration of 10^{-5} M affected neither mono- nor polysynaptic discharges of the spinal cord, 5×10^{-5} M of the drug increased the amplitude of mono-synaptic responses by approx. 30%, reaching a maximum level in 30 min, followed by a gradual recovery to the normal state in 60 min (Fig. 10).

When increased to 10^{-4} M, the promoting effect of NEM was so remarkable that the intensity of promotion reached approx. 60% in 30 min, followed by depression of the response 60 min after application. With a concentration above 5×10^{-4} M of NEM, a transient promotion of the response occurred, followed by marked depression. The depressive effect of the drug was no longer reversed by repeated washing.

FIG. 10. Effects of NEM on monosynaptic discharge of isolated bull-frog spinal cord in perfusion.

(●—● 5×10^{-5} M, ○—○ 10^{-4} M, △—△ 5×10^{-4} M, ▲—▲ 10^{-3} M)

DISCUSSION

In the isolated guinea pig atrial preparations, NEM in the concentration of 5×10^{-5} M $\sim 10^{-4}$ M produced an enhancement in the amplitude and rate of contraction at an earlier stage of application, resulting in an increase of resting tension and an irregularity in rate at later stages.

The combined use of cysteine with NEM produced neither initial effects of the drug nor depression at later stages, indicating that the effect of NEM is due to reactivity with the SH groups.

On the other hand, the initial promoting effect of NEM was blocked by pretreatment with adrenergic β -blocker, propranolol, nor was it observable in the reserpinized atria. Restoration of the initial effect by incubation of reserpinized atria with norepinephrine was obvious. It was also demonstrated that i.p. administration of NEM significantly decreased the epinephrine content in adrenal medulla and caused marked hyperglycemia in rats (17).

From these experiments, it is presumed that the appearance of initial effects of NEM are mediated by catecholamine released from stored sites in atrial tissues.

Consequently, comparing the initial action of NEM with that of tyramine producing positive inotropic action via norepinephrine released from sympathetic nerve terminals, the promoting effects of both NEM and tyramine were reduced by pretreatment with guanethidine, while the action of tyramine was antagonized by cocaine but that of NEM augmented. From these findings, it is suggested that cocaine does not interfere with noradrenaline release due to NEM and inhibits the re-uptake of catecholamine released by NEM, but blocks the uptake of tyramine itself.

Furthermore, in the sympathetically innervated ductus deferens preparations, NEM produced a transient augmentation of the response of ductus deferens to nerve stimulation at lower frequencies in earlier stages, resulting in the diminution of the response at later stages. Moreover, the diphasic effects of NEM were more prominent on the response to hypogastric nerve stimulation than to transmural stimulation.

It is generally accepted that impulses of hypogastric nerve stimulation are mainly transmitted to effector cells via ganglion, while transmural stimuli directly stimulates sympathetic nerve terminals. Thus, it is suggested that locus of actions of NEM are not only in sympathetic nerve terminals or in stored sites of catecholamine but also in ganglion, the latter being more sensitive to NEM than the former.

Consequently, the enhancement by NEM of the response to hypogastric nerve stimulation is more remarkable than that to transmural stimulation and even when suppression in ganglion cells begins, the augmenting effects on sympathetic nerve terminals still remain unaltered, resulting in a milder diminution of the response to transmural stimulation than that to hypogastric nerve stimulation. This assumption above noted is compatible with Komalahiranya et al's report (18).

Hasselbach and Taugner (19) and Kirsten and Kuperman (6) have already suggested that the primary site of NEM action is inhibition of calcium uptake by vesicles of the

sacroplasmic reticulum, on the outer surface in which SH groups are located. For this reason the NEM augmentation of enhancement by calcium should be given due consideration.

REFERENCES

- 1) JACOB, H.S. AND JANDEL, J.H.: *J. clin. Invest.* **41**, 779 (1962)
- 2) WEBB, J.L.: *Enzyme and Metabolic Inhibitors*, Vol. 3, p. 337, Academic Press, New York (1966)
- 3) OKAMOTO, M. AND KUPERMAN, A.S.: *Nature* **210**, 1062 (1966)
- 4) KIRSTEN, E.B. AND KUPERMAN, A.S.: *Br. J. Pharmacol. Chemother.* **40**, 814 (1970)
- 5) IWATA, H., YAMAMOTO, I. AND OKA, M.: *Japan. J. Pharmacol.* **18**, 471 (1968)
- 6) KIRSTEN, E.B. AND KUPERMAN, A.S.: *Br. J. Pharmacol. Chemother.* **40**, 827 (1970)
- 7) TODA, N. AND KONISHI, N.: *European J. Pharmacol.* **7**, 5 (1969)
- 8) GOODMAN, I. AND HIATT, R.B.: *Biochem. Pharmacol.* **13**, 871 (1964)
- 9) HUKOVIC, S.: *Br. J. Pharmacol. Chemother.* **16**, 188 (1961)
- 10) MATSUURA, S.: *Osaka City Med. J.* **15**, 29 (1969)
- 11) TRENDLENBURG, U.: *J. Pharmacol. exp. Ther.* **134**, 8 (1961)
- 12) FARMER, J.B. AND PETCH, B.: *J. Pharm. Pharmacol.* **15**, 639 (1963)
- 13) OHLIN, P. AND STROMBLAD, B.C.: *Br. J. Pharmacol. Chemother.* **20**, 299 (1963)
- 14) YAMAMOTO, H. AND NAKANISHI, H.: *Japan. J. Pharmacol.* **14**, 256 (1964)
- 15) SAKUMA, A., NONOMURA, Y., KANAMURA, H. AND OTSUKA, M.: *Japan. J. Pharmacol.* **15**, 357 (1965)
- 16) KURIYAMA, H.: *J. Physiol.* **169**, 213 (1963)
- 17) YAMAMOTO, H., MASUDA, Y., OKADA, N. AND MURANO, T.: *Japan. J. Pharmacol.* (in press)
- 18) KOMALAHIRANYA, A. AND VOLLE, R.L.: *J. Pharmacol. exp. Ther.* **139**, 304 (1963)
- 19) HASSELBACH, W. AND TAUGNER, G.: *Biochem. J.* **119**, 265 (1970)