STUDIES ON ANTINEPHRITIC EFFECT OF MIZORIBINE (p-INN, BREDININ®), A NEW IMMUNOSUPPRESSIVE AGENT, AND AZATHIOPRINE (1) EFFECT ON THE NEPHROTIC TYPE OF ANTI-GBM NEPHRITIS IN RATS

Kyoko OKAMOTO** **, Mikio ITO* and Yoshio SUZUKI*

*Department of Pharmacology, Faculty of Pharmacy,
Meijo University, Tenpaku-ku, Nagoya 468, Japan

**Research Laboratories, Toyo Jozo Co., Ltd, Tagata-gun,
Shizuoka 410-23, Japan

Accepted January 20, 1983

Abstract—The present study was made to investigate the antinephritic effect of mizoribine in comparison to that of azathioprine by using the nephrotic type of anti-rat glomerular basement membrane rabbit serum (anti-GBMserum)-induced nephritis in rats. The nephrotic type nephritis was induced in rats by two i.v. injections of anti-GBM serum at a 10 day interval. Both drugs were given orally, daily from the 2nd day following the first injection of anti-GBM serum to the 21st day. Mizoribine in doses of 5 and 7.5 mg/kg/day significantly inhibited urinary protein excretion by 30–40% on the 22nd day. Mizoribine at both doses showed an inhibitory tendency on urinary alkaline phosphatase (ALP) excretion on the 9th and 16th days. On the 22nd day, this drug at a dose of 7.5 mg/kg/day inhibited plasma cholesterol (CL) content by 59.6% and wet weight of kidneys by approx. 50%, but no significant difference was seen between the drug-treated and control groups. When renal tissues on the 22nd day were observed under light microscopy, mizoribine at both doses remarkably prevented glomerular changes such as mesangial proliferation and thickening of capillary walls and significantly inhibited the index of glomerular lesions (IGL) by over 60%. On the other hand, azathioprine at a dose of 20 mg/kg/day was as effective as mizoribine at 5 mg/kg/day in inhibiting urinary protein and ALP excretions, plasma CL content and kidney weight. However, azathioprine showed little effect on the IGL. From these results, mizoribine at the dose level of 1/4 to 1/3 of azathioprine showed a more potent effect than azathioprine in this model. Therefore, mizoribine is also expected to have a beneficial effect on nephrotic type nephritis in clinical fields.

It has been considered that an immunological mechanism may be involved in the initiation and progression of glomerulonephritis. In this connection, Germuth and Rodriguez (1) reported that more than 80% of renal diseases might be produced by triggering immunologic reactions. Practically, in clinical fields, immunosuppressive agents such as azathioprine, cyclophosphamide and 6-mercaptopurine have been widely used for

the treatment of glomerulonephritis. However, there are only a few reports regarding the effects of immunosuppressive agents on experimental glomerulonephritis. Feenstra et al. (2) reported that azathioprine showed a beneficial effect on heterologous immune complex glomerulonephritis in rats only when administered before or simultaneously with the onset of nephritis. Hurd (3) reported that treatment with cyclophosphamide pre-

vented interstitial nephritis and tubular cell proliferation in NZB/NZW mice. Contrary to those reports, Lim and Spargo (4) were unable to demonstrate any beneficial effects of 6-mercaptopurine and cyclophosphamide on experimental autologous immune complex glomerulonephritis in rats.

Mizoribine (p-INN, Bredinin®) [4-carbamoyl-1- β -D-ribofuranosyl-imidazolium-5-olate], a new immunosuppressive agent, which is an imidazole nucleoside isolated from the culture supernatant of *Eupenicillium brefeldianum*, has been shown to prevent the proliferation of lymphocytes in vitro (5) and to possess an immunosuppressive action in vivo (6). An application of this drug to renal transplantation has been recently shown to be apparently successful (7). Therefore, the present study was designed to estimate the antinephritic effect of mizoribine in comparison to that of azathioprine by using the nephrotic type of anti-rat glomerular basement membrane rabbit serum (anti-GBM serum)-induced nephritis of rats, which closely resembles human nephrotic syndrome (8).

Materials and Methods

Drugs: The drugs used in this experiment were mizoribine (Toyo Jozo Co., Ltd.) (Fig. 1) and azathioprine (Tanabe Seiyaku Co., Ltd.).

Experimental nephritis: Experimental nephritis was induced in male Wistar rats weighing approx. 180 g by the i.v. injection of 0.45 ml/animal of anti-GBM serum by the

method reported previously (8). In order to induce a nephrotic type of nephritis, 0.2 ml of anti-GBM serum per animal was again administered i.v. on the 10th day after the first injection.

Evaluation of the anti-nephritic effect of drugs: The 24 hr urine samples from the 1st to the 2nd day after the first injection of anti-GBM serum were collected and the rats divided into 4 groups, each consisting of 10 animals on the basis of the degree of proteinuria. Three groups were given p.o. daily 5 and 7.5 mg/kg of mizoribine and 20 mg/kg of azathioprine, respectively, from the 2nd day after the first injection of anti-GBM serum to the 21st day. In this case, mizoribine was dissolved in 0.9% saline and azathioprine suspended in saline solution of 1% gum arabic. Each drug was given in a volume of 1 ml/100 g of body weight. The remaining one group was given 0.9% saline p.o. daily instead of test drugs for periods of drug administration and served as the control. In addition to the above groups, a non-nephritic (normal) group was set in the experiment and was given 0.9% saline p.o. daily as was done with the control group. The 24 hr urine samples from the 8th to the 9th day, the 15th to the 16th day and the 21st to the 22nd day following the first injection of anti-GBM serum were collected. On the 22nd day, blood was drawn for determination of biochemical parameters in the plasma, and kidneys were taken immediately for histopathological studies. Evaluation of the antinephritic effect of the test drug was done by comparing biochemical parameters of urine and plasma and histopathological findings in kidneys of the drug-treated group with those of the control group.

Urine and blood collections: The 24 hr urine samples were obtained by keeping each animal in an individual metabolic cage for 24 hr. During the urine collection, each animal received 5 ml of distilled water twice

Fig. 1. Structure of mizoribine

orally without feeding. The urine thus obtained was then centrifuged at 3,000 rpm for 20 min at 4°C, and the supernatant was used for urinalysis. After the last collection of urine samples, each animal was lightly anesthetized with ether, and the blood was taken from the abdominal aorta into a centrifuge tube with the ratio of 9 parts blood to 1 part 3.13% trisodium citrate. The blood was then centrifuged at 3,000 rpm for 10 min at 4°C to obtain plasma.

Determination of biochemical parameters of urine: The protein content in urine was determined by the method of Kingsbury et al. (9) and expressed as mg/24 hr urine. As urinary enzymes, alkaline phosphatase (ALP) activity was determined by the method of Bessey et al. (10) by using disodium *p*-nitrophenyl phosphate (Daiichi Kagaku Co., Ltd.) as substrate. N-acetyl- β -glucosaminidase (NA- β -G) activity was determined according to the method of Hasebe (11) by using *p*-nitrophenyl-N-acetyl- β -D-glucosaminide (BDH Chemicals Co., Ltd., Poole, England) as substrate. Both enzymatic activities were expressed as mg of liberated *p*-nitrophenol/hr/24 hr urine.

Determination of biochemical parameters of plasma: Plasma urea nitrogen (UN) and cholesterol (CL) contents were determined by means of the method of diacetyl monoxime (12) and Zurkowski (13), respectively. Both results were expressed as mg/dl of plasma.

Kidney weight and assessment of histopathological preparations of kidneys: Kidneys were weighed and the result was expressed as mg of the mean of wet weight of both kidneys/100 g of body weight. For light microscopic observation, the kidneys weighed were fixed in 10% buffered formalin, embedded in paraffin and sectioned 3- μ thick. The sections were stained with hematoxylin and eosin (HE) and Masson trichrome (MT). As previously reported (14), the degree of glomerular lesions was assessed by light

microscopy, each glomerulus being scored from 0–2 as follows: Score 0—Normal glomerulus. Score 1—Although mild mesangial proliferation is seen, the capillary structures are well preserved. Score 2—In addition to moderate mesangial proliferation, there is partial destruction of capillary structures. According to the above criteria, 50 glomeruli/rat were observed. The number of glomerulus corresponding to each score was represented as n_0 , n_1 , and n_2 , and the index of glomerular lesions (IGL) was calculated as follows:
$$\frac{(0 \times n_0) + (1 \times n_1) + (2 \times n_2)}{n_0 + n_1 + n_2}$$

Statistical analyses: The data represented the mean $\pm$ S.E., and the results were statistically evaluated by the Kruskal-Wallis test and multiple comparison except for the normal group.

Results

1. Influences of mizoribine and azathioprine on body weight gain

The average body weight of all groups at the beginning of the experiment was 180 ± 2.1 g. However, on the 22nd day following the first injection of anti-GBM serum (the day when all animals were killed), weights of normal-; control-; mizoribine, 5 mg/kg/day p.o.-treated-; mizoribine, 7.5 mg/kg/day p.o.-treated-; and azathioprine, 20 mg/kg/day p.o.-treated groups were 275.6 ± 6.8 , 224.7 ± 4.0 , 212.8 ± 10.9 , 204.4 ± 6.5 and 206.4 ± 10.3 g, respectively. No significant difference in the gain in body weight between each drug-treated group and the control group was observed, although the gains in the mizoribine high dose- and azathioprine-treated groups were somewhat lower than that of the control group.

2. Effects of mizoribine and azathioprine on biochemical parameters of urine and plasma and kidney weight

Urinary protein excretion: As shown in Fig. 2, the average 24 hr proteinuria values of

Fig. 2. Effects of mizoribine and azathioprine on urinary protein excretion in the nephrotic type of anti-GBM nephritis in rats. Each column represents the mean value with S.E. obtained from 10 rats. The numbers in parentheses are the percent inhibition which is derived from the following formula: $\frac{C-T}{C-N} \times 100$ (C: Control, T: Test drug and N: Normal). * indicates a significant difference from the control at $P < 0.05$.

the control group on the 2nd day (the starting day of each drug treatment), the 9th day (the day before the second injection of anti-GBM serum), the 16th day and the 22nd day following the first injection of anti-GBM serum were 150.2 ± 25.9 , 213.4 ± 25.6 , 502.2 ± 38.4 and 599.3 ± 33.0 mg, respectively. On the 16th day, both drugs inhibited the protein excretion to the same extent as that in the case of the 22nd day, but no significant difference between each drug-treated group and the control group was recognized.

Urinary enzyme excretions: The results are illustrated in Fig. 3. Mizoribine (5 and 7.5 mg/kg/day p.o.), as well as azathioprine (20 mg/kg/day p.o.) only had an inhibitory tendency on the ALP excretion on the 9th and 16th days. Both drugs were ineffective in inhibiting the NA- β -G excretion on any days determined.

Plasma UN and CL contents: The results on plasma UN and CL contents on the 22nd day are indicated in Fig. 4. The UN content

of the control group was only 1.6 times as high as that of the normal group. There was no significant difference in the UN level between each drug-treated group and the control group. On the other hand, the CL level of the control group was 2.5 times higher than that of the normal group. Mizoribine at 7.5 mg/kg/day p.o. and azathioprine at 20 mg/kg/day p.o. inhibited the CL level by 59.6 and 45.9%, respectively, but no significant difference from the control was seen.

Kidney weight: The results on kidney weight on the 22nd day are given in Fig. 4. The average wet weight of both kidneys/100 g of body weight of the control group was 553 ± 41 mg, being 72.3% heavier than that of the normal group. Mizoribine at 7.5 mg/kg/day p.o. brought about approx. 50% inhibition on the kidney weight, but there was no significant difference from the control. Mizoribine at 5 mg/kg/day p.o. and azathioprine at 20 mg/kg/day p.o. also showed

an inhibitory tendency.

3. Histopathological findings of kidneys

treated with mizoribine and azathioprine

When renal tissues on the 22nd day were

Fig. 3. Effects of mizoribine and azathioprine on urinary alkaline phosphatase and N-acetyl- β -glucosaminidase excretions in the nephrotic type of anti-GBM nephritis in rats. Each column represents the mean value with S.E. obtained from 10 rats. The numbers in parentheses are the percent inhibition which is derived from the following formula: $\frac{C-T}{C-N} \times 100$ (C: Control, T: Test drug and N: Normal).

Fig. 4. Effects of mizoribine and azathioprine on plasma urea nitrogen and cholesterol contents in the nephrotic type of anti-GBM nephritis in rats. Blood was taken on the 22nd day after the first injection of anti-GBM serum. Each column represents the mean value with S.E. obtained from 10 rats. The numbers in parentheses are the percent inhibition which is derived from the following formula: $\frac{C-T}{C-N} \times 100$ (C: Control, T: Test drug and N: Normal).

Table 1. Effects of mizoribine and azathioprine on index of glomerular lesions in the nephrotic type of anti-GBM nephritis in rats

Groups	Index of glomerular lesions	% Inhibition
Control	0.106±0.021	
Mizoribine 5 mg/kg/day p.o.	0.039±0.005*, ‡	63.2
Mizoribine 7.5 mg/kg/day p.o.	0.032±0.008*, §§	69.8
Azathioprine 20 mg/kg/day p.o.	0.101±0.009	4.7

Kidneys were taken on the 22nd day after the first injection of anti-GBM serum. Results are the mean ±S.E. obtained from 10 rats. % Inhibition = $\frac{C-T}{C} \times 100$ (C: Control and T: Test drug). *indicates a significant difference from the control at $P < 0.05$. ‡ and §§ indicate a significant difference from azathioprine at $P < 0.05$ and 0.01 , respectively.

observed under light microscopy, the glomeruli of the control group were characterized by mild proliferation of mesangium and slight thickening of capillary walls (Photo 1-a). The results of effects on glomerular histopathological changes, quantitatively expressed as IGL, are shown in Table 1. The IGL of the control group was 0.106 ± 0.021 . Mizoribine at 5 and 7.5 mg/kg/day p.o. showed a potent inhibition of over 60% on the IGL. The glomeruli from animals treated with this drug were similar to that from the normal (Photo 1-b). However, azathioprine (20 mg/kg/day p.o.) had no effect on the IGL.

Discussion

In the present paper, the antinephritic effect of mizoribine was studied in comparison to that of azathioprine by using the nephrotic type of nephritis induced by two injections of anti-GBM serum at a 10 day interval into rats. We previously reported that two injections of anti-GBM serum caused more prominent nephrotic symptoms about one week following the second injection when compared to the case of one shot of anti-GBM serum (15). In the present experiment, the control group showed a urinary protein excretion of approx. 600 mg/day and

(a)

(b)

Photo 1. Kidney sections from control rat (a) and mizoribine, 7.5 mg/kg/day p.o.-treated rat (b) on the 22nd day after the first injection of anti-GBM serum. MT stain (a: $\times 200$, b: $\times 250$).

a plasma CL level of approx. 140 mg/dl (2.5 times higher than normal level) on the 22nd day following the first injection of anti-GBM serum. Thus, massive proteinuria and hypercholesterolemia characteristic of nephritis were observed in the control group.

In the preliminary experiment, we studied the effective dose of mizoribine. In this case, mizoribine was ineffective at the dose of 2.5 mg/kg/day p.o. On the other hand, with a 10 mg/kg/day p.o. dose of this drug, there was no gain in body weight during the treatment period, although this dosage resulted in a potent antinephritic effect. Therefore, in the present experiment, we used this drug at doses of 5 and 7.5 mg/kg/day p.o. Mizoribine at both doses remarkably inhibited not only urinary protein excretion but also histopathological changes in glomeruli evaluated by means of IGL. Moreover, this drug at both doses showed an inhibitory tendency on urinary ALP excretion, plasma CL content and kidney weight. On the other hand, azathioprine at 20 mg/kg/day p.o. was as effective as mizoribine at 5 mg/kg/day p.o. on urinary protein and ALP excretions, plasma CL content and kidney weight. However, azathioprine had little effect on the IGL. Thus, mizoribine at the dose level of 1/3 to 1/4 that of azathioprine showed more a potent antinephritic effect than azathioprine.

Recently, it has been revealed by Okubo et al. (16) that mizoribine significantly exerts a prolonged effect on the life of (NZB/NZW) F₁ mice with SLE nephropathy. They suggest that the effect of this drug may be due to suppression of antibody formation via the direct inhibition of B cell function. Possible mechanisms by which the enhancement of nephrotic symptoms is induced by two injections of anti-GBM serum are the following (15): One of the mechanisms may be due to the increased amount of anti-rat GBM rabbit antibody fixed to rat GBM by

reinjection of anti-GBM serum. Another mechanism is thought to be due to the augmentation of an immune response in the autologous phase through host antibody formation against rabbit γ -globulin in the reinjected anti-GBM serum. Therefore, it would appear that the mechanism of the antinephritic action of mizoribine may be explained by preventing the autoimmune response via the suppression of the auto-antibody formation. It has been shown that unlike azathioprine, mizoribine possesses few side effects such as the depression of bone marrow function and hepatic toxicity (7). In view of our results and the above reports on side effects, mizoribine is also expected to have a beneficial effect in clinical application.

References

- 1) Germuth, F.G., Jr. and Rodriguez, E.: Immunopathology of the Renal Glomerulus: Immune Complex Deposit and Antibasement Membrane Disease. p. 228. Little Brown, Boston (1973)
- 2) Feenstra, K., Lee, R.v.d., Greben, H.A., Arends, A. and Hoedemaeker, Ph. J.: Experimental glomerulonephritis in the rat induced by antibodies directed against tubular antigen II. Influence of medication with prednisolone and azathioprine: A histologic and immunohistologic study at the light microscopic and the ultrastructural level. *Lab. Invest.* **32**, 243–250 (1975)
- 3) Hurd, E.R.: Effect of cyclophosphamide on interstitial nephritis and tubule cell proliferation in NZB/NZW mice. *J. Immunol.* **119**, 1552–1555 (1977)
- 4) Lim, V.S. and Spargo, B.H.: Immunosuppressive treatment of autologous immune complex nephritis in rats. *J. Lab. Clin. Med.* **81**, 661–670 (1973)
- 5) Sakaguchi, K., Tsujino, M., Yoshizawa, M., Mizuno, K. and Hayano, K.: Action of bredinin on mammalian cells. *Cancer Res.* **35**, 1643–1648 (1975)
- 6) Mizuno, K., Tsujino, M., Takada, M., Hayashi, M., Atsumi, K., Asano, K. and Matsuda, T.: Studies on bredinin. I. Isolation, characterization and biological properties. *J. Antibiot. (Tokyo)* **27**, 775–782 (1974)
- 7) Inou, T., Kusaba, R., Takahashi, I., Sugimoto, H.,

- Kuzuhara, K., Yamada, Y., Yamauchi, J. and Otsubo, O.: Clinical trial of bredinin in renal transplantation. *Transplant. Proc.* **13**, 315–318 (1981)
- 8) Ito, M., Nagamatsu, T. and Suzuki, Y.: Pharmacological studies on experimental nephritic rats (10). Changes in coagulation-fibrinolysis system in the course of anti-GBM induced nephritis. *Japan. J. Nephrology* **23**, 297–308 (1981) (Abs. in English)
- 9) Kingsbury, F.B., Clark, C.P., Williams, G. and Post, A.L.: The rapid determination of albumin in urine. *J. Lab. Clin. Med.* **11**, 981–989 (1926)
- 10) Bessey, O.A., Lowry, O.H. and Brock, M.J.: A method for the rapid determination of alkaline phosphatase with five cubic millimeters of serum. *J. Biol. Chem.* **146**, 321–329 (1946)
- 11) Hasebe, K.: Biochemical studies on synovial fluid. (1) Mucopolysaccharase activity in synovial fluid in rheumatoid arthritis. *Fukushima J. Med. Sci.* **15**, 35–44 (1968)
- 12) Searcy, R.L. and Cox, F.M.: A modified technique for ultramicro estimations of urea nitrogen. *Clin. Chim. Acta* **8**, 810–812 (1963)
- 13) Zurkowski, P.: A rapid method for cholesterol determination with a single reagent. *Clin. Chem.* **10**, 451–453 (1964)
- 14) Suzuki, Y., Ito, M., Ogawa, Y. and Nagamatsu, T.: Pharmacological studies on experimental nephritic rats (6). Antinephritic effects of sodium chondroitin sulfate and other drugs on modified type of Masugi's nephritis. *Folia Pharmacol. Japon.* **75**, 585–599 (1979) (Abs. in English)
- 15) Suzuki, Y., Ito, M., Kito, T. and Nagamatsu, T.: Pharmacological studies on experimental nephritic rats (12). Nephritic model induced by the repeated administration of anti-rat GBM rabbit serum. *Pharmacometrics* **21**, 503–513 (1981) (Abs. in English)
- 16) Okubo, M., Kamata, K., Masaki, Y. and Uchiyama, T.: Effect of bredinin (BR) on lupus nephropathy of New Zealand Black/White F₁ hybrid mouse (B/WF₁). *Kidney Int.* **21**, 203 (1982)