THE BOTANICAL MAGAZINE.

VOL. X, No. 109.

PART II.

On the Different Views hitherto Proposed regarding the Morphology of the Flowers of *Ginkgo biloba*, L.

(*Preliminary note.*)

By

Kenjiro Fujii, *Rigakushi*.

With Pl. I.

(*Continued from* No. 108, p. 8.)

Provisional diagnosis of Ginkgo:—a deciduous tree, with “Langtriebe” and “Kurztriebe”; foliage leaves fan-shaped, dichotomously veined, with the upper margin wavy, usually bilobed, sometimes many-lobed in young trees, entire in flowering branches. Flowers⁽¹⁾ dioecious, single, in the axil of the uppermost scaly leaves and the lowermost foliage leaves; male flowers in the form of a loose, elongated, short stalked spike; stamens with two pollensacs and a very small round terminal-scale (“Endschuppen”); female flowers long-stalked, usually with two ovules placed opposite to each other near the apex of the stalk; ovules with a single integument and a 2-lipped micropyle, partially enclosed at the base in a small cup-shaped swelling, usually one ovule only in each flower becoming ripe seed; seeds drupe-like, the stony part 2-angled.

⁽¹⁾ whether what I designate here as a flower is really a single flower or an inflorescence, will be fully considered afterwards.

Historical account:—VAN TIEGHEM considers the female flower as a real single flower representing a carpellary leaf which is the first leaf of an undeveloped axillary bud; and the so-called flower-stalk as the leaf-stalk ending with two ovules, each of which corresponds to a half of the lamina of the carpellary leaf.

But there are often found flowers with more than two ovules, of which he remarks that though the leaves of *Ginkgo* are usually 2-lobed, 3—6-lobed ones are occasionally met with, and carpels with so many ovules may naturally occur.⁽¹⁾

STRASBURGER denies this view, and considers the female flower as a 2-flowered inflorescence, the ovule as the terminal formation of the secondary axillary shoot of the axis of inflorescence, where the subtending bract is abortive, and the cup-shaped basal swelling as either an arillus or a kind of pseudo-arillus formed by the swelling of the axis around the ovule.⁽²⁾ And he remarks, on comparing *Coniferae* with *Cycadaceae*, that in the former ovules are born on the "Achsenorgan," while in the latter on the "Blattorgan."⁽³⁾

According to EICHLER's latest view on the morphology in question the female flower is a real single flower, but the flower-stalk is a small shoot, and the cup-shaped swelling at the base of the ovule is the rudimentary carpel.⁽⁴⁾

SOLMS-LAUBACH also denies VAN TIEGHEM's view, and describes the the female flower as a female flowering shoot.⁽⁵⁾ SACHS,⁽⁶⁾ GÖEBEL,⁽⁷⁾ and also most other eminent authors regard it as a shoot; but it is not by every author that the question whether the female flower is to be regarded as an inflorescence or as a single flower is discussed.

SACHS states: "The single flower [*of Ginkgo*] consists of a stalk-like elongated axis which bears immediately beneath its apex two or more rarely three lateral ovules. Neither in this genus nor in *Taxus* are there any foliar structures close to the ovules, which either from

(1) Strasburger, *Die Coniferen u. d. Gnetaceen*, p. 16 (1872).

(2) His later view—*Die Angiospermen u. d. Gymnospermen*, p. 73-76. (1879).

(3) *Die Angiospermen u. d. Gymnospermen*, p. 134.

(4) Engler u. Prantl, *Pflanzenfamilien*, Embryophyta Siphonogama, p. 109 (1889).

(5) *Fossil Botany* (English translation), p. 61 (1891).

(6) *Text Book of Botany*, Second Edition (English translation), p. 516 (1882).

(7) *Outlines of Classification and Special Morphology of Plants* (English edition), p. 326 (1887).

their position or from any other circumstance can be regarded as carpels.”⁽¹⁾ He states, also, in the classification of ovules with respect to their position, that the lateral ovules may be regarded as the equivalent of whole leaves.⁽²⁾

BRAUN has also observed no carpels in *Taxaceæ*. GEBEL seems to be of opinion that the ovule of *Ginkgo* is an axial structure, the direct terminal formation of floral axis itself,⁽³⁾ and not of its branch as considered by STRASBURGER.

MASTERS regards the cup-shaped swelling at the base of the ovule as an imperfectly developed aril.⁽⁴⁾

I may state here in passing that ASA GREY's explanation of his figure of *Ginkgo* seed in his "Structural Botany" is not true. For he states: "A drupaceous seed of the same [*G. biloba*] in vertical section, exhibiting the mature disk which forms flesh."⁽⁵⁾ By this explanation we are led to suppose that the cup-shaped swelling at the base of the ovule becomes afterwards the pulpy portion of the seed.

As to the seed of *Ginkgo*, STRASBURGER states that since no fibrovascular bundles are developed, in the integument they cannot be regarded as the cause of the formation of the prominent edges of the stony part, though they may be regarded so in the case of *Taxus*; and since in the prominent edges there have never been detected even special cell rows, he considers the edges as a formation derived by inheritance from the seed which had no fibrovascular bundles in the integument.⁽⁶⁾

(To be continued.)

(1) Text Book of Botany, p. 516 (1882).

(2) l. c. p. 575.

(3) Outlines of Classification and Special Morphology, p. 385 (1887).

(4) Journal of the Linnean Society, Botany, Vol. 27, p. 300 (1891).

(5) loc. cit. p. 271 (1880).

(6) Die Coniferen und die Gnetaceen, p. 15 (1872).