K. Okamura del.

Fig. 1, *Microcladia borealis* Rupr., Fig. 2-7, *M. glandulosa* Grev.?,
 Fig. 8-13, *M. corallinae* (Mart.) Okam., Fig. 14-17, *Carpoblepharis*
Schmitzianum (Rbd.) Okam., Fig. 18-20, *C. flaccida* Kütz.

Loranthaceæ.

Loranthus Kræmpferi, *Maxim.* Monte Kiyosumi.

Abieti firmæ *S. et Z.* ac *Tsugæ diversifoliæ*, *Max.* innascitur.

Viscum japonicum, *Thunb.* Monte Kiyosumi.

Thesæ japonicæ *Nois.* ac *Euryæ ochraceæ* *Szys.* innascitur.

(*To be continued.*)

On *Microcladia* and *Carpoblepharis*.

By

K. Okamura *Rigaku-hakushi.*

With Pl. I.

Microcladia glandulosa Grev. (?).

Among those algæ, which are rather common and abundant along the pacific coast, there is a species of *Microcladia*, which appears to me after examination to be identical with *Microcladia glandulosa* Grev. This species, however, is not recorded, as far as I know, to occur in this country, and as I have not yet seen any specimen of it, I have no small difficulty in determining the plant in question. And so I think it better to give here the description of our plant, and as the consequence of my studying the minute structure of its frond, I would give some thought on the systematic position of *Microcladia* and *Carpoblepharis*.

Description: Fronds are linear, ancipito-compressed, membranaceous, and tufted, being erect or ascending from a shorter or longer decumbent base, where the plant adheres to other algæ by emitting monosiphonous, elongated and scutate root-fibres from the under-surface. When the plant is yet young, almost all of its under-surface adhere on the substratum; but when old, the greater part is set free excepting a few basal portions. In full-grown fronds there is a more or less traceable stem or rachis which branches dichotomo-alternately above in a distichous manner. Lower branches are longer than the upper with patent axils, and the plant is somewhat corymbose in general ramification. Ramuli are erect or erecto-patent, alternately arising along both sides of branches, and are many times dichotomous, sometimes assuming a flabellate appearance. Ramelli alternate toward the apex or bifid with both arms slightly incurved when young, but becoming usually patent when old.

Tetraspores are produced in ultimate and penultimate segments of lateral ramuli; they are arranged in a longitudinal row along the external side or along both sides in two rows (in the latter case they are transverse in order). This regular arrangement is sometimes deranged by the formation of one or more tetraspores in the lower transverse rows. Tetraspores are roundish and cruciate or irregularly triparted. Cystocarps are formed on the apex or at the side of ramuli, sessil, and are provided with simple or forked, flat involucre, embracing numerous neuclei.

With respect to the structure of the frond in general, I am quite at loss to see that there is no record about its minute details, as far as it is known to me. In the diagnosis of *Microcladia* given in Engler u. Prantl's *Planzenfamilien*, for example, no minute description is given about the structure of its frond, only giving that the shoot is thoroughly corticated by a broad rind which is internally constructed of larger and externally of smaller cells. Also, Mr. Reinbold, in discussing about *Gloiothamnion*, does not give the details about the structure of *Microcladia*, only expressing that *Gloiothamnion* does not present "eine scharfe Scheidung in zwei deutlich gesonderte Schichten, wie das bei *Microcladia* der Fall,....." From these lines, it seems to me that about the structure of *Microcladia* it is only known that the cortex consists internally of larger and externally of smaller cell.

In cutting a cross-section, two or three cells larger than the remaining are disposed in a transverse row between the axis and margins. Around the central axis there are six cells which form the bases of branches verticillately arising from the axis. In a longitudinal section cut parallel to the surface, a row consisting of a few cells ascends obliquely from both sides of the axial cell, at the height half way up. Cells in that row are gradually smaller upwards, and the ultimate ones form the cortical cellules; the remaining cells cut off minor ones towards both surfaces, and by further divisions they form cortical cells for the respective positions. This mode of construction of the cortex will be best seen in Figs. 11 and 12 of our plate. For the sake of comparison, I studied the structure of frond of *Microcladia borealis* Rupr. and found it to be similar to the things above stated.

Remarks: The plant, of which I have so far described above seems to me to be identical with *Microcladia glandulosa* Grev. It appears more so from Harvey's description and illustrations given in Phyc. Brit. tab. 29. He remarks that "*M. glandulosa* is often found tangled with other algæ, upon which it grows; and sometimes as Mr. Griffiths observes, creeps over

them in the manner of a *Cuscuta*, throwing out root-like fibres along the branches," and that tetraspores are "generally disposed in longitudinal series on the outer edge of the branchlet." Again Turner (Fuci, Vol. I. p. 82.) describes that tetraspores are "arranged sometimes in a two-fold series, but more commonly without any order" and that frond puts "forth through its whole length extremely minute root-like fibres from its margin, and consequently creeping." These remarks well coincide with the characters of our plant and from these reasons. I refer provisionally the plant in question to *M. glandulosa* Grev.

Herpochondria corallinæ Falkenb.

In 1866, von Martens⁽¹⁾ described a new species of *Rhizophyllis* growing on the fronds of some corallina which were collected at Yokohama and called it *R. corallinæ*. Recently Mr. Falkenberg⁽²⁾ referred this alga to the family *Rhodomelaceæ*, establishing it in a new genus *Herpochondria*. His materials, however, were destitute of cystocarps, and consequently, it may be doubted, he was led astray from placing it in a right systematic position. I have been fortunate enough to detect ripened cystocarps of this plant, and on the examination of the same, I have found that the cystocarp has the structure which is characteristic of *Ceramiceæ*. *Thus, this plant is ought not to be placed in Rhodomelaceæ but should be put in Ceramiceæ.* The plant now may be described as follows:

Fronds are small, lying flat and adhering to substratum by bundles of monosiphonous root-fibres which are emitted from the under surface. They have thick, fleshy, cartilaginous consistence when old, and are ancipitolinear, with dichotomo-alternate branches. Branches are more or less regularly lobed alternately along both sides, or dilated in a round and flabellate expansion in free extremities. Lateral shoots grow together in their lower portion with main shoots, so that the flat frond presents a branched midrib in surface view.

The apical cells of the main and lateral shoots are described by Mr. Falkenberg to be placed in a small notch at the apex of growing extremity, and on this ground he placed *Herpochondria* in the vicinity of *Chondria*. This description, however, is not correct. The apex of shoots is round and emarginate, and in or near the bottom of the notch there are two apical cells on both sides. Cortical cells there are converging and inwardly curving,

(1) von Martens die Preus. Exped. Tange p. 119, Taf. VIII., Fig. 1. (1866)—De Toni Phyc. Jap. nov. p. 40, (1895).

(2) Engl. u. Prantl, Pflanzenfam. Alg. p. 435, Fig. 244 E.

and two-to-three-celled prominences are seen along the growing margin. In stichidia, i.e. specially transformed branchlets bearing tetraspores, only one terminal cell has been observed in a shallow terminal notch, with similarly incurved arrangement of cortical cells. This is due to the special growth (into stichidia) of dichotomous branches developed from originally dichotomous, terminal cells of a short

Tetraspores are produced either in stichidia-like branches (having one terminal cell) developed from stunted shoots or in terminal portion of not specialized shoots (having two apical cells). They are at first regularly arranged in both transverse and longitudinal rows, presenting three or four in surface view, and are formed in a verticillate manner around the central axes, becoming more numerous and irregular afterwards. Sporiferous branches are often aggregated and are sometimes proliferated from the under-surface of frond.

Cystocarps are similarly produced and placed along the margin of frond, often heaping together, and are provided with some nine involucres. Involucres are compressed and either naked or provided with 2-3 celled teeth-like processes. Neuclei are few or numerous.

In cutting a cross-section of frond, a transverse row of cells having a wider diameter is seen stretched between the central axis and margins. For the sake of convenience, I call this row the middle layer. Outside of this is occupied by similar but smaller cells which become gradually smaller outward and pass to the cortical cells. Surrounding the axial cell there are six cells, which were taken by Mr. Falkenberg for pericentral cells, having been numbered by him in five. They are not, however, those comparable to pericentral cells of plants of *Rhodomelaceae*. A cross-section made from the adpressed portion of frond shows unequal thickness of cortex above and beneath the middle layer; but a section cut from the free portion shows radial structure, so that the plant does not possess a dorso-ventral organisation, as Mr. Falkenberg considered. On cutting a longitudinal section parallel to the flat surface, the middle layer is found to be composed of a chain of larger and roundish cells, arising obliquely and oppositely from the central cell. These chains lie in parallel to each other, keeping regularity and firmly coalesce together. Cells in the chain are gradually smaller toward the margin and they branch off minor ones towards both surfaces which give rise to cortex. A section cut through the plane perpendicular to the surface presents a similar appearance differing only in the shorter length of the chain of cells arising from the axial cell. They cut off minor cells towards surface and so form cortex.

Remarks: Now, by putting *Herpochondria* in *Ceramiaceæ*, the existence of that genus becomes doubtful and I am inclined to bring it under *Microcladia* on the ground that the structure of frond, and characters of apical cells, tetraspores, as well as cystocarps are all identical to those of the latter genus. The structure above stated will be found to be as just same as that described under *Microcladia glandulosa* (?). The slightly incurved growing apex of shoots terminated by two apical cells is also same as it is in the plant just mentioned, in which the terminal cells are also two and are either incurved or parted. The mode of ramification which is dichotomo-alternate and tetraspores which are arranged at the beginning in a subregular order, are also similar. and as so many characters are common to *Microcladia*, I think it is proper to refer this plant to that genus.

Gloiothamnion Schmitzianum Reinbold.

In doing these studies, I have consulted with the conspectus of genera of *Ceramiaceæ* given in 'Pflanzenfamilien' and have known that *Carpoblepharis* is separated from subfamily *Ceramiaceæ* chiefly by having verticillately arranged cortical filaments. Fortunately, I have a few specimens of *Carpoblepharis flaccida* Kütz. in my herbarium and had chance to study the structure of its frond. In cutting a cross-section, there is observed a central axis and, of cells of the intermediate layer, those disposed along the longer axis of the section are larger than the rest; and surrounding the central axis there are six cells, of which the two placed towards each surface are a little smaller than the cell on each side. In studying the longitudinal sections, there is found six verticillate branches arising from the axis, of which the two standing on the plane parallel to the surface of frond arise from the middle portion of the cell of central axis, while the remaining two, from a place a little above. For the sake of convenience of description, I shall call the former the lower branches and the latter, the upper. Cells forming the lower branches are slightly larger than those of the upper, and these larger ones are those stretched between margins, as is seen in a cross-section. Two proximal cells of the lower branch run parallel to the axis and the next cells succeeding above are turned toward margins. All those cells cut off minor ones toward periphery and thus cortical cells for that portion are formed. Again, the three proximal cells of the upper branch run parallel to the axis and come in contact with those of the series next both above and below. Each of these cuts off cells towards periphery, which give rise by further divisions into cortical cells for that portion. All the cells are firmly coalesced together and

consequently a cross-section of frond presents a parenchymatic appearance.

The state of things similar to that described above is seen in the structure of *Gloiothamnion Schmitzianum* Rbd. This genus was established by Mr. Reinbold⁽¹⁾ in 1895, who distinguished that genus by "die besondere Structur des Thallus, speciell der Rinde, welche eine scharfe Scheidung in zwei deutlich gesonderte Schichten, wie das bei *Microcladia* der Fall, nicht erkennen lässt." In his illustration there is no sharp distinction of cortical and intra-cortical layers. But according to my study of the same plant, the cortical layer is separated from the central axis by a few cells of an intra-cortical layer. In a cross-section there are six cells surrounding the central axis. In cutting a longitudinal section from a thicker portion of frond, we find 2-3 celled branches arising verticillately and obliquely from the axis. From the cells of those branches, cortical cells are derived directly or indirectly. These branches arise from the central axis at the same level with no marked difference in size of cells, as it is the case in *Carpobelepharis flaccida*. Judging from my study, I came to the conclusion that Mr. Reinbold's expression above quoted and his illustrations given, would have perhaps been done from the study either of a younger or of a narrower frond; because, the section of a very narrow portion of frond really presents an appearance similar to that illustrated in his figures 3 and 4. Even in such sections (made from a branch having the thickness of 130 μ) a few intra-cortical cells can be made out, only very much smaller and more closely placed beneath or almost among cortical cells.

Moreover, *Gloiothamnion Schmitzianum* much varies in the thickness of frond, habit of branches and robustness of ramification, as Mr. Reinbold has remarked for those growing on *Chondrus*. He has illustrated the narrowest forms of frond as the type of this species. Plants having such forms are slenderer and subcylindrical (attaining the diameter of about 130-180 μ), and have a less robust ramification. Even in those, the upper branches, either sterile or fructiferous, are more or less compressed. A very differently appearing plants have been collected by me at Bōshū, which have broader (375-500 μ in the broadest part) and ancipito-compressed fronds branching in a close pinnate. They seemed to me at first to be a species different from *Gloiothamnion Schmitzianum* Rbd. But, the basal portion of those plants are subterete and they grow in the same place together with the typical forms. In such extraordinary forms, the structure of frond like that described above is quite manifest.

(1) Reinbold in Hedwigia XXXIV, Taf. III. (1895).—Pflanzenfam. Alg. p. 502.—Reinboldiella Schmitziana (Reinb.) De Toni Phyc. Jap. Nov. p. 35. (1895).

Such is the result of my examination, which seems unfortunately quite contradicted with that of Mr. Reinbold. And if I am not mistaken, *I want to say that establishment of the new genus on the ground chosen by him is not preferable.* And here a question arises whether this plant should be kept as a distinct genus as stated by him or be put into some of the genera already established. To me, this plant does not seem to be referable to *Microcladia* as the author had at first inclined to do. The pinnate mode of ramification, non-curved growing apex of shoots, and slightly expanded stichidial ramuli are quite different from the last genus. On the contrary, *I am rather inclined to place it in Carpolepharis* by the structure of frond, straight apex of shoot, etc. Mr. Reinbold already remarked that *Gloiothamnion* will perhaps take its systematic position between *Microcladia* and *Carpoblepharis*, with the latter of which it is related by the characters being common with *Carpoblepharis*, as it seems to me, only the slenderness and minuteness of the frond, and its decumbent and root-bearing habit differ from that genus. Of its size, however, some have, as I have described above, very much compressed and broadened pinnate fronds, which remind some of *Carpoblepharis*; and then the decumbent habit only will not be sufficient enough for establishing a plant in a new genus.

Cystocarps are produced from stunted ramuli along branches and a single nucleus is surrounded by a few-compressed involucre.

From the study of *Microcladia*, *Herpochondria*, *Gloiothamnion* and *Carpoblepharis* chiefly done with respect to the structure of frond, I have a view about the systematic position of *Microcladia* and *Carpoblepharis*. By consulting with 'Pflanzenfamilien' the subfamily *Carpoblepharideae* seems to be separated from *Ceramieae* chiefly by having verticillate cortical filaments, if not by minute differences in the formation of cystocarps. If my consideration is to be justified, *Microcladia* and *Carpoblepharis* (as *Herpochondria* and *Gloiothamnion* have been reduced to the respective genera) ought to be placed in one and the same subfamily; or *Carpoblepharis* might be classed in *Ceramieae* together with *Microcladia*, *Ceramium*, etc., as Mr. Reinbold seems to be inclined to do. But, as the formation of cortex in *Ceramium* and its allies somewhat differs from that of *Carpoblepharis* and *Microcladia*, it will be proper to class the both genera in a subfamily, separating them from *Ceramieae* which will then contain *Ceramium*, *Campylæphora*, etc.

Nov. 24, 1899.

No. 6, Hinoki-cho, Akasaka,
Tokyo, Japan.

EXPLANATION OF FIGURES IN PLATE I.

Microcladia borealis Rupr.

Fig. 1. Diagram showing longitudinal section of the frond; a , the central axis.

Microcladia glandulosa Grev. (?)

Fig. 2. Frond bearing tetraspores in nat. size.

Fig. 3. Cystocarps. $\frac{1}{1}^2$.

Fig. 4. Pinnellæ bearing tetraspores; moderately magnified.

Fig. 5. Cross section of the frond; a , the central axis; moderately magd.

Fig. 6. Longitudinal section cut parallel to the surface of frond. $\frac{7}{1}^3$.

Fig. 7. " " " perpendicular " " $\frac{7}{1}^3$.

Microcladia corallinæ (Mart.) Okam.

Fig. 8. A piece of the sporiferous frond. $\frac{1}{1}^2$.

Fig. 9. The apical portion, a , of Fig. 8 more highly magnified; a , the apical cells. $\frac{2}{1}^0$.

Fig. 10. Longitudinal section cut perpendicular to the surface of frond; $\frac{2}{1}^0$.

Fig. 11. Cross section of a free portion of the frond; a , the central axis; $\frac{7}{1}^3$.

Fig. 12. Cystocarps; $\frac{1}{1}^5$.

Fig. 13. Cross-section of stichidial ramuli; $\frac{4}{1}^1$.

Gloiothamnion Schmitzianum (Rbd.) Okam.

Fig. 14. Longitudinal section perpendicular to the surface of frond; $\frac{3}{1}^0$.

Fig. 15. Cross-section of the frond; $\frac{3}{1}^0$.

Fig. 16. Two sterile fronds of the extraordinary broader form, $\frac{3}{1}^1$.

Fig. 17. Cross section of frond cut from the same; $\frac{2}{1}^0$.

Carpoblepharis flaccida Kütz.

Fig. 18. Diagram showing the longitudinal section cut parallel to the surface of frond; two points on the upper portion of the central cells indicate the starting points of the upper branches.

Fig. 19. Diagram showing the longitudinal section cut perpendicular to the surface of frond.

Fig. 20. Half of the cross-section of frond; a , the central axis; $\frac{2}{1}^0$.