Fig. 7.

× 2.5

Fig. 8.

× 280.

PINGUICULA RAMOSA sp. nov.

(Kōshinsō.)

Notes on *Pinguicula ramosa*, sp. nov.

(with Plate XI.)

By M. Miyoshi, *Rigakushi*.

On the 9th. August, 1890, I found a species of *Pinguicula* on Mount Kōshin in the province of Shimotsuke. It grows in great numbers on the moist exposed surface of the huge rocks which constitute the rugged outline of the peak. Unfortunately, the season was too late for the flower, so the specimens I collected were all in fruit. The following are the characters of the plant except the floral parts:—

Root fibrous.

Leaves 5 or 6, petiolate, radical, clustered, reflexed, light green, fleshy. Lamina (Fig. 6) elliptical, 1—1.5 cm. long, .5—1 cm. broad; margin entire, involute; apex rounded, often cleft at the middle; surface viscid, covered with minute glands (Fig. 8). Petiole (Fig. 6),

.5—1 cm. long, lighter-coloured than the lamina.

Scape 1, naked, light green, viscid, 7—9 cm. high, 1—2 mm. in diam., usually forked (Figs. 2, 3, 4) into 2 similar branches 1—2.5 cm. above the base, one of which often forks again, forming together with the other 3 similar branches (Fig. 5).

Flower 1, terminating the scape or branches.

Calyx persistent, 5-parted, 1.5 mm. in diam., lobes ovate-lanceolate.

Fruit a capsule (Fig. 7), ovate, 3 mm. in diam., longer than the calyx, opening in 2 valves.

Seeds minute 1 mm. long, .5 mm. broad, numerous, brown, elliptical, attached to the free central placenta.

As the plant grows on the perpendicular sides of rocks, the scapes assume an ascending position as shown in the accompanying figures.

Though my specimens, being destitute of the flower, are of course incomplete for the precise determination of the

species, yet the branching habit of the scape seems to me so peculiar a character, which I have noticed neither in any one of the descriptions of 32 species of this genus in De Candolle's *Prodromus* (Vol. VIII. p. 26-32), nor in the floras of different regions which I have examined, that I have provisionally named the plant *Pinguicula ramosa*. A strict determination, however, will be reserved until the next season, when I shall have an opportunity of examining the flower.

Explanation of Plate XI.

Figs. 2, 3.—Whole plants in the natural position (nat. size).

Figs. 4, 5.—Part of branching scapes (nat. size).

Fig. 6.—Leaf showing its petiolar portion (nat. size).

Fig. 7.—Capsule opening in 2 valves, showing the apex of the free central placenta ($\times 2.5$).

Fig. 8.—Two kinds of glands on the upper surface of the leaf ($\times 230$).

- a. Larger gland with usually 16 (sometimes 14—18) radiated cells.
- b. Side view of the larger gland showing its unicellular pedicel and the point of attachment.
- c. Smaller gland with usually 8 (sometimes 6—10) radiated cells.
- d. Stomata.
- e. Cells of the epidermis.

左記ハ予が今夏、日光ニ滞在シテ植物採集中、理科大學植物學撰科松田定久氏ト庚申山ニ赴キタル時ノ事ニシテ、予が則稿ノ植物採集旅行雜記中ニ收ムベキモノナレドモ、殊ニ茲ニ則記スルトナセリ、

○*Pinguicula ramosa*, sp. nov. (第十一圖) 二

就テ 理學士 三 好 學

かうしんそう(新報) (狸藻科)

明治廿二年^{カクシヤ}九月十八日、下野國^{カクシヤ}庚申山ニ於テ一種ノ *Pinguicula* ヲ採集セリ、最初ハ僅ニ六七株ヲ同山別所^{ベツシヨ}院寺ノ上ナル富士淺間^{フジノ}案内者^{サネノ}ノ岩ニテ發見^{ハツク}シタルガ、其レヨ