Geum pentapetalum MAKINO. *Ledum palustre* L.
Andromeda polifolia L. *Oxycoccus palustris* PERS.
Oxycoccus microcarpus TURCZ. *Vaccinium Vitis-idaea* L.
Trientalis europaea L. *Gentiana nipponica* MAKINO.

Botanical Institute, Faculty of Agriculture,
Hokkaido Imperial University, Sapporo,
(VI. 14, 1928.)

Explanation of the figures.

1. A beach-association near Onsenzaki, taken on IX. 8, 1927.
2. A grassy land-association near Tokotan, taken on IX. 12, 1927.
3. A heath-association near Tokotan, taken on IX. 9, 1927.
4. A swamp-association (a consociation of *Carex Lyngbyei*) near Hayakawa, taken on IX. 9, 1927.

On the number of chromosomes in *Diospyros Kaki* L. f. and *Diospyros Lotus* L.

By

Isawo Namikawa and Motoki Higashi.

As for *Diospyros Kaki*, YASUI¹⁾ mentioned that "the number of chromosomes is difficult to count by reason of their small size" and "usually there are 28 in pairs, but sometimes 27". The writers are attempting to look through the chromosome number in different cultural varieties of *Diospyros Kaki* and in *D. Lotus* which is widely used for the stock of Kaki. The result obtained does not agree with that of the previous investigator and seems worthy noting here.

The number of chromosomes was counted with pollen mother cells and root tips of the seedlings. The material was taken from the experimental orchard of the College of Agriculture, Kyoto Imperial University. The Kaki varieties studied here are non-astringent varieties. With *Diospyros Lotus*, different individuals from different Kaki-growing localities in Japan were examined.

The fixatives giving satisfactory results for the respective materials are as follows :

1) YASUI, K.—Studies of *Diospyros Kaki*, I. Bot. Gaz. 60: 362-373, 1915.

<i>Diospyros Kaki</i>		<i>Diospyros Lotus</i>	
Pollen mother cells	Root tips	Pollen mother cells	Root tips
BRASIL'S modification of BOUIN'S fluid and CHAMPY'S fluid.	The Bonn modification of FLEMING'S fluid.	BRASIL'S modification of BOUIN'S fluid.	BRASIL'S modification of BOUIN'S fluid.

FLEMMING'S solutions and chromo-acetic solutions of varying concentrations and CARNOY'S solution were also tried for the fixing of the pollen mother cells of these two species. The fixing of the prophase up to the diakinesis was secured fairly well, but good metaphase figures could not be obtained. The root tips of *Diospyros Lotus* fixed with those fixatives stained very poorly. BRASIL'S doris proved the most satisfactory in result for the pollen mother cells of the two species and for the root tips of *Diospyros Lotus*. MINOUCHI²⁾ recommends CHAMPY'S fluid for the fixing of mammalian chromosomes. This fixing agent gave good results for the stages later than the diakinesis of *Diospyros Kaki*. In the metaphase figure fixed with this solution, the chromosomes and even the nuclear plate itself (Fig. 6) show remarkably depressed size as compared with those fixed in other fixatives (Fig. 5), but the each pair of gemini is very clearly discernible. The prophase nuclei fixed in this mixture exhibit more or less of disintegration. This fixative can not be appropriated for *Diospyros Lotus*. The root tips of *Diospyros Kaki* are easily fixed in FLEMMING'S fluid.

Before the fixing of the pollen mother cells, preliminary observations were made with the boiling method with aceto-alcohol and aceto-carmines devised by HERTZ.³⁾ The approximate number of the chromosomes can be counted by this method.⁴⁾ HEIDENHAIN'S iron haematoxylin was applied for staining. Reliable countings were made in the metaphase and the anaphase. A few countings were also successfully tried in the diakinesis.

The number of chromosomes determined is as follows:

2) MINOUCHI, O.—On the fixation of chromosomes in mammals and other animals. Jap. Journ. Zool. 1: 219-234, 1928.

3) HERTZ, Z.—Der Nachweis der Chromosomen. Zeitschr. f. Bot. 18: 625-681, 1926.

4) Mr. SASAOKA, one of the research workers in the Imperial Horticultural Experiment Station, Okitsu, wrote recently to one of the writers that he examined the pollen mother cells of Zenji with the HERTZ method and counted approximately 45 pairs of gemini. The result agrees with the writers' observation.

	n	2n
<i>Diospyros Lotus</i>	15	30
<i>Diospyros Kaki</i> E-Gosho	45	—
Kurokuma	45	—
Nara-Gosho	45	—
Shôgatsu	45	—
Zenji	45	—
Seedlings of Anzai	—	90
Seedlings of Kubo	45	90
Seedlings of Tenjin-Gosho	—	90

In the case of *Diospyros Lotus*, which is dioecious, the haploid count was made in the metaphase and the anaphase of the heterotype as well as of the homotype division, and the result obtained in these different stages was the same. In *Kaki* varieties examined, no variation in the chromosome number was observed. The possibility of certain variations in the number of chromosomes in different cultural varieties might be expected if the present result is compared with that of Miss YASUI'S. A further survey will be interesting.

Fig. 1.

Fig. 2.

Fig. 3.

Fig. 4.

Fig. 5.

Fig. 6.

Fig. 1. Metaphase figure of somatic nuclear division of *Diospyros Lotus*. Fig. 2. Metaphase figure of heterotype division of *D. Lotus*. Fig. 3. Anaphase of the homotype division in *D. Lotus*. Fig. 4. Somatic metaphase of a seedling of Anzai (*D. Kaki*) fixed with FLEMING'S solution. Fig. 5. Anaphase of heterotype division of Shôgatsu (*D. Kaki*) fixed with BRASIL'S modification of BOVIN'S fluid. Fig. 6. Metaphase of heterotype division of E-Gosho (*D. Kaki*) fixed in CHAMPY'S mixture. Ca. 2300/1.