WIKSTROEMIA ALBIFLORA, SP. NOV.
NOM. JAP. H/O. ヒオウ.

植物學雜誌第五卷第五十三號

明治廿四年七月十日

THE BOTANICAL MAGAZINE.

[Vol. V.]

July 10, 1891.

No. 53.]

A New Japanese Wikstroemia.

Ryōkichi Yatabe, Sc. D.

Wikstroemia albiflora, nov. sp.

Nom. jap. *Hiō*. ヒオウ.

(Plate XXVI.)

A shrub 2—5 feet high; young branches brown, old branches grey. Leaves opposite, glabrous, ovate or broadly ovate-lanceolate, entire, pale or glaucescent and with 3—5 rather prominent pinnate veins on the under surface, $\frac{1}{2}$ —1 inch long, 3—6 lines broad; petioles short, $\frac{1}{2}$ line long. Flowers white, subsessile, 3 or 4 together, capitate, terminating the branches. Perianth-tube smooth, slender, slightly dilated towards the apex, 3 lines long; its limb 4-parted; the lobes patent, broadly ovate, obtuse, about 1 line long, imbricate in the bud. Stamens 8 in 2 series, those opposite the lobes of perianth inserted in the throat, the alternate ones in the middle portion of the tube; filaments short; anthers oblong, obtuse, adnate, orange in color. Hypogynous scale small, 1 or wanting. Ovary clavate, with the base stalk-like, slightly curved, 1-celled, about 1 line long; style very short; stigma capitate, hispidulous under the lens. Ovules solitary, anatropous, pendulous.—Fl. May.

Living specimens of this plant were collected by Mr. K. Watanabe, at Nanokawa-mura, in the province of Tosa, in March 1891, and are now kept in the Botanic Garden of the Imperial University. Its near relative is *W.*

japonica, Miq. The latter can be readily distinguished by its inflorescence, which, though short and contracted, is racemose and not truly capitate. Moreover it is not only terminal, but also axillary, which latter is not the case with the present species.

Plate XXVI. Fig. 1, branch with flowers; 2, flower cut vertically; 3, ventral and dorsal views of stamen; 4, pistil with a scale at the base; 5, ovule; 6, floral diagram:—*all except 1 enlarged.*

日本藥局方植物篇

(前號ノ續キ)

會員 澤田駒次郎

第五十二號正誤(189)ノ第十七行十二字目餘ノ下幅ノ字ヲ脱ス同第十九行二十一字目幅ノ下ニ在ルサノ字ハ不用(190)ノ第十三行三十二字目法ハ方ノ誤(191)ノ第九行十一字目粉濃ハ澱粉ノ誤ノ(192)第二行二字目ノ交糾ノ上「マヂレル」トアルハ「ヨヂレル」ノ誤全第十五行二十九字目産ノ下スノ字ヲ脱ス全十八行二十字目治明ハ明治ノ誤(193)ノ第二行二十一字目色ハ褐ノ誤

成分 加刺拔兒豆ノ主成分ハ有毒性亞爾加魯乙度ニシテ一千八百六十四年ニ於テ Jobst, Hesse ノ兩氏之ヲ發見シテ *Physostigma* ト命名セリ爾後一千八百六十七年ニ至リ Hesse 氏ハ原素分析ニ因リ其集成ヲ確定シ $C_{15}H_{21}N_3O_2$ ヨリナルモノトセリ

Physostigma ハ一名 *Physostigmin* ト稱シ褐色不品形ノ塊ニシテ設氏四十度(華氏百四度)ノ熱度ニ因テ軟化シ設氏四十五度(華氏百十三度)ノ熱度ニ逢フテ熔融ス酒精, 依的兒, 扁陣, 硫化炭素, 噶囉仿謨ニハ隨意ニ溶解シ水ニハ微ニ溶解ス又酸類ヲ中性ト爲シ鹽化鐵ノ溶液ヨリ含水酸化鐵ヲ沈澱ス

本品及ビ其他ノ鹽類ハ味ヲ有セス此等ノ溶液ニ Meyer 氏ノ試藥ヲ注