Orchidaceae Novae Micronesiacaе, III.^{1,2)}

Auctore

Takasi Tuyama

Received June 10, 1940.

25) *Moerenhoutia Hosokawae* (FUKUYAMA) TUYAMA stat. nov.

Moerenhoutia leucantha SCHLECHTER var. *Hosokawae* FUKUYAMA in Trans. Nat. Hist. Soc. Formos. 29-188, p. 97 (1939).

Ut FUKUYAMA indicat, toto ad *M. leucantham* Ponapensem valde affinis. Terrestri adscendens, usque ad 32 cm alta, rhizomate decumbente cauliformi, radicibus elongatis villosis, caule stricto basi folio donato glabro tereti, supra basin vaginis paucis amplexantibus obsesso, inflorescentiam versus paululo glanduloso-puberulo, foliis ca. 6 subremotis subrosulatis dispositis erecto-patentibus petiolatis, lamina elliptica utrinque acuminata glabra cum petiolis 3 cm longis 13-17 cm longa 3.2-4.1 cm lata, petiolo basi dilatato caulem vaginante glabro, spica erecta subdensa ca. 20-flora usque ad 20 cm longa, bracteis erecto-patentibus ovato-lanceolatis acuminatis ovario aequalibus vel brevioribus, floribus erecto-patentibus vel suberectis albis (fide TAKAMATSU), sepalis concavis basi exteriori tantum puberulis 10.5 mm longis 3 mm latis in explanato lineari-ellipticis basi latissimis valide 1-nerviatis, sepalis lateralibus cum sepalo intermedio conformibus sed apice subito acutis margine anteriore leviter dilatatis et 4 mm latis, petalis late falcatis unguis lineari-angustatis 1.5 mm latis supra medium margine anteriore subito ample dilatatis et 3 mm latis toto 11 mm longis, labello ca. 10 vel 9.5 mm longo 3 mm lato in explanato 5 mm lato e basi vesiculoso-cucullato, in tertia parte basilari subito valde constricto et iterum in laminam cucullato-orbiculatam productum, in tertia parte apicali leviter constricto et in lobum apicalem ovatum obtusum 3 mm longum et latum leviter composito-involutum productum, basi utrinque cornibus minoribus numerosis dispersis non fasciculatis, lamellis 2 e basi usque ad tertiam partem apicalem decurrentibus superiore latioribus et leviter divergentibus, carina superiore inter lamella incrassato-elevata, ovario cylindrico-fusiformi ca. 12 mm longo 2.5 mm lato pubescente, pedicello 1-2 mm longo, columna

1) Contributions from the Laboratory of Systematic Botany (Prof. T. NAKAI) of the Botanical Institute, Faculty of Science, Tokyo Imperial University.

2) The expense of the study has been partly defrayed from the subsidy granted by the Japan Society for the Promotion of Scientific Research.

gracili cum rostellis 7 mm longa antice utrinque alata anteriore superne orbiculato-dilatata et planata, rostellis gracillimis 2 donatis ca. 2.5 mm longis.

Nom. Jap. *Kusai-kukuriran* (nov.).

Spec. exam. Caroline, Ins. Kusai, Matante (leg. R. KANEHIRA, field. no. 4230, Mai. 18, 1937); ibidem, in monte Fenkol (leg. M. TAKAMATSU, no. 377, Jan. 28, 1936, steril. in Herb. B. P. Bishop Mus.).

Distr. In Insula Kusai endemica.

Specimen typicum non vidi, sed haec species cum labello sub apicem constricto, carina humilliore, cornibus minoribus in basi labellorum utrinque sparsa, petalis supra subito late falcatis dilatatis basi anguste unguiculatis ex *M. leucantha* distinguetur.

26) **Moerenhoutia leucantha** SCHLECHTER in ENGLER, Bot. Jahrb. **56**, p. 450 (1921).

var. **minor** TUYAMA var. nov.

Differt a forma typica, folio floribusque-minoribus, foliis 5.5–8.5 cm longis 2.0–2.3 cm latis, sepalis petalisque 8–9 mm longis.

Nom. Jap. *Hime-ponape-kukuriran* (nov.).

Spec. exam. Caroline, Ins. Ponape, in oppido Yû (leg. Y. KOBAYASI, Aug. 30, 1936—Typus in Herb. Univ. Imp. Tokyo.).

Distr. In Insula Ponape endemica.

27) **Nervilia Aragoana** GAUDICHARD in Freycinet, Voy. Bot. p. 422 (1826).

Pogonia flabelliformis LINDLEY in WALLICH, Cat. n. 7400 (1832); et Gen. Sp. Orch. p. 415 (1840).

Pogonia Nervilia BLUME, Mus. Bot. Lugd. Bat. **1**, p. 32 (1849).

Pogonia gracilis BLUME, Orch. Arch. Ind. p. 129 t. 57 (1858).

Nervilia yaeyamensis HAYATA, Icon. Pl. Formos. **2**, p. 140 (1912); et **4**, p. 118 (1914); **5**, p. 213 (1915), **syn. nov.**

Nom. Jap. *Awoi-bokuro*, *Yaeyama-hitotubaran* (NEMOTO, 1936).

Spec. exam. Micronesia, Caroline; Palau, Ins. Pelilin (leg. S. HATU-SIMA, field no. 4791 Apr. 7, 1938, floribunda).

Liukiu, Ins. Iriomote (leg. KANASHIRO, no. 86, Jul. 10, 1907); ibidem, Ins. Isigaki (leg.?, Jul. 7, 1911).

Distr. Japonia austr., Formosa, Philippine, Timor, Nova Guinea, Samoa.

Nova additio ad floram Palauensem.

28) **Nervilia palawensis** SCHLECHTER in ENGLER, Bot. Jahrb. **56**, p. 447 (1920).

Nervillia oxyglossa FUKUYAMA in Trans. Nat. Hist. Soc. Formos. 27, p. 278 (1937); **syn. nov.**

Nom. Jap. *Parao-hitotubokuro* (nov.).

Spec. exam. Caroline, Palau; Ins. Baobeldaob, Ngarmiskan (leg. T. TUYAMA, Sept. 1, 1937, fructifera); ibidem, in monte Ngarsul (leg. T. TUYAMA, Sept. 13 et 14, 1937, steril); ibidem, Neo-Ngathpan (leg. T. TUYAMA, Sept. 5, 1939, frutifera et florifera).

Distr. Endemica.

Ex auctoritate FUKUYAMAE, *N. oxyglossa* cum *N. palawensi* eadem est, cum exclusio magnitudinis corollae. *Nervillia palawensis* corollam post anthesin vulgo valde ampliata habet, id est, corolla primum 1–1.2 cm longa demum 3 cm vel ultra longa.

29) **Oberonia ponapensis** TUYAMA sp. nov. (sect. *Otoglossum*).

Epiphytica erecta vel pendula pluricaulis, 6–13 cm alta, radicibus filiformibus laxè undulatis elongatis glabris, caulibus leviter flexuosis 6–7-foliatis, basi vaginis paucis brevibus donatis, foliis suberectis linearibus apice falcatis incurvatis attenuatis 3–5.5 cm longis medio fere 2.5–4 mm latis basi vaginantibus, vaginis caulem omnino obtegentibus anceptibus marginibus hyalino-membranaceis anterioribus altius connatis. Inflorescentia erecta vel patula pro genere laxiuscula usque ad 6.5 cm longa 4–4.5 mm lata, bracteis lanceolatis longe acuminatis ca. 2.5 mm longis ovario manifeste longioribus, floribus in genere vix mediocribus subverticillatim dispositis aurantiaco-brunneis diametro 2 mm. Sepala ovata subacuta 0.9 mm longa 0.6 mm lata, lateralia obliqua conformia. Petala anguste elliptica 0.8 mm longa 0.3 mm lata margine irregulari undulata obtusiuscula. Labellum basi auriculatum, auriculis oblique semiovatis subacutis, lamina lineari-oblonga anteriore subito leviter dilatata antice in lobos 2 falcatos triangulos acutos producto, inter lobos latissime 1-dentata. Columna brevis a crassiuscula 0.5 mm crassa. Ovarium cum stipite brevi 1.5 mm longum. Capsula late obovoidea 2 mm longa 1.7 mm lata dehiscens.

Nom. Jap. *Ponape-yôrakuran* (nov.).

Spec. exam. Caroline, Ponape; in oppido Kityi, ad summum montis Troton alt. 600 m. (leg. S. HATUSIMA, no. 11102, Jul. 1939—typus in Herb. Univ. Imp. Tokyo.).

Distr. Endemica.

In sectione, cum *O. radicante*, *O. lineari*, et *O. brunneo* similis. Ex 3 his speciebus cum postremo simillima, sed bracteis ovario prominente longioribus, petalis latioribus, labello apice bicuspidato et medio multo obsolete latiusque acutis differt.

30) *Peristylus carolinensis* (SCHLECHTER) TUYAMA comb. nov.

Habenaria carolinensis SCHLECHTER in ENGLER, Bot. Jahrb. **56**, p. 445 (1920).

Nom. Jap. *Panape-sagiso* (nov.).

Spec. exam. Caroline, Ins. Ponape, Mataranium (leg. Y. KOBAYASI, Aug. 31, 1936); ibidem, in monte Nanalaut (leg. M. TAKAMATSU, no. 1077, Feb. 17, 1936, in Herb. B. P. Bishop Mus.).

Distr. In Insula Ponape endemica.

Peristylus format grex valde naturalis in genere *Habenaria* in sensu latiore, itaque combinatio nova proponatur.

31) *Peristylus palawensis* (TUYAMA) TUYAMA comb. nov.

Habenaria palawensis TUYAMA in Bot. Mag. Tokyo, **53**, p. 52 (1939).

Nom. Jap. *Parao-sagiso* (TUYAMA, 1939).

Spec. exam. Caroline, Palau; Ins. Baobeldaob, in monte Kattelwel (leg. T. TUYAMA, Sept. 3, 1937—typus in Herb. Univ. Imp. Tokyo.); ibidem, in monte Ngarsul (leg. T. TUYAMA, Sept. 13, 1937).

Distr. In Insula Baobeldaob endemica.

Cum aequali causa combinatio nova proponatur.

32) *Peristylus setifer* TUYAMA sp. nov.

Habenaria setifera TUYAMA nom. altern.

Habenaria sp. VOLKENS in ENGLER, Bot. Jahrb. **31**, p. 461 (1901); SCHLECHTER in ENGLER, Bot. Jahrb. **52**, p. 5 (1914); et in l. c. **56**, p. 445 (1921).

Terrestris erecta cum scapo 40 cm alta glaberrima, radicibus fusiformibus villosis paucis ca. 5 cm longis, tuberis albis flaccidis longe oblongis 1–2 donatis, caule ad 17 cm longo basi cum cataphyllis alte vaginatis paucis toto obtectis superioribus foliaceis, foliis ad apicem caulium approximatis horizontali patentibus 5–7 donatis, laminis lanceolatis apice longe acutatis vel leviter acuminatis basi subito angustatis et in petiolos vaginatis caulem ample amplectentibus margine integris interdum undululatis ad 15 cm longis 3 cm latis in sicco membranaceis costis infra elevatis, nervis utrinque 5 natis in vivo profunde viridibus, spica gracili subdense multiflora apice subnutanti ad 18 cm alta, pedunculo 13 cm longo vaginis adpressis paucis obsesso, bracteis foliaceis suberectis lanceolatis 1-carinatis apice subito longe acuminatis ca. 1 cm longis ovario brevioribus, floribus 0.8–1.4 cm distantibus erecto-patentibus sessilibus viridibus, sepalo intermedio convexo ovato-oblongo obtuso 2.5 mm longo in explanato 1.6 mm lato 1-nervato, lateralibus oblique oblongis obtusis 3.4 mm longis 1.4 mm latis, basi in unguis obliquis angustatis convexis dorso carinatis, carinis apice in parte 2/3 longitudinis

sepalorum liberis et setiforme productis, petalis oblique lateque ovatis 3 mm longis 2.1 mm latis valde obtusis 1-nervatis, labello fureatim 3-partito ca. ad 3.6 mm longo, lobis lateralibus 6 mm longis basi 0.5 mm latis apicem versus gradatim attenuatis et spiraliter incurvatis cum lobo intermedio in angulo 35° patentibus, lobo intermedio recte producto anguste oblongo apice acuto-obtuso, calcare ca. 1.7 mm longo basi 1 mm lato clavato supra medium levissime inflato et apicem versus angustato, apice apiculato et levissime incurvato 6 mm longo 1 mm lato ovario 2/3 breviori, antherae canalibus brevibus, processibus stigmatiferis utrinque infra 1 mm rotundato obtusis incurvatis, ovario sub anthesi ca. 1 cm longo 1.5 mm lato anguste cylindraceo erecto apice anteriore patente.

Nom. Jap. *Yappu-sagisô* (nov.).

Spec. exam. Caroline, Yap; Ins. Yap, in monte circa Tafgif alt. 150 m. (leg. T. TUYAMA, Sept. 19, 1939—typus in Herb. Univ. Imp. Tokyo.); ibidem, in monte Kaburu (Sankaku-rippyôno yama) alt. 100 m. (leg. T. TUYAMA, Sept. 17, 1939).

Distr. Endemica.

Habenaria papuana valde affinis sed foliis bracteisque angustioribus, floribus densioribus majoribus, calcaribus brevioribus, sepalis lateralibus setiferis exqua differt.

33) *Phreatia landronica* TUYAMA sp. nov. (Sect. *Euphreatia*).

Epiphytica erecta usque ad 18 cm alta, radicibus filiformibus albescentibus villosulis, caule subnullo, foliis erecto-patentibus ca. 7 donatis distichis digitatim dispositis lineari-ligulatis apice valde inaequaliter bilobatis 11–17 cm longis 1.3–1.6 cm latis coriaceo-papyraceis in sicco nervis lateralibus principalibus utrinque 3, vaginis compressis carnosus in sicco callosis 2–2.5 cm longis truncatim articulatis, inflorescentiis erecto-patentibus vaginulis paucis acuminatis ca. ad 1 cm longis donatis foliis prominente brevioribus, pedunculo 4.5–7 cm longo, racemo 4–5 cm longo densifloro, bracteis lanceolatis acuminatis fusciscentibus glabris patentissimis sub anthesi ovario aequilongis vel brevioribus, floribus in genere inter minores albis, sepalis uninervatis late obtuso-acutis, intermediis oblongo-ovatis ca. 1 mm longis, sepalis lateralibus late ovatis obliquis cum pede columnae mentum brevem obtusum formantibus, petalis ellipticis 1-nervatis sepalis brevioribus 0.8 mm longis, labello rhombeo-suborbiculari obtuso et apiculato 3-nervato margine inferioribus utrinque levissime sinuato basi contracto et brevissime unguiculato, columna perbrevis, anthera rotundata antice obtusissima, ovario sub anthesi cum pedicello 0.9 mm longo filiformi 1.5 mm longo obovoideo, capsula juvenili ca. 1.7 mm longa ellipsoidea.

Nom. Jap. *Mariana-hureran* (nov.).

Spec. exam. Marianne, Saipan; ad summum montis Tappôtyô, alt. 470 m. (leg. S. HATUSIMA, Jul. 5, 1939, no. 10673—typus in Herb. Univ. Imp. Tokyo.); ibidem (leg. T. TUYAMA, Sept. 25, 1930, steril.).

Distr. Endemica.

34) **Phreatia palawensis** (SCHLECHTER) TUYAMA comb. nov.

Rhynchophreatia palawensis SCHLECHTER in ENGLER, Bot. Jahrb. **56**, p. 488 (1921).

Phreatia Ryozoana TUYAMA in Bot. Mag. Tokyo, **53**, p. 54 (1939), syn. nov.

Nom. Jap. *Honaga-hureran* (TUYAMA, 1939).

Spec. exam. Caroline, Palau; Ins. Baobeldaob, sine loco speciali (leg. G. KOIDZUMI, Feb. 1–3, 1915); ibidem, Aimiriik (leg. R. KANEHIRA, field no. 1948, Aug. 1932); ibidem, Ngarsul (leg. T. TUYAMA, Sept. 13, 1937); ibidem, Ngathpan (leg. T. TUYAMA, Aug. 16, 1939).

Distr. Endemica.

Discriptio *R. palawensis* SCHLECHTERI cum illa *P. Ryozoanae* valde aequalis est, cum exclusione calli retrorsi labelli. Etenim, in Palau, specimen *Rhynchophreatiae* cum callo retrorso non legitur. Mea sententia, hae 2 species valde proximae sunt.

35) **Phreatia pseudo-Thompsonii** TUYAMA sp. nov. (sect. *Euphreatia*).

Epiphytica acaulis humilis erecta 5–10 cm alta caespitosa, radicibus filiforme fibrosis flexuosis pallide fuscis, foliis 4–6 digitatim laxe dispositis lineari faleatim ligulatis 3–10 cm longis 2.5–4 mm latis in textu papyraceis apice obtusis bilobulatis infra medium latissimis et inferiore gradatim angustatis, basi vaginatis, vaginis pallidis ample dilatatis margine membranaceis dorso callosis multi-nervatis 3 mm diametientibus, inflorescentiis e axillis foliorum inferiorum erecto-patentibus folia leviter brevioribus pedunculo 2.5–3 mm longo cum vaginis amplexantibus superioribus in

Fig. 7. A. Flos *Phreatiae Thompsonii* (×20).

B. Flos *Phreatiae pseudo-Thompsonii* (×20).

bracteam exeuntibus subremote obtegentibus, racemo florifero cum pedunculo aequilongo vel breviori dense florato diametro ca. 2.5 mm, bracteis lanceolatis acuminatis patentissimis floribus manifeste longioribus, floribus erectopatentibus albis minutis diametro 2.4 mm, sepalis intermediis oblongo-ovatis acutis uninervatis,

lateralibus late ovatis apice longe acuminatis obliquis intermedio aequilongis uninervatis 1.1 mm longis 0.7 mm latis, petalis oblongis acuto-obtusis 0.8 mm longis 1-nervatis, labello obovato apice brevissime trilobulato margine undululato medio 3-nervato basi angustato breviter unguiculato 11 mm longo 8 mm lato, lobis obtusis intermedio prominentissimis, columna brevissima cylindrica, anthera cucullato-globosa, ovariis sub anthesi cum pedicello 1.2 mm longis clavatis.

Nom. Jap. *Ponape-hime-yureran* (nov.).

Spec. exam. Caroline, Ins. Ponape (leg. G. KOIDZUMI, Jan. 1915—typus in Herb. Univ. Imp. Tokyo.); ibidem (leg. KAMIYA, anno?); ibidem, Kolonia, ad truncos *Artocarpus communis* (leg. S. HATSUSIMA, no. 10739, Jul. 14, 1939, in Herb. Univ. Imp. Kyûshû.).

Distr. Endemica.

Cum *P. Thompsonii* valde affinis, sed foliis tenuioribus floribus omnino minoribus, bracteis inflorescentiis longioribus, labello angustiori et margine saepe grosse undulatis apice emarginato et medio mucronulato differt. Iterum cum *P. inversa* novo-Hiberniensis affinis, sed labello non emarginato, petalis quam sepalis comparatione minoribus differt.

Clavis specierum *Phreatiarum* cum labellis rhombicis
in sectione *Euphreatia*.

1. Labellum basi bicallosum2.
1. Labellum basi non appendiculatum4.
2. Flores longe pedicellati (ca. 3 mm)3.
2. Flores breviter pedicellati*P. carolinensis* (Ponape).
3. Flores majores, sepalis 1.25 mm longis.....
P. collina (Kaiser Wilhelmsland).
3. Flores minores, sepalis 1 mm longis*P. pacifica* (Kusai).
4. Labellum obtusum*P. oxyantheroides* (Neu Mecklenburg).
4. Labellum apiculatum*P. ladronica* (Saipan).

In his 5 speciebus, *P. ladronica* solum cum inflorescentiis folio brevioribus et labellis apiculatis praedita. In callo labelli, haec species cum *P. carolinensis* affinis, sed toto minor, bracteis cum ovariis brevioribus, petalis late ovatis differt.

36) **Pseuderia micronesiaca** SCHLECHTER in ENGLER, Bot. Jahrb. 56, p. 473 (1921).

Arundina Kanehirae YAMAMOTO in Trans. Nat. Hist. Soc. Formosa. 23, p. 20 (1933), syn. nov.

Nom. Jap. *Inu-sekkoku* (nov.).

Spec. exam. Caroline; Ins. Ponape (leg. R. KANEHIRA, field. no. 1610,

Jul. 1931—co-typus *Arundinae Kanehirae*); ibidem (leg. G. KOIDZUMI, Jan. 1915, steril.); ibidem in monte Ngargairu (leg. S. UMEMO, Sept. 26, 1939.).

Distr. In Insula Ponape endemica.

Specimen co-typicum *Arundinae Kanehirae* YAMAMOTO examinatum cum *Pseuderia micronesiaca* conspecificum est. Verisimiliter *Arundina* sp. KANEHIRA cum hac specie congeneris est.

37) **Zeuxine palawensis** TUYAMA in Bot. Mag. Tokyo, **53**, p. 58 (1939).

var. **variegata** TUYAMA var. nov.

Folia medio alba secus venas primarias et secundarias tantum viridia versus marginem gradatim intensius viridescuntia.

Nom. Jap. *Huiri-parao-kinuran* (nov.).

Spec. exam. Caroline, Palau; Ins. Baobeldaob, Ngathpan (leg. T. TUYAMA, Sept. 6, 1939—typus in Herb. Univ. Imp. Tokyo.).

Distr. Endemica.

Explicatio tabulae.

- a. *Peristylus setifer* TUYAMA.
 - b. *Bulbophyllum kusaiense* TUYAMA.
 - c. *Phreatia pseudo-Thompsonii* TUYAMA.
 - d. *Bulbophyllum Hatusimanum* TUYAMA.
 - e. *Agrostophyllum kusaiense* TUYAMA.
 - f. *Phreatia ladronica* TUYAMA.
 - g. *Cestichis dolichostachya* TUYAMA.
 - h. *Disperis palawensis* TUYAMA.
 - i. *Oberonia ponapensis* TUYAMA.
 - j. *Hetaeria Ogurae* TUYAMA.
 - k. *Didymoplexis fimbriata* SCHLECHTER.
 - l. *Eria Uchiyamae* TUYAMA.
 - m. *Bulbophyllum desmanthum* TUYAMA.
 - n. *Cestichis Yamadae* TUYAMA.
 - o. *Liparis palawensis* TUYAMA.
-