DIANELLA STRAMINEA, YATABE.
KIBANA-NO-KIKYÖRAN. キハナノキキヤウラン

K. WATANABE DEL.

New or Little Known Plants of Japan.

No. 30.

Ryōkichi Yatabe, Sc. D.

Dianella straminea, nov. sp.

Nom. jap. Kibana-no-kikyōran. キバナノキ、ヤウラン

(Order Liliaceæ).

Plate I.

Rhizome short, branched. Stem with short internodes, complanate, green, a few inches to nearly a foot high, 4—5 lines broad. Leaves evergreen, distichous, ensiform, shortly sheathing at the base, keeled on the back, scabrous with fine hardened teeth on the edges as well as the keel, 1—2 feet long, 1—1½ inches broad. Peduncle terminal, subterete, with sheathing and keeled reduced leaves or empty bracts, 1—3 feet long including the panicle, the empty bracts 1—3 inches long. Panicle terminal, a few inches

New or Little Known Plants of Japan.

to a foot long; bracts broadly lanceolate, or deltoid, amplexicaul, keeled and often scabrous on the back, the lowest about 1 inch long, upper ones much smaller. Flowers $\frac{1}{2}$ inch across; pedicels slightly arcuate, 2—6 lines long, articulated under the flower. Perianth-segments 6, in 2 series, pale-yellow (sulphureous in bud and stramineous in flower), marcescent, subequal, oblong, obtuse or retuse and with a few minute hairs at the apex, with thin margins and 5 parallel nerves, reflexed when fully open, closed over the ovary after flowering, about 3 lines long, 1 line broad. Stamens 6, hypogynous, opposite the perianth-segments; filaments linear, $1\frac{1}{4}$ lines long, dilated at the apex, the dilated part yellow and $\frac{1}{2}$ line long; anthers 2-celled, erect, linear, adnate, yellow, connivent and forming a tube around the style in bud, 1 line long, the cells opening introrsely in an apical oblique pore. Ovary sessile, short, subtrigonal, 3-celled, pale-green, $\frac{1}{2}$ line long; style filiform, 2 lines long; stigma terminal, small; ovules 4 in 2 rows in each cell, oblong, anatropous; placentæ axile. Berry subglobose, somewhat depressed, blue, 3—5 lines across, when young crowned with the detached and shriveled upper portion of the perianth, the lower portion of which remains attached to the base of the berry. Seeds few, subovoid, with 3—4 prominent ridges; testa black, smooth, shining; embryo clavate, somewhat curved, in the axis of a white fleshy albumen.—Fl. Summer—autumn.

This plant is cultivated from Luchu in the Botanic Garden of the Imperial University.

This species differs from *Dianella ensifolia*, Redoute, chiefly in the following respects:—The leaves of the former are scabrous with fine teeth on the edges and keel, while those of the latter are smooth all over. The flowers of the former are pale-yellow, while those of the latter are pale-blue. The berries of the former are depresso-globose, while those of the latter are globoso-ovoid. The seeds of the former are subovoid with 3–4 prominent ridges, while those of the latter are ovoid and somewhat compressed.

Plate I. Fig. 1, lower part of plant; 2, panicle; 3, vertical section of flower; 4, pistil; 5, ovule; 6, ventral and dorsal views of stamen; 7, outer perianth-segment; 8, inner perianth-segment; 9, floral diagram; 10, berry; 11, vertical section of berry; 12, transverse section of berry; 13, seed; 14, vertical section of seed:—*all except 1, 2, and 10 enlarged.*

二三植物ノ化學的實驗

醫科大學教授

下山順一郎氏口演

澤田駒次郎筆記

下山順一郎氏ハ去ル十一月東京植物學會ニ於テ二三植物ノ化學的實驗ト云ヘル演題ニ就キテ演述セラレタ

リ左ノ一編ハ其演述ノ大要ヲ記スルモノトス

余ハ二三ノ植物ヲ化學的實驗ニ徴セリ茲ニ其成績ニ就キ聊カ陳述セントス

凡ソ植物ハ其莖、根、葉、花、果實、種子等何レノ部分ヲ問ハズ概シテ各々特異ノ物質ヲ含有スルハ已ニ諸君ノ