P-189 TP-10による前立腺高温度治療と経尿道的前立腺摘除術との比較

杏林大学

奴田原紀久雄、太田雅也、金城真美、野田治久、渡辺和吉、加藤可顕、宮田晃臣、村田明弘、米田龍生、桶川隆嗣、小島美保、東原英二

【目的】前立腺高温度治療装置TP-10と経尿道的前立腺摘除術(TUR-P)の治療成績を比較検討した。

【対象と方法】TP-10による治療を施行した30例（平均年齢72.9歳）とTUR-Pを施行した11例（平均年齢70.6歳）の前立腺肥大症患者を対象として、術前、術後1カ月目、3カ月目にIPSS, QOL index、尿流量検査、前立腺容量を比較検討した。副作用の検討のため術前、術直後、術後7日目に血算、血液生化学検査を行った。TP-10による治療は外来局所麻酔下、尿道温度50℃、60分単回治療とした。

【結果】TP-10による治療ではIPSS, QOL indexはともに術後有意に減少した（IPSSは術前18.4±7.1、術後1カ月12.8±7.1、3カ月9.7±6.2、QOL indexは術前4.5±1.1、術後1カ月2.8±1.6、3カ月2.4±1.2）が、Qmaxと前立腺容量は有意の変化を示さず、残尿量は術後1カ月、3カ月で、術前より有意に減少した（術前89.7±18.7、術後1カ月48.9±41.3、3カ月49.9±38.1ml）。一方TUR-PではIPSS, QOL index, Qmax、残尿量、前立腺容量のすべてにおいて術前より有意の改善と縮小を認めた。術後のヘマトクリット、血清アルブミン値はTP-10では減少しなかったが、TUR-Pでは有意に減少した。

【結論】TP-10による前立腺肥大症の治療は、TUR-Pほど顕著な治療効果は示さないが、自覚症状、他覚症状の改善をもたらすため、局所麻酔しか施行できない症例や、外来での治療を希望する症例に対して有効な方法となりうると考えられた。

P-191 HIGH ENERGY TRANSURETHRAL MICROWAVE THERMOTHERAPY (TUMT) VERSUS TRANSURETHRAL RESECTION OF THE PROSTATE (TURP) IN THE TREATMENT OF BENIGN PROSTATIC HYPERPLASIA (BPH)

Frans M.J. Debruyne, Jean J.M.C.H. de la Rosette, Nijmegen, The Netherlands.

PURPOSE : In the evaluation of a new technology for the treatment of BPH, the results should be compared with those obtained using a generally accepted method. We report on the 1-year follow-up of a prospective randomized study comparing high energy TUMT and TURP in patients with voiding complaints because of BPH.

METHODS : Patients with symptomatic BPH were treated by either TURP (21 patients : mean age 69.6 ± 8.5 years) or TUMT (31 patients : mean age 69.3 ± 5.9 years) and assessed using the Madsen symptom score, measurements of voiding parameters, transrectal ultrasound and cystometry including pressure flow analyses. Examinations were repeated at fixed intervals for up to 12 months post treatment.

RESULTS : After both TURP and TUMT there was a significant improvement in all clinical parameters. At 1 year follow-up, a symptomatic improvement of 78 % in the TURP group was found and of 68 % in TUMT group. The improvements in free flow rate are 100 % and 69 % in the TURP and TUMT group respectively. For both groups a significant relief of bladder outlet measures was found. No serious complications occurred in either group while 1 patient in each group required a retreatment.

CONCLUSION : Satisfactory results were obtained after both treatments with improvements following high energy TUMT in the same range as following TURP treatment.

P-190 THERMOTHERAPY IN LARGE (> 60 g) BENIGN PROSTATE HYPERTROPHY WITH PROSTATRON 2.5.

VALLANCIEN Guy, VEILLON Bruno, HAROUNI Mohamed, GUILLONNEAU Bertrand. Institut Mutualiste Montsouris - Paris - France.

PURPOSE : Transurethral microwave thermotherapy (TUMT) at high temperature (60 to 70 °C) with Prostatron 2.5 (EDAP Technomed) was proposed to symptomatic patients with large obstructive prostates, originally candidates for open prostatectomy.

TECHNIQUE : ambulatory procedure, 1 session of 1 hour with patients under neuroleptic analgesia. Post TUMT bladder catheterisation was systematic.

INDICATIONS : 25 patients (mean age 67) with large obstructed prostate, average weight 78 g (55-133) entered the protocol after signing the consent form.

RESULTS : The energy delivered by the equipment averaged 181 kJ. Bladder catheterisation averaged 2 weeks (8 to 45 days). 50 % of catheters were removed at day 8.

RESULTS	PRE	3 MONTHS	6 MONTHS
IPSS	17	7	4
Q MAX (ml/s)	7.8	14.3	19.8
Residual Vol.(ml)	100	25	26

No significant side effects were noticed : no impotence, no incontinence, no urethral stenosis, no transfusion. A secondary TURP (due to median lobe) had to be performed on one patient.

CONCLUSION : TUMT with Prostatron 2.5 in large (> 60 g) and obstructive prostates considerably improves both objective (Qmax +150 %) and subjective (IPSS -75 %) symptoms at 6 months. Median lobe remains an exclusion criteria. Results are continuously improving over the 6 month period.

P-192 TUMT AT 1296 MHZ: A MINIMAL INVASIVE ALTERNATIVE FOR OBSTRUCTED HIGH RISK BPH PATIENTS

THÜROFF, STEFAN; CHAUSSY, CHRISTIAN
MUNICH HARLACHING, GERMANY (presented by Prof. Chaussy)

INTRODUCTION AND OBJECTIVES: Geriatric obstructed BPH patients with indwelling catheter are often scored as high risk ASA III/IV patients and are not preferable clients for anaesthesia requiring endourological surgery. This results often in a life with indwelling catheter for these patients, handicapped by many other age related health problems.

MATERIAL AND METHODS: Offering high intensive transurethral microwave therapy at 1296 MHZ to these patients we analyzed a group of 102 patients within the last two years who underwent this therapy in sedoanalgesation.

The mean age of 80 years and the prostate weight of ~ 40 gramm showed this group different to our TURP patients of the last two years (ø age 66 ø resection weight 29 gramm).

With a bladder volume > 150 cc, exclusion of bladder-/prostate cancer and two unsuccessful catheter releases, the treatment was performed. All patients had a suprapubic tube and perioperative antibiotic prophylaxis.

A 60 min computer supported TUMT with an average of 160 kJ was performed.

RESULTS: Severe side effects of high intensive thermotherapy did not occur. No sphincter lesion, no bleeding, no rectum wall damage or fistula was noticed. After an average period of 6 (1-12) weeks suprapubic drainage almost 80% of the patients voided spontaneously with tolerable residual urine and uroflow. The differentiated data of 102 patients with follow up to two years will be presented.

CONCLUSION: High intensive TUMT at 1296 MHZ proved to be a valid therapy option for this special patient group. It's a minimal invasive, anaesthesia free, outpatient procedure with simple selection criteria and a high success rate of 80%.