EW2-2**Role of ureteroscopic biopsy in the management of upper urinary tract malignancies**

Department of Urology, Saiseikai Shimonoseki General Hospital
Koji Shiraishi, Satoshi Eguchi, Jun Mohri, Yoriaki Kamiryo

Purpose : With recent progress in ureteroscopy, ureteroscopic inspection and pathological diagnosis became more important in the management of upper urinary tract malignancies. We examined the accuracy of ureteroscopic biopsy in the diagnosis of upper urinary tract transitional cell carcinoma (TCC). Furthermore, we discussed whether conservative management (Holmium YAG laser, transurethral resection or partial ureterectomy) is possible or not based on the pathological diagnosis.

Patients and Methods : Forty consecutive patients underwent ureteroscopic observation and biopsy using a 3 Fr cold cup forceps. Pathological diagnoses of the biopsy sample and the grade or stage of surgically resected tumors were compared. In the patients with grade 1 or 2 TCC diagnosed by ureteroscopic biopsy, disease-free and cancer-specific survival rates were determined whether conservative management is performed or not.

Results : There were no major complications by ureteroscopic biopsy. Nineteen percent of specimens appeared as false-negative, whereas malignancies were suspected in most cases based on the results of urine cytology and ureteroscopic inspection. Pathological grading of biopsy was almost coincide with that of surgically resected specimen, while 85% of grade 2 or 3 TCC appeared muscle invasive disease ($P_t \leq$). There were no significant differences in disease-free and the survival rate between the group of nephroureterectomy and those of conservative management, except for grade 3 or P_t tumors.

Conclusions : Because the disease is often multifocal and has a high likelihood of recurrence, it has been traditionally treated by nephroureterectomy. Ureteroscopic biopsy is safe and accurate if sufficient sample is obtained. Partial ureterectomy, but not endoscopic management, may be a feasible option for patients with grade 2 tumor. Ureteroscopic biopsy should be aggressively performed to the patients who require conservative management.

EW2-3**Endoscopic treatment of upper urinary tract tumors**

Department of Urology, Saga Medical School
Tsunehiro Tsukahara, Jiro Uozumi, Zenjiro Masaki

(Purpose) We present our experiences of endoscopic treatment for upper urinary tract tumors, attempting to outline the usefulness and problems of this treatment.

(Materials and Methods) We have endoscopically treated 15 cases of suspected upper urinary tract malignancy since April 1996. The patients consist of 12 male and 3 female. Age ranged 22 to 83 years old. Rigid pyeloscope (19.5 and 25 Fr.) were used in two renal pelvic tumors and mini-TU-resectoscope was utilized for the treatment of 1 tumor located at the lowermost ureter. Rigid or fiber ureteroscope (6.9–12 Fr.) were utilized in the other 12 cases depending on tumor size or location. Biopsy forceps were most frequently used for tumor resection in combination with electro-coagulation and Ho-YAG LASER. A "Snare" with electro-currenty was used in resecting large tumor.

(Results) Among 15 patients, 9 had malignant lesions and 8 of these 9 cases were transitional cell carcinomas, less than 15 mm in diameter. The largest tumor occurring in the renal pelvis was percutaneously treated in 2 separate sessions using Snare. Tumor recurrences in the ureter were observed in 2 cases. One was treated endoscopically again but another one who had had contralateral nephrectomy in the past underwent open resection of the entire ureter and placement nephrostomy. Although 4 cases ultimately underwent nephroureterectomy following the endoscopic treatment, one of 4 had no malignant lesion proven by histological examination of the resected specimen. Six benign lesions (2 renal pelvis and 4 ureter) consist of polyp, polypoid lesion and papilloma. Two cases were treated with loop resection and the remaining 4 with biopsy forceps. As a major complication, postoperative loss of the renal function was observed. Complete obstruction of the ureter seen in this case was considered to be caused by too early removal of double-J catheter.

(Conclusions) Endoscopic treatment could be one of choice for upper urinary tract tumors in the selected case. Since the procedure had to be done in the very limited space, biopsy forceps and a Snare with electro-currenty seemed to be very useful tool. Further study is needed.