YW3-15 Modulation of phenotype of human prostatic stromal cells by transforming growth factor- β s.

Department of Urology, Sapporo Medical University, Sapporo, JAPAN

Toshihiro Hisataki

Background : We investigated the effects of transforming growth factor (TGF) - β s on morphological and receptor phenotypes, as well as proliferation of human prostatic myofibroblast cell line (PMF) and prostatic stromal cell line (PSC).

Methods : The effects of TGF- β s on morphological changes and proliferation of the cells were studied by immunohistochemistry and bromodeoxyuridine assay, respectively. The expression of alpha 1-receptor subtypes was measured by real time quantitative RT-PCR and the radioligand binding assay for the receptors was also performed.

Results : TGF- β 1,2 and 3 induced expression of desmin and myosin of cells of the PMF, and significantly inhibited their growth. The alpha 1a -receptor was expressed only in the PSC and alpha 1b and 1d, in all cell lines. TGF- β 1 suppressed the expression of all three subtypes of the alpha 1-receptor. The binding sites of cells of all the cell lines were reduced by treatment with this growth factor.

Conclusion : TGF- β s may induce human prostatic stromal cells to express the smooth muscle phenotype and inhibited their growth. However, the growth factor reduced the binding sites of the receptor and suppressed mRNA expression of its subtypes, suggesting that morphological and receptor phenotypes may be regulated via more than one pathway by TGF- β (s).

YW3-16 A sub-renal capsule model for benign prostatic hyperplasia

Department of Urology, Osaka University Graduate School of Medicine

Tetsuya Takao

BACKGROUND. One of the main constraints in understanding the etiology of benign prostatic hyperplasia is the lack of suitable model systems. To address this issue, we have used murine prostatic cell lines to establish a novel in vivo model for studying prostatic cell interactions.

METHODS. Luminal, basal, and smooth muscle (SM) cell lines were inoculated alone or in combinations under the renal capsule of intact or castrated male mice, and the growth and composition of prostatic tissue in the absence or presence of doxazosin was determined.

RESULTS. Both the luminal and basal cell lines reconstituted prostatic tissue if co-inoculated under the renal capsule with normal SM cells, whereas none of the lines formed significant tissue when inoculated alone. The growth of prostatic tissue formed from co-inoculation of basal and SM cells was androgen responsive. In addition, a significant reduction in prostatic tissue was noted in animals treated with doxazosin.

CONCLUSION. We have established an in vivo model that uses prostatic epithelial and SM cell lines for investigating cellular interactions between epithelial and SM cells that regulate prostatic growth and function. This model will be useful for elucidating the cellular interactions that regulate tissue growth in vivo.

YW3-17 NEONATAL EXPOSURE TO ENVIRONMENTAL DISRUPTORS DECREASES ESTROGEN RECEPTOR- β AND SUPPRESSES APOPTOSIS OF THE PROSTATE AFTER PUBERTY

Department of Urology, Yaizu Municipal General Hospital

Shinsuke Hayami

Objective : We examined the mechanism of prostate enlargement to estimate ER β , androgen receptor (AR), morphometry, and apoptosis of prostates in rats which were exposed to environmental endocrine disruptors during the neonatal period.

Materials and Methods : Subjects were 32 male Wistar rats. Sixteen rats were injected with 25 μ g Estradiol on days 1,3, and 5 after birth (neonatal estrogen exposure group) ; the others served as the control group. Each groups' rats were sacrificed on days 80,125,160, and 180. Serum testosterone, AR and ER β mRNA, their histological compositions and PCNA of the ventral prostate were measured.

Results : Prostate weights showed no significant differences between groups. Epithelial components of the exposure group showed relatively increased at day 160. At days 160 and 180, AR and ER β mRNA were low in the exposure group. Expression of PCNA at day 160 was increased significantly for rats exposed to estrogen.

Conclusions : Low-dose estrogen exposure in the neonatal period decreases ER β after puberty and suppresses apoptosis after middle age. In puberty, decreasing ER β led to glandular hyperplasia of the prostate.

YW3-18 The characterization of proprotein convertases in the prostate

Department of Urology, Mie University Faculty of Medicine

Katsunori Uchida

Introduction and Objective : Proprotein convertases (PCs) have been known as processing proteases that convert a wide variety of precursor proteins (e. g. TGF- β s, MMPs, and integrins) to their mature active form. The aim of this study was to evaluate the function of PCs in the development and carcinogenesis of the prostate

Methods : PCs gene expressions in neonatal rat ventral prostate (VP) and human prostate cancer cell lines were examined and PCs activity was measured. Effects of the PCs inhibitor , decanoyl-Arg-Val-Lys-Arg-chloromethylketone (CMK) on neonatal rat VPs and LNCaP were analyzed.

Results : Selective PCs gene expressions were detected by RT-PCR. PCs activity was modulated in the presence of testosterone. Addition of CMK inhibited Testosterone-induced growth and branching of rat VPs. Exposure of LNCaP to CMK resulted in the loss of luminal cell phenotype and induction of basal cell phenotype.

Conclusions : These data suggest that PCs may be responsible for the development and differentiation of the prostate.