AS-3

Laparoscopic Radical Cystoprostatectomy

Catholic University of Korea, Republic of Korea

Tae-Kon, Hwang

An advance in manufacturing laparoscopic instruments and improvement of surgical technique, there has been a relatively recent interest in laparoscopic pelvic surgery. This interest initially led to the development of techniques for laparoscopic radical prostatectomy (LRP). With the realization of the technical feasibility of LRP and encouraging short-term outcome, the interest moves to the bladder which is the only remaining intra-abdominal organ naturally.

The laparoscopic radical cystoprostatectomy (LRC) and urinary diversion is one of the most challenging operations in Urology. The laparoscopic procedure of the radical cystoprostatectomy is closely related with Laparoscopic radical prostatectomy (LRP). Nowadays, the LRP patients are increased dramatically in Asia. But the number of LRC patients is much less than LRP patients. The surgical experience on LRP makes the surgeon comfortable to do LRC. Intestinal surgery for urinary diversion, which is most difficult part in LRC, can be performed extracorporeally through the minilaparotomy where the specimen was extracted. Laparoscopic surgery is also associated with better preservation of the immune system than open surgery, which results in a decreased incidence of infectious complications and leak from the intestinal anastomosis site.

It is clear that the LRC is becoming more accepted worldwide. As LRC becomes better standardized it is likely that operative times will reduce further, as was the case with LRP. Satisfactory long-term oncological and functional outcomes will determine whether the procedure develops into a real contender in the future.

Robotic-assisted laparoscopic radical cystectomy, intra-abdominal formation of an orthotopic ileal neobladder and laparoscopic nerve- and seminal-sparing cystectomy with orthotopic ileal neobladder can be performed in the advanced hospital.

Even though there remained the technical difficulty in the field of LRC with intracorporeal urinary diversion, the problem of night time incontinence and erectile dysfunction, the LRC technique would be popularized as the LRP in the future.

With the improvement of easy intracorporeal intestinal anastomotic technique or the development of device like a push button intestine anastomosis, the LRC could be performed in general hospital with experienced doctors.