S14-1

Recurrence hazards of superficial bladder cancer after TURBT.

Department of Urology and Andrology, Graduate School of Comprehensive Human Sciences, University of Tsukuba, Tsukuba, Japan

Shiro Hinotsu, Hideyuki Akaza

Introduction

Superficial bladder cancer can be treated by TURBT followed by intravesical instillation of anti-cancer drug or BCG with minimal morbidity, but the patients are at high risk for tumor recurrence. We introduced the smoothed hazard function plots of tumor recurrence, and this technique made it possible to understand the time course change in the hazard rate.

Natural history of superficial bladder cancer

In Japan, several clinical trials have been conducted for the purpose of finding the optimal dose and schedule of anthracycline instillation after TURBT. In our meta-analysis of JUCRGA (Japanese Urological Cancer Research Group for Adriamycin/Farmorubicin) clinical trials, we were able to depict graphically the characteristic natural history of tumor recurrence after TURBT, and found that there were two phases of recurrence, an early phase and a late phase. In the early phase, it was demonstrated that each of subgroups classified on the basis of the background factors showed a distinguishable pattern of tumor recurrence. One feature is that patients with multiple tumors show an early peak of recurrence, and another feature is that patients with recurrent tumor show a higher hazard.

Intravesical chemotherapy and immunotherapy

From our meta-analysis of JUCRGA data, it was revealed that intravesical chemotherapy provided protection against tumor recurrence only in the early phase, and no prophylactic effect in the late phase. From the data of the trial conducted BCG Tokyo 172 Strain Study Group, BCG instillation may prevent tumor recurrence not only in the early phase but also in the late phase of recurrence compared with Doxorubicin instillation.

Conclusion

From the data of superficial bladder cancer trials, there are two patterns of tumor recurrence after TURBT, early phase and late phase. Intravesical chemotherapy is effective mainly in the early phase, and intravesical immunotherapy may reduce the risk of recurrence both in early phase and in late phase.

S14-2

Risk factors and prediction of recurrence in superficial bladder cancer

Department of Urology, Nara Medical University, Nara, Japan¹⁾, Oncogene Research Unit/Cancer Prevention Unit, Tochigi Cancer Center, Utsunomiya, Japan²⁾

Kiyohide Fujimoto¹⁾, Yoshitomo Chihara¹⁾, Yoshihiko Hirao¹⁾, Kokichi Sugano²⁾

Superficial bladder cancers will recur in 60-70% of cases when they are untreated after transurethral resection (TUR), and 15-25% will show grade and stage progression. To accurately assess recurrence, complete TUR of all visible tumors is firstly required, and secondly, a reliable histopathological diagnosis of appropriate TUR specimens bearing the muscle layer and marginal mucosa and systematic random biopsy specimens are mandatory. It is well-known that the number, size, grade, and stage of the tumors are important pathological risk factors of recurrence. Especially, T1G3 tumors as well as associated carcinoma in situ often have high risk of recurrence presenting with stage progression or resistance to the conservative treatment. On the contrary, our clinical trials for intravesical instillation therapy revealed that the tumor multiplicity at the initial presentation was the sole risk factor of recurrence for the Ta-I, G1-2 tumors (n = 144) in the non-instillation control groups. On the other hand, recurrence is considered to be metachronous multi-focality, and recent genetic studies have demonstrated that individual multi-focal tumors develop initially with the common genetic alterations, and progress later with independent multi-step genetic alterations. In our comprehensive survey of the chromosomal and genetic alterations detected by microsatellite markers in Ta tumors, the chromosomal arms 9p and 9q were the most common allelic-loss regions in Ta tumors, but the high-grade Ta tumors clearly showed wide-range chromosomal and genetic alterations including the chromosome 17p where the p53 gene alterations are frequently seen in invasive diseases. Therefore, chromosomal or genetic alterations identical to the primary tumor in the exfoliated urothelial cells are likely to be promising markers for monitoring recurrence and progression in the individual patient. Herein, we present the results of our pathological and genetic survey on the superficial bladder cancer.