The Application of Copper Waterline on Laying Performance and Gut Health of Aged Laying Hens

Ning Ma^{*}, Min Liu^{*}, Mengze Song, Sheng Li, Xiaoyan Lin, Hongchao Jiao, Xiaojuan

Wang, Jingpeng Zhao, Shuhong Sun and Hai Lin

Shandong Agricultural University, Shandong Provincial Key Laboratory of Animal Biotechnology and Disease Control and Prevention, No. 61 Daizong Street, Taian City, Shandong Province, 271018, China

Running Title: Copper waterline in laying hens

^{*}These two authors contributed equally to this work and should be considered co-first authors.

Correspondence: Dr. Hai Lin and Dr. Shuhong Sun, Shandong Agricultural University, Shandong Provincial Key Laboratory of Animal Biotechnology and Disease Control and Prevention, No. 61 Daizong Street, Taian City, Shandong Province, 271018, China.

(Email: hailin@sdau.edu.cn, sunshuhong@sdau.edu.cn)

Abstract

The effect of the application of copper waterline on the performance and gut health of aged laying hens was evaluated in this study. Forty-eight 70-week-old laying hens were divided into two groups (three replicates of eight hens each): control and copper (Cu) groups provided with normal polyvinyl chloride (PVC) waterline or Cu waterline. The laying performance was measured during the four-week period of the experiment. The intestinal antioxidant status and the microbiota diversity of the cecal content were determined. Moreover, a bacteriostasis test on *Escherichia coli* and *Salmonella enteritidis* was conducted after inoculation in waterline and hens, respectively. The water Cu^{2+} content was increased by Cu waterline compared to the control ($P < 0.05$). Cu waterline had no detectable effect on most production performances, however, it increased the egg weight ($P < 0.05$). Cu waterline increased the Cu level in the eggshell. Cu level in excreta increased with time, especially in the final two weeks, however, there was no significant change in fecal Cu excretion. The lipid peroxidation product malondialdehyde content in ileum decreased ($P < 0.01$), while the activities of CuZn-superoxide dismutase (SOD) of ileum and glutathione peroxidase (GSH-PX) activity of jejunum and ileum increased after Cu treatment. The relative abundance and richness of cecal microbiota increased after Cu treatment ($P < 0.05$). Cu waterline changed the microbial composition, including the increased proportion of *Methanocorpusculum*, *Paludibacter*, and decreased proportion of *Fucobacterium*, *Anaerobiospirillum*, and *Campylobacter*. The colonization of *E. coli* and *S. enteritidis* in

Cu waterline was suppressed by Cu treatment, indicating that Cu waterline had potential antibacterial properties. The result suggests that Cu waterline could inhibit the colonization of pathogenic microorganisms such as *E. coli* and *Salmonella* and facilitate the enrichment of cecal microbiota diversity.

Keywords: copper waterline, antioxidant capacity, microbial diversity, antibacterial activity, laying hens

Introduction

Copper (Cu) is essential in several enzyme systems, including cytochrome oxidase, tyrosinase, lysyl oxidase, and superoxide dismutase, by serving as co-factor of a variety of intracellular and extracellular enzymes (Klasing 1998). The Cu requirement for laying hens is not available in the National Research Council (1994) list. Dietary Cu supplementation level had no significant influence on the laying performance of hens but decreased serum cholesterol concentration, suggesting that 125 mg/kg of Cu provide adequate supplementary concentrations for laying hens (Balevi and Coskun 2004; Lien *et al.* 2004). For late-phase hens, a corn-soybean basal diet (containing 10.3 mg Cu/kg) might be sufficient to meet their maintenance and production requirements (Li *et al.* 2018). The lack of copper in laying hens leads to anemia and abnormal egg size and shape (Baumgartner *et al.* 1978).

In practice, Cu addition at prophylactic levels is used as a growth promoter in poultry production (Leeson, 2009; Bortoluzzi *et al.* 2020). However, high levels of Cu supplementation may have two negative effects: oxidative damage induced by Cu overloading (Zhang *et al.* 2000; Toplan *et al.* 2005) and elevated excretion of Cu in the excreta (Skrivan *et al.* 2006). More than 90% of copper is excreted with droppings when supplemented at a pharmacological concentration (Mohanna and Nys, 1998), leading to heavy metal waste in livestock and poultry breeding.

The bactericidal action of Cu depends on the concentration of free ionic Cu in solution (Zevenhuizen *et al.* 1979). Acidic copper sulfate-based commercial sanitizer is

successfully used at various intervention points of poultry processing (Russell, 2008). It has recently been proven that Cu could reduce the conjugative transfer of resistance plasmids from extended-spectrum β -lactamase-producing *E. coli* (Buberg *et al.* 2020). *In vivo*, supplementation with 187.5 mg/kg of Cu from Cu sulfate pentahydrate had no effect on the number of ileal lactobacilli of birds (Pang *et al.* 2009).

In this study, the use of a copper waterline was evaluated in laying hens. The laying performance, Cu intake and excretion, antioxidant capacity, and microbial composition in cecal content were measured. The inhibition of *E. coli* and *S. enteritidis* by Cu waterline was respectively measured *in vitro* and *in vivo*. *E. coli*, a pathogenic bacteria that exists widely in animals and nature, was used to detect the inhibitory effect of Cu waterline on the spread of bacteria. *S. enteritidis*, an epidemic pathogen, was employed to evaluate the inhibitory effect of Cu waterline on the bacterial colonization in the gut of chickens. Our research sheds new light on copper waterline applications in aged laying hens by explaining its potential effect of improving antioxidative and antibacterial effects and changes in microbial composition.

Materials and Methods

Animal Ethics

All study procedures were approved by the Animal Care and Use Committee of Shandong Agricultural University and were in accordance with the Guidelines for Experimental Animals established by the Ministry of Science and Technology (Beijing, P. R. China).

Birds and experiment design

Forty-eight 70-weeks old Hy-line Brown laying hens with an average body weight of 2.25 ± 0.01 kg were randomly assigned to two groups and reared in cages equipped with a normal poly vinyl chloride (PVC) waterline (control) or copper waterline (Cu). Each treatment had three replicates of eight hens. The laying hens were reared in cages (60-cm length $\times$ 45-cm width $\times$ 50-cm height), with two birds in each cage. Drinking water was provided using nipple drinkers (1 nipple/cage.) via waterline. Duplicate water samples (10 mL) were obtained from each nipple drinker 4 times (6 repeats each time) every week from the control and Cu waterlines, and the water samples were mixed and stored at 4 °C for further measurement. All laying hens had free access to feed and water. The lighting program was 16L:8D. The experimental diet (Table 1) was formulated to meet the recommendation by NRC (1994). The experiment lasted four weeks. Feed intake, water intake, and egg production were recorded daily. The feces were collected daily for every replicate. After drying and weighing, the Cu content was measured, and the average daily fecal Cu excretion was calculated. At the end of the experiment, blood samples were obtained from the wing vein using heparinized syringes and placed into ice-cold tubes. Plasma was obtained after centrifugation at 400 g for 10 min at 4 °C and was stored at -20 °C for further analysis. After the blood samples were obtained, laying hens were killed by exsanguination (Close, 1997; Wang *et al.* 2020). The content of the cecum was collected, snap-frozen in liquid nitrogen, and stored at -80 °C. The duodenum, jejunum, and ileum mucosae were collected, snap-frozen in liquid nitrogen,

and stored at -80 °C for enzyme activity assay.

Microbial analysis of the cecal contents

Total bacterial genomic DNA samples were extracted, and the quantity and quality were measured. After the detection of DNA, the PCR amplification of the bacterial 16S rRNA gene V4-V5 region was performed on the Illumina Miseq PE250 platform. After initial denaturation, annealing, and extension, the PCR amplicons were obtained, purified, quantified, and pooled in equal amounts. The separation and amplification 16S rDNA and quantification of the microflora composition were performed at the Annoroad Gene Technology Company (Beijing, China). The average number of observed operational taxonomic units (OTU) at 97% identity was determined using UCLUST in QIIME software (Version 1.9.1). To estimate the portion of the diversity covered by our subsampling, the diversity of the OTUs found in the samples and their evenness were measured. The OTUs were taxonomically annotated based on the Silva (<http://www.arbsilva.de>) taxonomic databases. The QIIME software was used to generate the species abundance table at different levels (phylum, class, order, family, genus, species) and displayed with R software (Version 2.15.3). The alpha diversity of samples (Chao 1, Shannon index, Rank abundance curve) was analyzed using R software. To assess beta diversity, the unweighted UniFrac distance metrics were produced by QIIME software, and principal coordinate analysis (PCoA) was performed.

Cu Analysis

Diet, fecal, and serum Cu levels were analyzed by graphite furnace atomic absorption

as described by Meeravali and Arunachalam (1997). Cu concentration in water was determined by ICP-AES 7000SERIES atomic absorption spectrometer (Thermo Fisher) at an absorption wavelength of 324.7 nm.

T-AOC, CuZn-SOD, GSH-PX and Malondialdehyde (MDA) Measurement

The activity of total superoxide dismutase (T-SOD), CuZn-SOD, total antioxidant activity (T-AOC), and activity of glutathione peroxidase (GSH-PX) of the serum and intestinal mucosa homogenate were measured with commercial kits (Jiancheng Bioengineering Institute, Nanjing). The level of lipid peroxides released from the small intestinal tracts, estimated as the level of lipid peroxides released from the small intestinal tracts, was estimated as TBARS according to Huang *et al.* (2015).

Escherichia coli Bacteriostasis Test

E. coli was obtained from the Avian Disease Centre of Shandong Agricultural University (Shandong, China). The *E. coli* strain was seeded on LB (Solarbio, China) agar to obtain isolated pure colonies, and a single colony was inoculated into LB broth and incubated overnight at 37 °C while shaking at 180 rpm (QYC-2102, Fuma, China) for 10 h. The concentration of *E. coli* was adjusted to 8.22×10^6 colony-forming unit (CFU)/mL in sterile saline.

Three pipes of Cu waterline and control waterline (1.8 m long) filled with tap water were respectively injected with 1 mL sterile saline or *E. coli* solution at a concentration of 8.22×10^6 CFU/mL. Afterward, a water sample was obtained every 24 h for five times. The water samples (100 µL) were serially diluted 10-fold with sterile

phosphate-buffered saline (1:10, w/v), and then screened on MacConkey plates (Hopebio, Qingdao, China). After incubation at 37 °C for 24 h, the CFU of *E. coli* was enumerated.

Salmonella Challenge Test

A strain of *S. enteritidis* (SDJX-1) isolated from chickens was obtained from the Avian Disease Centre of Shandong Agricultural University. A single colony was selected from the xylose lysine deoxycholate agar plate and transferred into a tube containing 5 mL tryptic soy broth and then incubated at 37 °C while shaking at 180 rpm (QYC-2102, Fuma, China) for 16 h. The concentration of *S. enteritidis* was adjusted to 2.53×10^5 CFU/mL in sterile saline.

Ten one-day-old SPF White Leghorn layer-type chicks were obtained from Jinan SPAFAS Poultry Company (Jinan, China). The chickens were reared individually in a cage within an environmentally controlled chamber. Each cage had a feeder and nipple drinker. The chicks were fed a commercial diet and had free access to feed and water throughout the experimental period. On Day 8, the chicks were orally challenged with 0.2 mL of *S. enteritidis* (2.53×10^5 CFU/mL). Afterward, the chicks were randomly divided into two groups and were reared in cages equipped with a control waterline or Cu waterline. The experiment lasted 12 days, and a water sample was obtained every day from every nipple ($n = 5$ for each treatment). The water sample (100 µL) was serially diluted 10-fold with sterile phosphate-buffered saline (1:10, w/v), screened on xylose lysine deoxycholate plates (Hopebio, Qingdao, China), incubated at 37 °C for 24

h, and then the CFU of *S. enteritidis* was enumerated.

Statistical analysis

Prior to analysis, all data were examined for the homogeneity and normal distribution plots of variances among the treatments by using the UNIVARIATE procedure. The main effect was evaluated using a one-way ANOVA with the Statistical Analysis Systems statistical software package (Version 8e, SAS Institute, Cary, NC, USA). The antibacterial effect against *Salmonella* was analyzed by two-way ANOVA with SAS software. The data of 16S rRNA sequencing were calculated with QIIME (Version 1.9.1) and displayed with R software (Version 2.15.3). $P < 0.05$ was considered statistically significant.

Results

There were no significant differences in body weight, average feed intake, water intake, and egg production between the control and Cu groups. Egg weight, however, increased with Cu treatment ($P < 0.05$, Table 2). Cu treatment increased the Cu level in the water and Cu intake from water ($P < 0.01$, Table 3). Similarly, the daily Cu intake significantly increased ($P < 0.01$). Cu waterline had no significant influence on Cu concentration in serum and egg content, whereas increased Cu level was observed in the eggshell ($P < 0.05$, Table 3). Cu level in the excreta was not significantly changed by the Cu treatment in the first and second week but increased at Week 3 ($P < 0.05$) and Week 4 ($P < 0.1$, Fig. 1A). In contrast, the average daily fecal Cu excretion during the four-week period was not significantly changed by Cu treatment (Fig. 1B).

Cu treatment had no significant influence on serum T-SOD and CuZn-SOD activity and T-AOC (Fig. 2A, B, C). The activity of CuZn-SOD did not change in the duodenum and jejunum but increased ($P < 0.05$) with Cu treatment in the ileum (Fig. 3 A). In contrast, the activities of T-SOD in the duodenum, jejunum, and ileum mucosa did not change with Cu treatment (Fig. 3B), same as the T-AOC level ($P = 0.139, 0.311,$ and 0.820 , Fig. 3C). GSH-PX activity increased with Cu treatment in the jejunum and ileum ($P < 0.05$) but not in the duodenum (Fig. 3D). MDA level was not influenced in the duodenum and jejunum, however, it decreased ($P < 0.01$) with Cu treatment in the ileum (Fig. 3E). The protein carbonyl concentration was not altered by Cu treatment (Fig. 3F).

Alpha diversity was applied to analyze the complexity of species diversity per sample. Chao 1 diversity analysis showed that Cu treatment increased ($P < 0.05$) the alpha diversity of cecal microbiota (Fig. 4A). The Shannon index in Cu-hens showed an increasing tendency (Fig. 4B). Furthermore, the rank abundance curve was longer on the X-axis for Cu-administered laying hens, thus, verifying the increased microbiota diversity in Cu-hens (Fig. 4C). Consistent with the alpha diversity, the microbial composition increased significantly at order, family, and species levels in Cu-hens, compared with the control ($P < 0.05$). The Cu-hens had 36, 53, and 17 more genera at order, family, and species levels than the control hens ($P < 0.05$, Fig. 4D). The cecal microbiota was characterized by 18 major phyla, in which *Firmicutes* and *Bacteroides* were the most predominant phyla. Although there was no significant difference in the abundance of major phyla between the Cu-treated and control groups, in Cu-hens, the

respective abundance of phylum *Elusimicrobia* increased and phylum *Fusobacteria* decreased (0.12% and 4.63%) compared with the control (0.03% and 8.84%, Fig. 4E). At the lower taxonomic levels, the genera, *Methanocorpusculum*, *Paludibacter*, *Elusimicrobium*, *Gracilibacter*, and *Syntrophomonas* (0.05%, 0.60%, 0.06%, 0.0008%, and 0.001%), and the family, Peptostreptococcaceae (0.003%) were enriched in the cecal content of Cu-hens (0.17%, 1.24%, 0.11%, 0.0021%, 0.004% and 0.076%, Fig. 4F). Cu treatment respectively increased *Bacteroidales*, *Clostridiales*, and *Alphaproteobacteria* (14.61%, 6.90%, and 2.30%) and decreased *Fucobacterium*, *Anaerobiospirillum*, and *Campylobacter* (4.62%, 1.86%, and 0.13%) at different levels of relative abundance (over 0.5%)(Fig. 4G), compared with those in control group (11.07%, 6.38%, 2.10%, 8.84%, 6.03%, and 0.49%). The samples in the control group did not form a distinct cluster, but they were essentially separated from those in the Cu treatment in the PCoA plot (Fig. 4H), indicating that Cu treatment had a potential effect on cecal microbiota structure.

At 24 h time point after inoculation treatment, the *E. coli* content was significantly higher ($P < 0.001$) in the inoculation groups, and there was no difference between the Cu and normal waterline groups (Fig. 5A). At 48 and 72 h after *E. coli* treatment, the *E. coli* content significantly decreased with Cu treatment ($P < 0.001$) compared to the control (Fig. 5A). In the Cu treatment, the *S. enteritidis* content was significantly lower ($P < 0.01$) than that of the control 3 d after inoculation (Fig. 5B).

Discussion

Copper is an essential mineral element for chickens. In broilers, copper promotes growth performance. Recently, based on meta-analysis, it was concluded that higher copper content is beneficial for the production performance of broilers (Feng *et al.* 2020).

In laying hen, copper deficiency results in anemia and eggs with abnormal sizes, shapes, and eggshells (Baumgartner *et al.* 1978). The hen-day egg production, egg weight, daily feed intake, and feed efficiency values were not affected by dietary levels of copper sulfate from 0 to 143 mg/kg in laying hens fed a corn-soybean basal diet that provided 6 mg Cu/kg diet (Christmasr and Harms, 1983). In contrast, supplementing hens with high Cu levels exceeding 250 ppm impairs the laying performance of hens by reducing feed consumption and decreasing the circulation of 17 β -oestradiol concentrations (Pearce *et al.* 1993).

In this study, the laying rate and feed intake were not significantly altered while Cu treatment increased egg weight and feed efficiency (Table 2), indicating that the Cu waterline is beneficial for the laying performance of hens. This result was contrary to the previous result of high dose of dietary Cu by Ankari *et al.* (1998), who reported that a significant reduction in egg production and a negative effect on food conversion was found when 250 mg/kg of copper was added compared to the control (no added copper). In the Cu treatment group, feed Cu intake was similar to that of the control hens, whereas Cu water intake was significantly higher than in the control hens (Table 3). The result implies that Cu source from water improves feed efficiency. Indeed, the quadratic

response of food intake and the adverse effects on food intake, egg production, and body-weight has been observed following the addition of CuSO_4 but not the addition of CuO (Jackson and Stevenson, 1981).

The Cu levels in serum and egg content were not changed in the hens of the Cu group. This result was contrary to that observed in broiler breeder hens by Berwanger *et al.* (2018), who reported that serum and yolk Cu levels increased with dietary Cu supplementation for 19 weeks in a dose-dependent manner. In laying hens, egg Cu was increased by dietary Cu supplementation at a dose of 250 mg/kg for 24 weeks (Pekel and Alp, 2011). The lack of obvious change in egg Cu appears to be related to the relatively shorter experimental period in the present study (4 weeks) and the unchanged Cu intake. The Cu content in the eggshell increased with Cu-treatment, which is thought to result from a relatively higher Cu serum level in the Cu treatment (+ 21.5%) compared with the control (Table 3). Whether the result positively influences eggshell quality remains to be investigated further.

In the present study, though the fecal Cu concentration was higher in the Cu treatment, the total fecal Cu excretion did not change, indicating that the increased Cu intake via waterline had no obvious influence on Cu excretion and, in turn, can affect the environment. The excreta Cu was significantly increased with dietary Cu supplementation (Pekel and Alp, 2011).

Cu is an integral part of CuZn-SOD, which is involved in the maintenance of redox balance, and catalyzes the formation of highly reactive hydroxyl radicals when in its

'free' form. Dietary Cu ranging from 0.5 mg/kg to 20 mg/kg primarily influences antioxidant capacity in rats (Roughead *et al.* 1999). In broiler chickens, dietary Cu level is associated with the antioxidant capacity (Song *et al.* 2011). In the present study, T-SOD and CuZn-SOD activities and T-AOC did not significantly change, suggesting that the Cu waterline had no obvious influence on the antioxidant ability of hens (Fig. 2). The unchanged serum Cu level may be a possible explanation for the unchanged antioxidant capacity. Similarly, dietary Cu levels from 5 to 15 mg/kg did not influence plasma Cu and liver CuZn-SOD activity (Koh *et al.* 1996). As a biomarker of lipid peroxidation, MDA is increased by stress (Huang *et al.* 2015; Gao *et al.* 2010). In the ileum, the reduced MDA concentration should result from the elevated CuZn-SOD and GSH-PX activities (Fig. 3). Collectively, the present result implies that the Cu waterline has no unexpected effect on the antioxidant capacity.

Gut microbiota plays a vital role in maintaining normal gastrointestinal and normal digestion of nutrients (Gong *et al.* 2021). The richness and diversity of microbial communities can be reflected by single sample diversity (alpha diversity), including two marker indicators, Chao 1 estimator and Shannon index. Chao 1 is indicative of species richness, while the Shannon index represents microbial population diversity. In this study, copper waterline application increased the Chao 1 and Shannon indexes to a certain extent (Fig. 4), indicating that copper waterline application increased the microbial alpha diversity of cecal content. This result implies a novel finding since previous studies on dietary organic copper or inorganic copper addition emphasized the

antimicrobial effect of dietary Cu (Fuller *et al.* 1960), but did not fully investigate the effects of copper on microbial diversity in the hindgut of laying hens.

A similar finding was observed in the OTU levels, which are classified via phylogenetic or population genetics to reflect the abundance of bacteria and genera in the sample. The application of copper waterline significantly increased the number of OTUs at the order, family, and species levels compared to the control ($P < 0.05$) but not at the phylum, class, and genus levels, indicating that copper waterline increases the abundance of bacteria at the order, family, and species level (Fig. 4D). The result suggests that hens with copper waterline have richer microbial abundance with an increased proportion of *Methanocorpusculum*, *Paludibacter*, *Elusimicrobium*, *Gracilibacter*, *Syntrophomonas*, and decreased proportion of *Fucobacterium*, *Anaerobiospirillum*, and *Campylobacter* (Fig. 4). The increased proportion of *Methanocorpusculum* was consistent with the previous work by Wu *et al.* (2020), who reported that it was one of the dominant methanogens that encoded various antioxidant enzymes. The decreased abundance of *Anaerobiospirillum* and *Campylobacter*, two potential pathogenic bacteria in poultry (Zhao *et al.* 2019; Sahin *et al.* 2015), indicate that copper waterline is beneficial for intestinal health. The enhancement of microbial abundance is important in maintaining intestinal stability, protecting the host against certain pathogenic and zoonotic organisms (Nava *et al.* 2005), stimulating immune responses (Mead, 2000), and consequently improving overall animal health. Therefore, the changes in the microbial composition, especially the increase in the relative

abundance of cecal bacteria, could be responsible for the improvement in egg quality (egg weight) in our study. It has been reported that the alteration in the microbiota, such as *Sphaerochaeta* and *Enorma*, may affect the metabolism of laying hens to improve egg quality, which is in agreement with our findings (Gong *et al.* 2021).

The antibacterial effects of copper waterline might be partly responsible for the changes in the microbial composition of hens. *E. coli* and *Salmonella* are two main pathogenic microorganisms that lead to foodborne illnesses transmitted from poultry products to humans (Pontin *et al.* 2021). When existing water in Cu^{2+} is released from the copper waterline and enters the bacterial cytomembrane, it breaks the enzyme systems and kills the bacteria (Gregor *et al.*, 2011). Similarly, antibacterial activity has been reported for *S. enteritidis* as it inhibits biofilm formation (Pontin *et al.* 2021). In this study, the inhibitory effect of copper treatment on *Salmonella* was observed simultaneously with the increased alpha diversity of bacteria in cecum content, suggesting the selectivity of Cu on microorganisms. Xia *et al.* (2004) found that Cu administration could significantly reduce the total pathogenic organism in the gut and have a positive effect on weight gain. Forouzandeh *et al.* (2021) and Villagómez-Estrada *et al.* (2020) reported that the supplementation of Cu could modulate bacterial community by decreasing the abundance of Streptococcaceae, a major pathogen that causes many diseases (Qiao *et al.*, 2014). In contrast, Forouzandeh *et al.* (2021) reported an increase in Clostridiaceae, Peptostreptococcaceae, and Enterococcaceae abundance at the family level in CuSO_4 treated chicken. Collectively,

the increased microbial diversity and inhibited bacterial colonization should be responsible at least partially for the improved feed efficiency of hens fed with Cu waterline.

In conclusion, the result indicates that copper waterline application is beneficial for the performance and gut health of aged laying hens. In addition, it suggests that copper waterline can inhibit the colonization of pathogenic microorganism such as *E. coli* and *Salmonella* and facilitate the enrichment of cecal microbiota diversity, which improves the performance of hens.

Acknowledgments

This research was supported by grants from the National Key Research Program of China (2016YFD0500510), Modern Agro-industry Technology Research System (CARS-40), the Key Technology Research and Development Program of Shandong province (2019JZZY020602), and Shandong Provincial Natural Science Foundation (ZR2020QC180), and the Taishan Scholars Program (No. 201511023),

Conflict of Interest

The authors declare that they have no conflict of interest. We are grateful to the reviewers for their valuable comments and suggestions on the paper.

Author Contributions

Ning Ma, Mengze Song, Sheng Li, and Xiaoyan Lin conducted the experiments; Min Liu and Ning Ma processed the experimental data, performed the analysis, drafted the paper and designed the figures; Hongchao Jiao, Xiaojuan Wang, and Jingpeng Zhao

were involved in planning and supervision of the work; Hai Lin and Shuhong Sun contributed to the design and implementation of the research, as well as the funding acquirement, project supervision, paper reviewing, and editing. All authors discussed the results and commented on the paper.

Advance View Proof

References

- al Ankari A, Najib H and Al Hozab A. Yolk and serum cholesterol and production traits, as affected by incorporating a supraoptimal amount of copper in the diet of the leghorn hen. *British Poultry Science*, 39: 393-397. 1998.
- Arthur JR. The glutathione peroxidases. *Cellular and Molecular Life Sciences*, 57: 1825-1835. 2000.
- Balevi T and Coskun B. Effects of dietary copper on production and egg cholesterol content in laying hens. *British Poultry Science*, 45: 530-534. 2004.
- Banks KM, Thompson KL, Rush JK and Applegate TJ. Effects of copper source on phosphorus retention in broiler chicks and laying hens. *Poultry Science*, 83: 990-996. 2004.
- Baumgartner S, Brown DJ, Salevsky E and Leach RM. Copper deficiency in the laying hen. *Journal of Nutrition*, 108: 804-811. 1978.
- Berwanger E, Vieira SL, Angel CR, Kindlein L, Mayer AN, Ebbing MA and Lopes M. Copper requirements of broiler breeder hens. *Poultry Science*, 97: 2785-2797. 2018.
- Bortoluzzi C, Vieira BS and Applegate TJ. Influence of dietary zinc, copper, and manganese on the intestinal health of broilers under *Eimeria* challenge. *Frontiers Veterinary Science*, 7: 13. 2020.
- Buberg ML, Witsø IL, L'Abée-Lund TM and Wasteson Y. Zinc and copper reduce conjugative transfer of resistance plasmids from extended-spectrum

- beta-lactamase-producing *Escherichia coli*. Microbial Drug Resistance, 26: 842-849. 2020.
- Buchanan NP, Hott JM, Cutlip SE, Rack AL, Asamer A and Moritz JS. The effects of a natural antibiotic alternative and a natural growth promoter feed additive on broiler performance and carcass quality. Journal of Applied Poultry Research, 17: 202-210. 2008.
- Chandra D, Shrivastava HP, Mandal AB, Tyagi PK and Singh DP. Effect of feeding different levels of dietary copper on production performance, egg quality traits and egg yolk cholesterol content in CARI-Red layers. Indian Journal of Poultry Science, 46: 79-82. 2011.
- Chang LY, Slot JW, Geuze HJ and Crapo JD. Molecular immunocytochemistry of the CuZn superoxide dismutase in rat hepatocytes. Journal of Cell Biology, 107: 2169-2179. 1988.
- Lai CC, Huang WH, Askari A, Klevay LM and Chiu TH. Expression of glutathione peroxidase and catalase in copper-deficient rat liver and heart. The Journal of Nutritional Biochemistry, 6: 256-262. 1995.
- Close B, Banister K, Baumans V, Bernoth EM, Bromage N, Bunyan J, Erhardt W, Flecknell P, Gregory N, Hackbarth H, Morton D and Warwick C. Recommendations for euthanasia of experimental animals: Part 2. DGXT of the European Commission. Laboratory Animals, 31: 1-32. 1997.
- Crapo JD, Oury T, Rabouille C, Slot JW and Chang LY. Copper, zinc superoxide

- dismutase is primarily a cytosolic protein in human cells. Proceedings of the National Academy of Sciences, 89: 10405-10409. 1992.
- Dalle-Donne I, Rossi R, Giustarini D, Milzani A and Colombo R. Protein carbonyl groups as biomarkers of oxidative stress. International Journal of Clinical Chemistry, 329: 23-38. 2003.
- David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, Ling AV, Devlin AS, Varma Y, Fischbach MA, Biddinger SB, Dutton RJ and Turnbaugh PJ. Diet rapidly and reproducibly alters the human gut microbiome. Nature, 505: 559-563. 2014.
- Demirel G, Pekel AY, Alp M and Kocabağlı N. Effects of dietary supplementation of citric acid, copper, and microbial phytase on growth performance and mineral retention in broiler chickens fed a low available phosphorus diet. The Journal of Applied Poultry Research, 2010: 1-31. 2012.
- Dewulf EM, Cani PD, Claus SP, Fuentes S, Puylaert PG, Neyrinck AM, Bindels LB, de Vos WM, Gibson GR, Thissen JP and Delzenne NM. Insight into the prebiotic concept: lessons from an exploratory, double blind intervention study with inulin-type fructans in obese women. Gut, 62: 1112-1121. 2013.
- Duncan DB. Multiple range and multiple F-test. Biometrics 11: 1-42. 1955.
- Ebeid T, Eid Y, Saleh A and EI-Hamid HA. Ovarian follicular development, lipid peroxidation, antioxidative status and immune response in laying hens fed fish oil-supplemented diets to produce n-3-enriched eggs. Animal, 2: 84-91. 2007.

- Eyssen H. The mode of action of antibiotics in stimulating growth of chickens. *Journal of Experimental Medicine*, 117: 127-128. 1963.
- Fairweather-Tait and Susan J. The concept of bioavailability as it relates to iron nutrition. *Nutrition Research*, 7: 319-325. 1987.
- Feng C, Xie B, Wuren Q and Gao M. Meta-analysis of the correlation between dietary copper supply and broiler performance. *PLoS One*, 15: e0232876. 2020.
- Fields M, Lewis CG, Lure M and Antholine WE. The influence of gender on developing copper deficiency and on free radical generation of rats fed a fructose diet. *Metabolism*, 41: 989-994. 1992.
- Forouzandeh A, Blavi L, Abdelli N, Melo-Duran D, Vidal A, Rodríguez M, Monteiro ANTR, Pérez JF, Darwich L and Sola-Oriol D. Effects of dicopper oxide and copper sulfate on growth performance and gut microbiota in broilers. *Poultry Science*, 100: 101224. 2021.
- Fuller R, Newland LGM, Briggs CAE, Braude R and Mitchell KG. The normal intestinal flora of the pig. iv. the effect of dietary supplements of penicillin, chlortetracycline or copper sulphate on the faecal flora. *Journal of Applied Bacteriology*, 23: 195-205. 1960.
- Gao J, Lin H, Wang XJ, Song ZG and Jiao HC. Vitamin E supplementation alleviates the oxidative stress induced by dexamethasone treatment and improves meat quality in broiler chickens. *Poultry Science*, 89: 318-327. 2010.
- Gemma C, Vila J, Bachstetter A and Bickford P C. Oxidative stress and the aging brain:

from theory to prevention. *Brain Aging: Models, Methods, and Mechanisms*.

PubMed, Chapter 15. 2007.

Gong H, Yang Z, Celi P, Yan L, Ding X, Bai S, Zeng Q, Xu S, Su Z, Zhuo Y, Zhang K and Wang J. Effect of benzoic acid on production performance, egg quality, intestinal morphology, and cecal microbial community of laying hens. *Poultry Science*, 100: 196-205. 2021.

Gregor G, Christopher R and Marc S. Metallic copper as an antimicrobial surface. *Applied and Environmental Microbiology*, 2011: 1541-1547. 2011.

Huang C, Jiao H, Song Z, Zhao J, Wang X and Lin H. Heat stress impairs mitochondria functions and induces oxidative injury in broiler chickens. *Journal of Animal Science*, 93: 2144-2153. 2015.

Iji P A, Saki A. A and Tivey DR. Intestinal development and body growth of broiler chicks on diets supplemented with non-starch polysaccharides. *Animal Feed Science and Technology*, 89: 175-188. 2001.

Jackson N and Stevenson MH. A study of the effects of dietary added cupric oxide on the laying, domestic fowl and a comparison with the effects of hydrated copper sulphate. *British Journal of Nutrition*, 45: 99-110. 1981.

Jung T, Höhn A, Catalgol B and Grune T. Age-related differences in oxidative protein-damage in young and senescent fibroblasts. *Archives of Biochemistry and Biophysics*, 483: 127-135. 2009.

Kaya H and Macit M. Effect of inclusion of garlic (*Allium sativum*) powder at different

- levels and copper into diets of hens on performance, egg quality traits and yolk cholesterol content. *International Journal of Poultry Science*, 11: 166-171. 2012.
- Keller GA, Warner TG, Steimer KS and Hallewell RA. Cu, Zn superoxide dismutase is a peroxisomal enzyme in human fibroblasts and hepatoma cells. *Proceedings of National Academy of Sciences of the United States of America*, 88: 7381-7385. 1991.
- Klasing KC. Comparative avian nutrition. *Quarterly Review of Biology*. 1998.
- Koh TS, Peng RK and Klasing KC. Dietary copper level affects copper metabolism during lipopolysaccharide-induced immunological stress in chicks. *Poultry Science*, 75: 867-872. 1996.
- Kovatcheva-Datchary P and Arora T. Nutrition, the gut microbiome and the metabolic syndrome. *Best Practice & Research. Clinical Gastroenterology* 27: 59-72. 2013.
- Leeson S. Copper metabolism and dietary needs. *Worlds Poultry Science Journal*, 65: 353-366. 2009.
- Li WX, Chen YQ, Zhao LH, Ma QG, Zhang JY and Ji C. No copper supplementation in a corn-soybean basal diet has no adverse effects on late-phase laying hens under normal and cyclic high temperatures. *Poultry Science*, 97: 1352-1360. 2018.
- Lien TF, Chen KL, Wu CP and Lu JJ. Effects of supplemental copper and chromium on the serum and egg traits of laying hens. *British Poultry Science*, 45: 535-539. 2004.
- Mahmoud KZ and Edens FW. Influence of selenium sources on age-related and mild heat stress-related changes of blood and liver glutathione redox cycle in broiler

- chickens (*Gallus domesticus*). Comparative Biochemistry and Physiology. Part B, Biochemistry & Molecular Biology, 136: 921-934. 2003.
- Mead GC. Prospects for 'competitive exclusion' treatment to control salmonellas and other foodborne pathogens in poultry. Veterinary Journal, 159: 111-123. 2000.
- Meeravali NN and Arunachalam J. Graphite furnace atomic absorption spectrometric (GFAAS) determination of Cu, Cd, Cr, Mn, Ni and Pb in tellurium metal using precipitation and ion-exchange procedures. Fresenius Journal of Analytical Chemistry, 358: 484-488. 1997.
- Miles RD, O'Keefe SF, Henry PR, Ammerman CB and Luo XG. The effect of dietary supplementation with copper sulfate or tribasic copper chloride on broiler performance, relative copper bioavailability, and dietary prooxidant activity. Poultry Science, 77: 416-425. 1998.
- Mohanna C and Nys Y. Influence of age, sex and cross on body concentrations of trace element (zinc, iron, copper and manganese) in chickens. British Poultry Science, 39: 536-543. 1998.
- National Research Council. Nutrition requirements of poultry. Washington (DC): National Academy Press. 1994.
- Nava GM, Bielke LR, Callaway TR and Castañeda MP. Probiotic alternatives to reduce gastrointestinal infections: the poultry experience. Animal Health Research Reviews, 6: 105-118. 2005.
- Olukosi OA, van Kuijk SJA and Han Y. Sulfate and hydroxychloride trace minerals in

- poultry diets-comparative effects on egg production and quality in laying hens, and growth performance and oxidative stress response in broilers. *Poultry Science*, 98: 4961-4971. 2019.
- Oteiza PI, Olin KL, Fraga CG and Keen CL. Zinc deficiency causes oxidative damage to proteins, lipids and DNA in rat testes. *Journal of Nutrition*, 125: 823-829. 1995.
- Pang Y, Patterson JA and Applegate TJ. The influence of copper concentration and source on ileal microbiota. *Poultry Science*, 88: 586-592. 2009.
- Pappas AC, Acamovic T, Sparks NHC, Surai PF and Mcdevitt RM. Effects of supplementing broiler breeder diets with organic selenium and polyunsaturated fatty acids on egg quality during storage. *Poultry Science*, 80: 865-874. 2005.
- Pearce J, Jackson N and Stevenson MH. The effects of dietary intake and of dietary concentration of copper sulphate on the laying domestic fowl: effects on some aspects of lipid, carbohydrate and amino acid metabolism. *British Poultry Science*, 24: 337-348. 1983.
- Pekel AY and Alp M. Effects of different dietary copper sources on laying hen performance and egg yolk cholesterol 1, 2, 3. *Journal of Applied Poultry Research*, 20: 506-513. 2011.
- Pekel AY, Demirel G, Alp M and Kocabagli N. Influence of different dietary copper sources on eggshell quality and phosphorus retention in laying hens. *Journal of Applied Poultry Research*, 21: 460-466. 2012.
- Peña MM, Lee J and Thiele DJ. A delicate balance: homeostatic control of copper

- uptake and distribution. *Journal of Nutrition*, 129: 1251-1260. 1999.
- Ponukalina EV, Afanas'eva GA, Chesnokova NP and Kirichuk VF. Lipid peroxidation and the antioxidant blood system in dynamics of pestilential and choleraic intoxication. *Patologicheskaya Fiziologiya I Eksperimental'naya Terapiya*, 9: 17-18. 2001.
- Pontin KP, Borges KA, Furian TQ, Carvalho D and Nascimento V. Antimicrobial activity of copper surfaces against biofilm formation by *Salmonella enteritidis* and its potential application in the poultry industry. *Food Microbiology*, 94: 103645. 2021.
- Prohaska JR and Gybina AA. Intracellular copper transport in mammals. *Journal of Nutrition*, 134: 1003-1006. 2004.
- Qiao Y, Sun J, Xie Z, Shi Y and Le G. Propensity to high-fat diet-induced obesity in mice is associated with the indigenous opportunistic bacteria on the interior of Peyer's patches. *Journal of Clinical Biochemistry and Nutrition*, 55: 120-128. 2014.
- Roughead ZK, Johnson LK and Hunt JR. Dietary copper primarily affects antioxidant capacity and dietary iron mainly affects iron status in a surface response study of female rats fed varying concentrations of iron, zinc and copper. *Journal of Nutrition*, 129: 1368-1376. 1999.
- Russell SM. The effect of an acidic, copper sulfate-based commercial sanitizer on indicator, pathogenic, and spoilage bacteria associated with broiler chicken

- carcasses when applied at various intervention points during poultry processing. *Poultry Science*, 87: 1435-1440. 2008.
- Sahin O, Kassem II, Shen Z, Lin J and Zhang Q. *Campylobacter* in poultry: ecology and potential interventions. *Avian Diseases*, 59: 185-200. 2015.
- Skrivan M, Skrivanová V and Marounek M. Effect of various copper supplements to feed of laying hens on Cu content in eggs, liver, excreta, soil, and herbage. *Archives of Environmental Contamination & Toxicology*, 50: 280-283. 2006.
- Skrivan M, Skrivanová V, Marounek M, Tůmová E and Wolf J. Influence of dietary fat source and copper supplementation on broiler performance, fatty acid profile of meat and depot fat, and on cholesterol content in meat. *British Poultry Science*, 41: 608-614. 2000.
- Song Z, Zhao T, Liu L, Jiao H and Lin H. Effect of copper on antioxidant ability and nutrient metabolism in broiler chickens stimulated by lipopolysaccharides. *Archives of Animal Nutrition*, 65: 366-375. 2011.
- Sturtz LA, Diekert K, Jensen LT, Lill R and Culotta VC. A fraction of yeast Cu,Zn-superoxide dismutase and its metallochaperone, CCS, localize to the intermembrane space of mitochondria. A physiological role for SOD1 in guarding against mitochondrial oxidative damage. *Journal of Biological Chemistry*, 276: 38084-38089. 2001.
- Toplan S, Dariyerli N, Özçelik D and Akyolcu MC. The effects of copper application on oxidative and antioxidant systems in rats. *Trace Elements & Electrolytes*, 22:

178-181. 2005.

Villagómez-Estrada S, Pérez L, Darwich L, Vidal A, van Kuijk S, Melo-Durán D and Solà-Oriol D. Effects of copper and zinc sources and inclusion levels of copper on weanling pig performance and intestinal microbiota. *Journal of Animal Science*, 98: 1-15. 2020.

Wu Z, Nguyen D, Lam TYC, Zhuang H, Shrestha S, Raskin L, Khanal SK and Lee PH. Synergistic association between cytochrome bd-encoded Proteiniphilum and reactive oxygen species (ROS)-scavenging methanogens in microaerobic-anaerobic digestion of lignocellulosic biomass. *Water Research*, 190: 116721. 2021.

Xia MS, Hu CH and Xu ZR. Effects of copper-bearing montmorillonite on growth performance, digestive enzyme activities, and intestinal microflora and morphology of male broilers. *Poultry Science*, 83: 1868-1875. 2004.

Yagi K. A simple fluorometric assay for lipoperoxide in blood plasma. *Biochemical Medicine*, 15: 212-216. 1976.

Yang XJ, Sun XX, Li CY, Wu XH and Yao JH. Effects of copper, iron, zinc, and manganese supplementation in a corn and soybean meal diet on the growth performance, meat quality, and immune responses of broiler chickens. *The Journal of Applied Poultry Research*, 20: 263-271. 2011.

Yin YL, Huang RL, Libao-Mercado AJ, Jeaurond EA, De Lange CFM and Rademacher M. Effect of including purified jack bean lectin in casein or hydrolysed

- casein-based diets on apparent and true ileal amino acid digestibility in the growing pig. *Animal Science*, 79: 283-291. 2004.
- Yokode M, Kita T, Kikawa Y, Ogorochi T, Narumiya S and Kawai C. Stimulated arachidonate metabolism during foam cell transformation of mouse peritoneal macrophages with oxidized low-density lipoprotein. *Journal of Clinical Investigation*, 81: 720-729. 1988.
- Roughead ZK, Johnson LK and Hunt JR. Dietary copper primarily affects antioxidant capacity and dietary iron mainly affects iron status in a surface response study of female rats fed varying concentrations of iron, zinc and copper. *Journal of Nutrition*, 129: 1368-1376. 1999.
- Zhang SSZ, Noordin MM, Rahman SOA and Haron J. Effects of copper overload on hepatic lipid peroxidation and antioxidant defense in rats. *Veterinary and Human Toxicology*, 42: 261-264. 2000.
- Zhao Y, Li X, Sun S, Chen L, Jin J, Liu S, Song X, Wu C and Lu L. Protective role of dryland rearing on netting floors against mortality through gut microbiota-associated immune performance in shaoxing ducks. *Poultry Science*, 98: 4530-4538. 2019.
- Zhou X, Wu X, Yin Y, Zhang C and He L. Preventive oral supplementation with glutamine and arginine has beneficial effects on the intestinal mucosa and inflammatory cytokines in endotoxemic rats. *Amino Acids*, 43: 813-821. 2012.

Table 1 Ingredient and nutrition composition of the basal diet

Ingredient	Content %
Corn	59.65
Soybean meal	22.91
Wheat bran	5.00
Soya-bean oil	1.14
Limestone powder, ground	8.93
Calcium hydrophosphate	1.53
Iodized salt	0.35
Lysine (99%)	0.04
Methionine (98%)	0.10
Choline chloride (50%)	0.10
Vitamin mix ¹	0.05
Mineral mix ²	0.20
Nutrition level ³	
ME, kcal/kg	2650
Crude protein, %	16.5
Calcium, %	3.54
Available phosphorus, %	0.40
Crude fiber, %	2.6
Cu mg/kg ⁴	27.16

¹Vitamin premix provided (per kilogram of diet): Vitamin A (from retinyl acetate), 5,000 IU; vitamin B1, 345 mg; vitamin B2, 1,375 mg; Vitamin D3, 1000 IU; Vitamin E, 6,600 mg; Vitamin K, 458 mg; Pantothenic acid, 1,340 mg; Nicotinic acid, 5,658 mg; Pyridoxine, 673 mg; Biotin, 550 mg; Folic acid, 122 mg; Cobalamin, 442 mg.

²Mineral premix provided (per kilogram of diet): MnSO₄, 13.84 g; ZnSO₄, 12.75 g; FeSO₄, 9.17 g; Na, 33.56 mg; KI, 113.48 mg.

³ Calculated values.

⁴ Measured value.

Table 2 Effect of copper waterline on the laying performance of hens

	Control	Cu	P-Value
Egg production, %	74.4 ± 2.15	75.6 ± 2.59	0.741
Egg weight, g	66.0 ± 0.37 ^b	68.3 ± 0.67 ^a	0.039
Feed intake, g/hen/day	113.2 ± 0.4	107.5 ± 2.7	0.103
Feed efficiency	2.30 ± 0.03 ^b	2.08 ± 0.04 ^a	0.003
Water intake, mL/hen/day	278.3 ± 48.3	313.7 ± 42.1	0.610
Body weight, g	2.34 ± 0.04	2.32 ± 0.04	0.829

Data were presented as Mean ± SEM (n = 3);

^{a, b} Means within the same line with different superscript differ significantly (P < 0.05).

Table 3 Effect of copper waterline on Cu intake and Cu content in serum and egg

	Control	Cu	P-Value
Cu intake			
Cu content in diet, mg/kg	27.16	27.16	
Dietary Cu intake, mg/hen/d	3.07 ± 0.01	2.92 ± 0.07	0.103
Water Cu content, mg/L	0.043 ± 0.02 ^b	0.373 ± 0.06 ^a	0.008
Cu intake from water, mg/hen/d	0.012 ± 0.002 ^b	0.117 ± 0.016 ^a	0.003
Total Cu intake, mg/d	3.09 ± 0.01	3.04 ± 0.06	0.536
Serum Cu level, µg/mL	1.16 ± 0.06	1.41 ± 0.16	0.178
Cu level in egg content, µg/g	2.32 ± 0.76	4.23 ± 0.14	0.235
Cu level in eggshell, µg/g	7.13 ± 0.26 ^b	8.08 ± 0.27 ^a	0.019

Values are presented as Mean ± SEM (n = 6).

^{a, b} Means within the same line with different superscripts differ significantly (P < 0.05).

Figure captions

Fig. 1 Effect of copper waterline on copper content in excreta of laying hens. (A) fecal Cu concentration (mg/kg), and (B) average daily fecal Cu excretion (mg/d/bird). Data are shown as Mean $\pm$ SEM (n = 3); *P < 0.05, #0.05 < P < 0.1.

Fig. 2 Effect of copper waterline on serum anti-oxidant properties. (A) T-SOD (B) CuZn-SOD and (C) T-AOC activities. Values are presented as Mean $\pm$ SD (n = 8)

Fig. 3 Effect of copper waterline on redox balance in duodenum, jejunum, and ileum mucosae. (A) CuZn-SOD, (B) T-SOD, (C) T-AOC, (D) GSH-PX, (E) Content of MDA, and (F) Content of Protein Carbonyl. Data are presented as Mean $\pm$ SD (n = 8); *P < 0.05, **0.05 < P < 0.01.

Fig. 4 Effect of copper waterline on the alpha diversity and microbial abundance of microbiota in the cecal content of laying hens. (A) Chao 1, (B) Shannon index, (C) Rank abundance curve, (D) OTU, (E) Phylum-level relative abundance (%) represented as a stacked graph, (F) The top 20 changes in genus-level relative abundance (%), (G) genus-level relative abundance (> 0.5%) represented as a stacked graph, and (H) PCoA analysis based on unweighted UniFrac distance metric matrices for each individual hens among groups. Data are shown as Mean $\pm$ SD (n = 4); *P < 0.05; **0.05 < P < 0.1.

Fig. 5 The antibacterial effect of copper waterline on (A) *E. coli* (n = 3), and (B) *S. enteritidis* (n = 5). Data are shown as Mean $\pm$ SD; *P < 0.05; **0.05 < P < 0.01; ***P < 0.0001.

Fig.1

Fig. 2

Fig. 3

Fig. 4

Fig. 5