1-O-11

Oral Sessions

Significance of SGLT2 in glucagon secretion from α -TC cells

Licht Miyamoto¹, Suguru Nakayama¹, Aya Hatano¹, Mana Hattori¹, Haruka Inoue¹,
Yasumasa Ikeda², Koichiro Tsuchiya¹

¹Dept. Medical Pharmacology, Inst. of Biomedical Sciences, Tokushima University, ²Dept. Pharmacology, Inst. of Biomedical Sciences, Tokushima University

SGLT2 inhibitors lower blood glucose levels by preventing renal glucose reuptake, but they often cause hyperglucagonemia. Recent studies suggested its expression and potential role as a glucagon suppressor in pancreatic alpha cells, though it has been under debate. Thus, we conducted functional analyses of SGLT2 in a typical model of pancreatic alpha cell, α -TC cells to unveil roles of SGLT2 in the glucagon secretion. Glucagon secretion as well as intracellular ATP level decreased in response to glucose deprivation. SGLT2 inhibitors reduced glucose uptake, but glucagon secretion nor ATP level was affected. An inhibitor of KATP channel increased glucagon secretion without changing ATP level. Therefore, glucose starvation should not facilitate but mitigate glucagon secretion in α -TC cells possibly by raising AMP/ATP ratio which mitigates membrane potential through KATP channel. We also found SGLT2-mediated glucose uptake in α -TC cells. Nevertheless, the glucose influx is supposed to be too small to take effects on ATP level, and SGLT2 inhibitors should not directly alter glucagon secretion. Glucose starvation-induced glucagon secretion may require interaction among different types of the cells in islets.